



Invariant Measures for Uncountable Random Interval Homeomorphisms

Janusz Morawiec¹ · Tomasz Szarek²

Received: 9 December 2022 / Accepted: 1 June 2023
© The Author(s) 2023

Abstract

A necessary and sufficient condition for the iterated function system $\{f(\cdot, \omega) \mid \omega \in \Omega\}$ with probability P to have exactly one invariant measure μ_* with $\mu_*((0, 1)) = 1$ is given. The main novelty lies in the fact that we only require the transformations $f(\cdot, \omega)$ to be increasing homeomorphisms, without any smoothness condition, neither we impose conditions on the cardinality of Ω . In particular, positive Lyapunov exponents conditions are replaced with the existence of solutions to some functional inequalities. The stability and strong law of large numbers of the considered system are also proven.

Keywords Iterated function systems · Invariant measures · Stability · Strong law of large numbers · Functional equations

Mathematics Subject Classification 37C40 · 39B12 · 47B80 · 39B22

1 Introduction

In this note we study random one-dimensional systems consisting of increasing homeomorphisms defined on the interval $[0, 1]$. These systems can be looked upon from various points of view. Here we are interested in the existence of a unique invariant measure on $(0, 1)$ (unique ergodicity). Observe that existence of such a measure may not be obtained by the Krylov–Bogolubov theorem since we are looking for a measure on a non-compact interval. On the other hand, every convex combination of the Dirac

✉ Janusz Morawiec
janusz.morawiec@us.edu.pl

Tomasz Szarek
tszarek@impan.pl

¹ Institute of Mathematics, University of Silesia, Bankowa 14, 40-007 Katowice, Poland

² Faculty of Physics and Applied Mathematics, Gdańsk University of Technology, Gabriela Narutowicza 11/12, 80-233 Gdańsk, Poland

measures δ_0 and δ_1 is invariant, therefore the aforementioned theorem, applied to the closed interval $[0, 1]$, does not carry information about the problem we are interested in.

Alsedá and Misiurewicz studied similar problem for some function systems consisting of piecewise linear homeomorphisms (see [2]). More general iterated function systems were considered by Gharaei and Homburg in [14]. Recently Malicet obtained unique ergodicity as a consequence of the contraction principle for time homogeneous random walks on the topological group of homeomorphisms defined on the circle and interval (see [17]). In turn, Zdunik and the second author [24] defined a class of iterated function systems (the so-called admissible iterated function systems) satisfying unique ergodicity on $(0, 1)$. All the results were formulated for the systems consisting of finitely many transformations.

Here we give a necessary and sufficient condition for the existence and uniqueness of an invariant measure for more general IFS's with probabilities. Our condition is of the same type as that given in [22]. In fact, this condition is expressed in the language of functional equations. Also the proof of uniqueness is based upon some results on existence of solutions of some functional equations.

The second aim of this paper is to study properties of the considered iterated function system with probabilities such as properties of supports of its invariant measures, asymptotic stability, and strong law of large numbers.

2 Preliminaries

Fix a probability space (Ω, \mathcal{A}, P) and a function $f: [0, 1] \times \Omega \rightarrow [0, 1]$ such that:

- (H₁) for every $x \in [0, 1]$ the function $f(x, \cdot): \Omega \rightarrow [0, 1]$ is \mathcal{A} -measurable;
- (H₂) for every $\omega \in \Omega$ the function $f_\omega: [0, 1] \rightarrow [0, 1]$ defined by $f_\omega(x) = f(x, \omega)$ is an increasing homeomorphism of $[0, 1]$ onto itself.

The family $\mathcal{F} = \{f_\omega \mid \omega \in \Omega\}$ forms an *iterated function system* (IFS for short) and the pair (\mathcal{F}, P) , in which we are interested in this paper, forms an *iterated function system with probabilities* (IFSP for short). Many authors have widely studied such systems (see [8] and the references therein). It is worth mentioning here that we do not make any assumptions on contractivity or at least local contractivity on average of the system as authors usually do (see e.g. [3, 19, 20]). The contractivity, for instance, does not hold for homeomorphisms. In our setting we do not assume smoothness of homeomorphisms f_ω .

Note that f is a *Carathéodory function* (i.e., continuous with respect to the first variable and \mathcal{A} -measurable with respect to the second one), and hence it is measurable with respect to σ -algebra $\mathcal{B} \otimes \mathcal{A}$, where \mathcal{B} denotes the σ -algebra of all Borel subsets of $[0, 1]$ (see [1, Lemma 4.51]). Therefore, f is also a *random-valued function* (rv-function for short), which *iterates* were introduced independently in [5] and [10] for different needs, as follows

$$f^1(x, \omega) = f(x, \omega_1) \quad \text{and} \quad f^{n+1}(x, \omega) = f(f^n(x, \omega), \omega_{n+1})$$



for all $x \in [0, 1]$, $n \in \mathbb{N}$ and $\omega = (\omega_1, \omega_2, \dots) \in \Omega^\infty$; here Ω^∞ is defined as the one-sided shift space on Ω , which plays an essential role in topological and symbolic dynamics (for more details see [23]). Observe that for every $n \in \mathbb{N}$ the iterate $f^n : [0, 1] \times \Omega^\infty \rightarrow [0, 1]$ is again an rv-function, but now on the product probability space $(\Omega^\infty, \mathcal{A}^\infty, P^\infty)$. More precisely, the iterate f^n is measurable with respect to the product σ -algebra $\mathcal{B} \otimes \mathcal{A}_n$, where \mathcal{A}_n denotes the σ -algebra of all sets of the form

$$\{(\omega_1, \omega_2, \dots) \in \Omega^\infty \mid (\omega_1, \dots, \omega_n) \in A\}$$

with A from the product σ -algebra \mathcal{A}^n .

Fix $B \in \mathcal{B}$. Since f is an rv-function, it follows that $f^{-1}(B) \in \mathcal{B} \otimes \mathcal{A}$. Hence the Ω -section $f^{-1}(B)_\omega$ of the set $f^{-1}(B)$, determined by $\omega \in \Omega$, is Borel and

$$f^{-1}(B)_\omega = \{x \in [0, 1] \mid f(x, \omega) \in B\} = \{x \in [0, 1] \mid f_\omega(x) \in B\} = f_\omega^{-1}(B).$$

Moreover, if μ is a Borel probability measure, defined on $[0, 1]$, then the function " $\omega \mapsto \mu(f_\omega^{-1}(B))$ " is \mathcal{A} -measurable and $\int_\Omega \mu(f_\omega^{-1}(B))dP(\omega) = P \otimes \mu(f^{-1}(B))$. This justifies the following definition.

A Borel probability measure μ_* , defined on $[0, 1]$, is said to be an *invariant measure* for the IFSP (\mathcal{F}, P) , if

$$\mu_*(B) = \int_\Omega \mu_*(f_\omega^{-1}(B))dP(\omega) \tag{2.1}$$

for every Borel set $B \subset [0, 1]$. Note that μ_* is invariant measure for (\mathcal{F}, P) if and only if $\mu_*(B) = P \otimes \mu_*(f^{-1}(B))$ for every Borel set $B \subset [0, 1]$.

Denote by $B([0, 1], \mathbb{R})$ the set of all bounded functions from $[0, 1]$ to \mathbb{R} , and by $C([0, 1], \mathbb{R})$ its subset of all continuous functions. Put

$$\begin{aligned} \mathcal{I} &= \{\varphi \in B([0, 1], \mathbb{R}) \mid \varphi \text{ is increasing, } \varphi(0) = 0 \text{ and } \varphi(1) = 1\}, \\ \mathcal{D} &= \mathcal{I} \cap C([0, 1], \mathbb{R}). \end{aligned}$$

Following the idea from [22] we introduce the following conditions:

- (a) for any $x \in (0, 1)$ there exists a set $\Omega_-^x \in \mathcal{A}$ with $P(\Omega_-^x) > 0$ and such that $f_\omega(x) < x$ for every $\omega \in \Omega_-^x$ or for any $x \in (0, 1)$ there exists a set $\Omega_+^x \in \mathcal{A}$ with $P(\Omega_+^x) > 0$ and such that $x < f_\omega(x)$ for every $\omega \in \Omega_+^x$;
- (b) there exist $\rho_1, \rho_2 \in \mathcal{I}$ which are right-continuous, continuous at 1, such that $\rho_1 \leq \rho_2$ and

$$\rho_1(x) \leq \int_\Omega \rho_1(f_\omega^{-1}(x))dP(\omega) \quad \text{and} \quad \int_\Omega \rho_2(f_\omega^{-1}(x))dP(\omega) \leq \rho_2(x)$$

for every $x \in [0, 1]$.

Note that if the point $x \in (0, 1)$ is a common fixed point of all functions of the family \mathcal{F} , then the Dirac measure δ_x is an invariant measure for (\mathcal{F}, P) . In such a case, the



search for invariant measures for (\mathcal{F}, P) reduces to an independent search for invariant measures for $(\mathcal{F}|_{[0,x]}, P)$ and $(\mathcal{F}|_{[x,1]}, P)$, where families $\mathcal{F}|_{[0,x]}$ and $\mathcal{F}|_{[x,1]}$ consist of all functions from \mathcal{F} restricted to $[0, x]$ and $[x, 1]$, respectively. Condition (A) excludes this situation. In particular, the invariant measure we are looking for cannot be the Dirac measure of any point of the interval $(0, 1)$. Moreover, as we will see later that condition (A) guarantees the atomlessness of the invariant measure we are interested in, as well as its uniqueness. On the other hand, it is well known that to prove the existence of an invariant measure on $(0, 1)$, some conditions on the behaviour in the neighbourhood of 0 and 1 have to be assumed. Condition (B) is parallel with the existence of positive Lyapunov exponents of smooth functions.

From now on we fix an IFSP (\mathcal{F}, P) with $f: [0, 1] \times \Omega \rightarrow [0, 1]$ satisfying (H_1) and (H_2) , and consider a function $g: [0, 1] \times \Omega \rightarrow [0, 1]$ given by

$$g(x, \omega) = f_{\omega}^{-1}(x). \quad (2.2)$$

It is clear that $g_{\omega} = f_{\omega}^{-1}$ for every $\omega \in \Omega$, and hence g satisfies (H_2) . To see that g satisfies also (H_1) it suffices to fix $x, a \in [0, 1]$ and note that

$$\{\omega \in \Omega \mid g(x, \omega) \leq a\} = \{\omega \in \Omega \mid (\omega, a) \in f^{-1}([x, 1])\} \in \mathcal{A}.$$

Therefore, g is a Carathéodory function as well as an rv-function.

In the proofs of our results it will be more convenient to use the function g instead of f .

3 Key Observation

The following simple observation (cf. e.g. [12, 9.1.1]) is motivated by [18] and reads as follows.

Proposition 3.1 (i) *If μ_* is an invariant measure for (\mathcal{F}, P) , then the function $\varphi_*: [0, 1] \rightarrow [0, 1]$ given by*

$$\varphi_*(x) = \mu_*([0, x]) \quad (3.1)$$

is increasing, right-continuous, $\varphi_(1) = 1$ and*

$$\varphi_*(x) = \int_{\Omega} \varphi_*(g(x, \omega)) dP(\omega) \quad \text{for every } x \in [0, 1]. \quad (3.2)$$

(ii) *If $\varphi_*: [0, 1] \rightarrow [0, 1]$ is an increasing and right-continuous function satisfying (3.2) with $\varphi_*(1) = 1$, then formula (3.1) determines uniquely an invariant measure for (\mathcal{F}, P) .*

Proposition 3.1 says that we have a mutual, one-to-one, correspondence between invariant measures for (\mathcal{F}, P) and right-continuous and increasing functions satisfying (3.2) taking value 1 at 1. Using for subsets $A \subset [0, 1]$ the notation χ_A to denote



the characteristic function of A defined on $[0, 1]$, we observe that the function $\chi_{[0,1]}$ corresponds to the Dirac measure δ_0 and the function $\chi_{\{1\}}$ corresponds to the Dirac measure δ_1 .

The next observation shows that if φ_* solves (3.2), then there is a fairly large family of pair of functions ρ_1, ρ_2 satisfying (B).

Remark 3.2 Fix $\varphi_* \in \mathcal{I}$ satisfying (3.2) and assume that it is right-continuous and continuous at 1. If $\psi \in \mathcal{D}$ is convex and $\phi \in \mathcal{D}$ is concave, then $\psi \circ \varphi_*, \phi \circ \varphi_* \in \mathcal{I}$ are right-continuous, continuous at 1, and, by the Jensen inequality (see e.g. [12, 10.2.7]), we have

$$\begin{aligned} (\psi \circ \varphi_*)(x) &\leq \int_{\Omega} (\psi \circ \varphi_*)(f_{\omega}^{-1}(x)) dP(\omega) \leq \int_{\Omega} (\phi \circ \varphi_*)(f_{\omega}^{-1}(x)) dP(\omega) \\ &\leq (\phi \circ \varphi_*)(x) \end{aligned}$$

for every $x \in [0, 1]$.

Proposition 3.3 Assume that

$$\{x \in (0, 1) \mid f_{\omega}(x) = x \text{ for } P\text{-almost all } \omega \in \Omega\} = \emptyset. \tag{3.3}$$

If μ is an invariant measure for (\mathcal{F}, P) , then $\mu(\{x\}) = 0$ for every $x \in (0, 1)$.

Proof Let $\varphi_*: [0, 1] \rightarrow [0, 1]$ be the function that corresponds to μ by Proposition 3.1. It follows from [21, Lemma 1] that φ_* is continuous at every point of $(0, 1)$, or equivalently, that $\mu(\{x\}) = 0$ for every $x \in (0, 1)$. \square

Corollary 3.4 Assume that (\mathcal{F}, P) satisfies (3.3). Then the existence of an invariant measure for (\mathcal{F}, P) different from all convex combinations of Dirac measures δ_0 and δ_1 is equivalent to the existence of a continuous and increasing function $\varphi_*: [0, 1] \rightarrow [0, 1]$ satisfying (3.2) with $\varphi_*(0) = 0$ and $\varphi_*(1) = 1$.

Proof (\Rightarrow) Let μ be an invariant measure for (\mathcal{F}, P) that is not a convex combination of δ_0 and δ_1 . By assertion (i) of Proposition 3.1 there exists a right-continuous and increasing function $\varphi: [0, 1] \rightarrow [0, 1]$ satisfying (3.2) with $\varphi(1) = 1$ and such that $\varphi \neq c\chi_{[0,1]} + (1 - c)\chi_{\{1\}}$ for every $c \in [0, 1]$. Moreover, Proposition 3.3 implies that φ is continuous at every point of $(0, 1)$. Therefore, the function $\varphi_*: [0, 1] \rightarrow [0, 1]$ given by

$$\varphi_*(x) = \begin{cases} \frac{\varphi(x) - \varphi(0)}{\lim_{y \rightarrow 1^-} \varphi(y) - \varphi(0)}, & \text{if } x \in [0, 1) \\ 1, & \text{if } x = 1 \end{cases}$$

is continuous, increasing, satisfies (3.2), $\varphi_*(0) = 0$ and $\varphi_*(1) = 1$.

(\Leftarrow) It is enough to apply assertion (ii) of Proposition 3.1. \square

Remark 3.5 If (\mathcal{F}, P) satisfies (A), then (3.3) holds.



In view of Remark 3.5 we see that Corollary 3.4 says that if there is an invariant measure μ for (\mathcal{F}, P) satisfying (A) and such that $\mu((0, 1)) > 0$, then there is an extreme invariant measure μ_* for (\mathcal{F}, P) such that $\mu_*((0, 1)) = 1$; recall that an extreme invariant measure for (\mathcal{F}, P) is an invariant measure for (\mathcal{F}, P) that cannot be represented as a convex combination of other invariant measures for (\mathcal{F}, P) .

We now prove that under condition (A) the considered IFSP has at most three extreme invariant measures.

Theorem 3.6 *Assume that (\mathcal{F}, P) satisfies (A). Then there exists at most one invariant measure μ_* for (\mathcal{F}, P) with $\mu_*((0, 1)) = 1$.*

Proof According to Propositions 3.1 and 3.3, and Remark 3.5, it suffices to prove that there exists at most one continuous function $\varphi_*: [0, 1] \rightarrow [0, 1]$ satisfying (3.2) with $\varphi_*(0) = 0$ and $\varphi_*(1) = 1$.

Suppose, on the contrary, that there are two different continuous functions $\varphi_1, \varphi_2: [0, 1] \rightarrow [0, 1]$ satisfying (3.2) such that $\varphi_1(0) = \varphi_2(0) = 0$ and $\varphi_1(1) = \varphi_2(1) = 1$. Put $\varphi = \varphi_1 - \varphi_2$. Clearly, φ is a non-trivial continuous solution of equation (3.2). Put $M = \sup\{|\varphi(x)| \mid x \in [0, 1]\}$ and $S = \{x \in [0, 1] \mid |\varphi(x)| = M\}$. Obviously, $a = \min S \in (0, 1)$ and $b = \max S \in (0, 1)$. By (A) and (2.2), at least one of the following cases occurs: there exists a set Ω_-^b with $P(\Omega_-^b) > 0$ such that

$$b < g(b, \omega) \quad \text{for every } \omega \in \Omega_-^b \tag{3.4}$$

or there exists set Ω_+^a with $P(\Omega_+^a) > 0$ such that

$$g(a, \omega) < a \quad \text{for every } \omega \in \Omega_+^a. \tag{3.5}$$

Note that for every $x \in S$ we have

$$M = |\varphi(x)| \leq \int_{\Omega} |\varphi(g(x, \omega))| dP(\omega) \leq M,$$

and hence $g(x, \omega) \in S$ for P -almost all $\omega \in \Omega$. In consequence,

$$a \leq g(a, \omega) \leq g(b, \omega) \leq b$$

for P -almost all $\omega \in \Omega$, which contradicts both (3.4) and (3.5). □

We end this section with an application of results obtained up to now.

Example 3.7 Fix a real number $a > 1$, an \mathcal{A} -measurable function $c: \Omega \rightarrow (0, a)$ with $\int_{\Omega} c(\omega) dP(\omega) = 1$ that is not equal to 1 almost everywhere, and consider the family \mathcal{F} consisting of functions of the form

$$f_{\omega}(x) = \begin{cases} \frac{x}{c(\omega)} & \text{for } x \in [0, \frac{c(\omega)}{a}], \\ \frac{(a-1)x+1-c(\omega)}{a-c(\omega)} & \text{for } x \in (\frac{c(\omega)}{a}, 1]. \end{cases}$$

Clearly, (A) holds and for every $\omega \in \Omega$ we have

$$g(x, \omega) = \begin{cases} c(\omega)x & \text{for } x \in [0, \frac{1}{a}], \\ \frac{c(\omega)(1-x)+ax-1}{a-1} & \text{for } x \in (\frac{1}{a}, 1]. \end{cases}$$

Moreover, if $x \in [0, \frac{1}{a}]$, then

$$\int_{\Omega} g(x, \omega)dP(\omega) = x \int_{\Omega} c(\omega)dP(\omega) = x,$$

and if $x \in (\frac{1}{a}, 1]$, then

$$\int_{\Omega} g(x, \omega)dP(\omega) = \frac{1-x}{a-1} \int_{\Omega} c(\omega)dP(\omega) + \frac{ax-1}{a-1} = x.$$

Thus we see that the function $\text{id}_{[0,1]}$ solves equation (3.2); in particular, (B) holds with $\rho_1 = \rho_2 = \text{id}_{[0,1]}$. Therefore, Proposition 3.1 implies that the one-dimensional Lebesgue measure on $[0, 1]$ is an extreme invariant measure for the considered (\mathcal{F}, P) . Moreover, by Theorem 3.6, there is no other extreme invariant measure μ for (\mathcal{F}, P) with $\mu((0, 1)) > 0$.

4 The Case of Two Extreme Measures

Combining Proposition 3.1 with [4, Theorem 2.2] we obtain the following result.

Proposition 4.1 *Assume that the family \mathcal{F} consists of pairwise commuting functions. If*

$$\int_{\Omega} g(x, \omega)dP(\omega) \neq x \text{ for every } x \in (0, 1), \tag{4.1}$$

then δ_0 and δ_1 are the only extreme invariant measures for (\mathcal{F}, P) .

The next result is a counterpart to Proposition 4.1.

Proposition 4.2 *Assume that (\mathcal{F}, P) satisfies (A). If there exists an increasing homeomorphism $h: [0, 1] \rightarrow [0, 1]$, commuting with every member of \mathcal{F} , such that*

$$h(x) \neq x \text{ for every } x \in (0, 1), \tag{4.2}$$

then δ_0 and δ_1 are the only extreme invariant measures for (\mathcal{F}, P) .



Proof Suppose that, on the contrary, there is an invariant measure μ_* for (\mathcal{F}, P) with $\mu_*((0, 1)) = 1$. Let φ_* be the function that corresponds to μ_* , by Proposition 3.1. From Theorem 3.6, Remark 3.5 and Proposition 3.3 we conclude that φ_* is the unique increasing and continuous function satisfying equation (3.2) with $\varphi_*(0) = 0$ and $\varphi_*(1) = 1$. Since the function $\varphi = \varphi_* \circ h$ is increasing, continuous $\varphi(0) = 0$, $\varphi(1) = 1$, and for every $x \in [0, 1]$ we have

$$\begin{aligned}\varphi(x) &= \int_{\Omega} \varphi_*(g(h(x), \omega)) dP(\omega) = \int_{\Omega} \varphi_*(h(g(x, \omega))) dP(\omega) \\ &= \int_{\Omega} \varphi(g(x, \omega)) dP(\omega)\end{aligned}$$

it follows that $\varphi = \varphi_*$, i.e. $\varphi_*(h(x)) = \varphi_*(x)$ for every $x \in [0, 1]$. This jointly with the assumptions imposed on h implies that φ_* is constant on $(0, 1)$, which contradicts its continuity. \square

Note that (4.1) implies (A), but (A) does not yield (4.1). Moreover, if the family \mathcal{F} consists of pairwise commuting function, then (4.1) yields the existence of at least one member in the family \mathcal{F} , which satisfies (4.2) and commutes with every member of \mathcal{F} .

The next example shows the possible application of Propositions 4.1 and 4.2.

Example 4.3 Fix an \mathcal{A} -measurable function $\alpha: \Omega \rightarrow (0, \infty)$ and consider the family \mathcal{F} that consists of functions of the form

$$f_{\omega}(x) = x^{\alpha(\omega)}.$$

We first observe that the function $h: [0, 1] \rightarrow [0, 1]$ given by $h(x) = x^2$ is an increasing homeomorphism satisfying (4.2) that commutes with every member of \mathcal{F} . If there exists $\Omega_0 \subset \Omega$ with $P(\Omega_0) > 0$ such that $\alpha(\omega) \neq 1$ for every $\omega \in \Omega_0$, then (\mathcal{F}, P) satisfies (A), and hence δ_0 and δ_1 are the only extreme invariant measures for (\mathcal{F}, P) , by Proposition 4.2. If there is no set Ω_0 with the above property, then clearly any probability measure on $[0, 1]$ is invariant measure for (\mathcal{F}, P) .

Note that in specific situations of the considered family \mathcal{F} , Proposition 4.1 can also be applied. For example, assume that $\Omega = [0, 1]$, $\mathcal{A} = \mathcal{B}$, P is the one-dimensional Lebesgue measure on $[0, 1]$ and $\alpha(\omega) = \frac{1}{\omega + \alpha}$, where $\alpha \in (0, \infty)$ is fixed. Then for every $x \in (0, 1)$ we have

$$\int_0^1 g(x, \omega) d\omega = \int_0^1 x^{\omega + \alpha} d\omega = x^{\alpha} \frac{x - 1}{\log x},$$

and hence, $x^{\alpha} \frac{x-1}{\log x} \geq \sqrt{x} \frac{x-1}{\log x} > x$ if $\alpha \in (0, \frac{1}{2}]$, whereas $x^{\alpha} \frac{x-1}{\log x} \leq x \frac{x-1}{\log x} < x$ if $\alpha \in [1, \infty)$. Therefore, Proposition 4.1 can be applied for every $\alpha \in (0, \frac{1}{2}] \cup [1, \infty)$; however it cannot be applied if $\alpha \in (\frac{1}{2}, 1)$ because (4.1) does not hold.



5 The Case of Three Extreme Measures

We are now in a position to prove the main results of this paper. For this purpose define the operator $\mathbf{T}: \mathcal{I} \rightarrow \mathcal{I}$ by

$$\mathbf{T}\varphi(x) = \int_{\Omega} \varphi(g(x, \omega))dP(\omega)$$

for every $x \in [0, 1]$.

Theorem 5.1 *Assume that (\mathcal{F}, P) satisfies (A) and (B). Then there exists exactly one invariant measure μ_* for (\mathcal{F}, P) with $\mu_*((0, 1)) = 1$.*

Proof The uniqueness is settled by Theorem 3.6. It is only the existence that must be proved.

Since

$$\rho_1 \leq \mathbf{T}^n \rho_1 \leq \mathbf{T}^{n+1} \rho_1 \leq \mathbf{T}^{n+1} \rho_2 \leq \mathbf{T}^n \rho_2 \leq \rho_2 \quad \text{for every } n \in \mathbb{N},$$

the sequence $(\mathbf{T}^n \rho_1)_{n \in \mathbb{N}}$ pointwise converges and its limit φ is an increasing function, $\rho_1 \leq \varphi \leq \rho_2$ and, by the Monotone Convergence Theorem,

$$\varphi(x) = \int_{\Omega} \varphi(g(x, \omega))dP(\omega) \quad \text{for every } x \in [0, 1].$$

Consequently, the function $\varphi_*: [0, 1] \rightarrow [0, 1]$ defined by

$$\varphi_*(x) = \lim_{y \rightarrow x^+} \varphi(y) \quad \text{for every } x \in [0, 1), \quad \varphi_*(1) = 1,$$

is increasing, right-continuous, satisfies (3.2), and $\rho_1 \leq \varphi_* \leq \rho_2$. In particular, φ_* is continuous at 1. By Proposition 3.1 formula (3.1) determines an invariant measure for (\mathcal{F}, P) , which vanishes on $\{0, 1\}$. □

It turns out that condition (B) is not only sufficient, but also necessary for the IFS (\mathcal{F}, P) satisfying (A) to have the third extreme measure. More precisely, we have the following consequence of Theorem 5.1.

Corollary 5.2 *Assume that (\mathcal{F}, P) satisfies (A). Then there exists an invariant measure μ_* for (\mathcal{F}, P) with $\mu_*((0, 1)) = 1$ if and only if (B) holds.*

Proof (\Leftarrow) This implication follows directly from Theorem 5.1.

(\Rightarrow) Suppose that there exists an invariant measure μ_* for (\mathcal{F}, P) with $\mu_*((0, 1)) = 1$. Let φ_* be the function that corresponds to μ_* by Proposition 3.1. From Proposition 3.3, which we may apply by Remark 3.5, we see that φ_* is continuous on $(0, 1)$ and since $\mu_*(\{0\}) = \mu_*(\{1\}) = 0$ we conclude that φ_* is continuous. Hence $\varphi_* \in \mathcal{D}$. Now, it is enough to apply Remark 3.2. □



Returning to the family \mathcal{F} from Example 4.3, note that it does not satisfy (B) if (A) holds, which is an immediate consequence of Corollary 5.2 or Theorem 5.1. However, if (A) does not hold, then (B) is satisfied with any right-continuous and continuous at 1 functions $\rho_1, \rho_2 \in \mathcal{I}$.

6 Supports of Invariant Measures

Given a family $\mathcal{H} = \{h_\omega \mid \omega \in \Omega\}$ of increasing homeomorphisms of an interval $I \subset \mathbb{R}$ onto itself, we say that an interval $J \subset I$ is \mathcal{H} -invariant if $h_\omega(J) \subset J$ happens P -a.s., and we say that it is \mathcal{H} -expansive if $J \subset h_\omega(J)$ happens P -a.s.

Put

$$\mathcal{J}_1 = \{[a, b] \subset (0, 1) \mid [a, b] \text{ is a minimal in the sense of inclusion } \mathcal{F}\text{-expansive interval}\},$$

$$\mathcal{J}_2 = \{[a, b] \subsetneq [0, 1] \mid [a, b] \text{ is a maximal in the sense of inclusion } \mathcal{F}\text{-expansive interval disjoint from } \bigcup \mathcal{J}_1 \text{ with } a = 0 \text{ or } b = 1\},$$

$$\mathcal{J} = \mathcal{J}_1 \cup \mathcal{J}_2.$$

Note that (3.3) implies that the family \mathcal{J}_1 consists of pairwise disjoint non-degenerate intervals, whereas (A) yields $\mathcal{J}_1 = \emptyset$.

Theorem 6.1 *Assume that (\mathcal{F}, P) satisfies (3.3). If μ is an invariant measure for (\mathcal{F}, P) , then $\text{supp } \mu \cap (a, b) = \emptyset$ for every $[a, b] \in \mathcal{J}$.*

Proof Let μ be an invariant measure for (\mathcal{F}, P) . If μ is a convex combination of δ_0 and δ_1 , then there is nothing to prove. Therefore, we assume that $\mu((0, 1)) > 0$. Put $\mu_* = \frac{1}{\mu((0, 1))} \mu$ and let φ_* be the function that corresponds to μ_* by Proposition 3.1. Note that φ_* is continuous by Proposition 3.3.

Let $\psi: \mathbb{R} \rightarrow (0, 1)$ be an increasing homeomorphism. For every $\omega \in \Omega$ we put $h_\omega = \psi^{-1} \circ g_\omega \circ \psi$ and define a function $h: \mathbb{R} \times \Omega \rightarrow \mathbb{R}$ putting $h(x, \omega) = h_\omega(x)$. It is clear that for every $x \in \mathbb{R}$ the function $h(x, \cdot): \Omega \rightarrow \mathbb{R}$ is \mathcal{A} -measurable and for every $\omega \in \Omega$ the function h_ω is an increasing homeomorphism of \mathbb{R} onto itself. By (3.3) we have

$$\{x \in \mathbb{R} \mid h(x, \omega) = x \text{ for } P\text{-almost all } \omega \in \Omega\} = \emptyset.$$

Moreover, φ_* satisfies (3.2) if and only if $\varphi = \varphi_* \circ \psi$ satisfies

$$\varphi(x) = \int_{\Omega} \varphi(h(x, \omega)) dP(\omega) \quad \text{for every } x \in \mathbb{R}.$$

Fix $[a, b] \in \mathcal{J}_1$. Then $[\psi^{-1}(a), \psi^{-1}(b)]$ is minimal $\{h_\omega^{-1} \mid \omega \in \Omega\}$ -expansive. Applying [16, Theorem 2] we obtain $\varphi(\psi^{-1}(a)) = \varphi(\psi^{-1}(b))$, and hence $\varphi_*(a) = \varphi_*(b)$. In consequence, $\text{supp } \mu = \text{supp } \mu_* \subset [0, a] \cup [b, 1]$.



Let $[0, b] \in \mathcal{J}_2$. Then $(-\infty, \psi^{-1}(b))$ is maximal $\{h_\omega^{-1} \mid \omega \in \Omega\}$ -expansive disjoint from any $\{h_\omega^{-1} \mid \omega \in \Omega\}$ -expansive compact interval $I \subset \mathbb{R}$. Applying [16, Remark 2] we conclude that φ is constant on $(-\infty, \varphi(\psi^{-1}(b)))$, i.e., φ_* is constant on $(0, b]$. In consequence, $\text{supp } \mu = \text{supp } \mu_* \subset \{0\} \cup [b, 1]$.

If $[a, 1] \in \mathcal{J}_2$, then arguing as above we come to $\text{supp } \mu \subset [0, a] \cup \{1\}$. □

The next result gives additional information about supports of invariant measures for (\mathcal{F}, P) satisfying (A).

Theorem 6.2 *Assume that (\mathcal{F}, P) satisfies (A). If μ is an invariant measure for (\mathcal{F}, P) , then $\text{supp } \mu \cap \{0, 1\} \neq \emptyset$.*

Proof Without loss of generality we can assume that $\mu((0, 1)) = 1$. Let φ be correspond to μ by Proposition 3.1.

Put

$$a = \sup\{x \in [0, 1] \mid \varphi(x) = 0\} \quad \text{and} \quad b = \inf\{x \in [0, 1] \mid \varphi(x) = 1\}.$$

Proposition 3.3 jointly with Remark 3.5 imply $\varphi(a) = 0$ and $\varphi(b) = 1$. In particular, $a < b$. Since

$$0 = \varphi(a) = \int_{\Omega} \varphi(g(a, \omega)) dP(\omega) \quad \text{and} \quad 1 = \varphi(b) = \int_{\Omega} \varphi(g(b, \omega)) dP(\omega),$$

it follows that $\varphi(g(a, \omega)) = 0$ and $\varphi(g(b, \omega)) = 1$ for P -almost all $\omega \in \Omega$. Hence $g(a, \omega) \leq a < b \leq g(b, \omega)$ for P -almost all $\omega \in \Omega$, i.e.

$$a \leq f_\omega(a) < f_\omega(b) \leq b \quad \text{for } P - \text{almost all } \omega \in \Omega. \tag{6.1}$$

Suppose that $\text{supp } \mu \cap \{0, 1\} = \emptyset$. Then we would have $0 < a < b < 1$, which jointly with (6.1) contradicts (A). □

Remark 6.3 Let μ be an invariant measure for (\mathcal{F}, P) , $x \in (0, 1)$ and $n \in \mathbb{N}$. Then $\{\omega \in \Omega^\infty \mid f^n(x, \omega) \in (\inf \text{supp } \mu, \sup \text{supp } \mu)\} \subset \{\omega \in \Omega^\infty \mid f^{n+1}(x, \omega) \in (\inf \text{supp } \mu, \sup \text{supp } \mu)\}$ holds P^∞ -a.s.

Proof Put $a = \inf \text{supp } \mu$, $b = \sup \text{supp } \mu$. Clearly, the interval $[a, b]$ is \mathcal{F} -invariant. Therefore, if $\omega = (\omega_1, \omega_2, \dots) \in \Omega^\infty$ and $a < f^n(x, \omega) < b$, then $a \leq f(a, \omega_{n+1}) < f(f^n(x, \omega), \omega_{n+1}) = f^{n+1}(x, \omega) = f(f^n(x, \omega), \omega_{n+1}) < f(b, \omega_{n+1}) \leq b$ for P -almost all $\omega_{n+1} \in \Omega$. □

7 Stability

Using some martingale techniques developed in [13] (see also [8]), we prove the stability of (\mathcal{F}, P) . Before formulating our result, let us denote by \mathbf{P} the Markov–Feller operator that corresponds to (\mathcal{F}, P) , i.e.

$$\mathbf{P}\mu(B) = \int_{[0,1]} \left(\int_{\Omega} \chi_B(f(x, \omega)) dP(\omega) \right) d\mu(x) \tag{7.1}$$



for all $\mu \in \mathcal{M}_1$ and $B \in \mathcal{B}$; here \mathcal{M}_1 denotes the family of all probability Borel measures on $[0, 1]$.

Theorem 7.1 *Assume that (\mathcal{F}, P) satisfies (A) and (B). If μ_* is the unique invariant measure for (\mathcal{F}, P) with $\mu_*((0, 1)) = 1$, then for every $x \in (0, 1)$ we have*

$$\lim_{n \rightarrow \infty} \int_{\Omega^\infty} \psi(f^n(x, \omega)) dP^\infty(\omega) = \int_{[0,1]} \psi(y) d\mu_*(y) \tag{7.2}$$

for any $\psi \in C([0, 1], \mathbb{R})$.

Proof Following [15], for every $n \in \mathbb{N}$, we consider the operator $\sigma_n: \Omega^\infty \rightarrow \Omega^\infty$ given by

$$\sigma_n(\omega_1, \omega_2, \dots) = (\omega_n, \dots, \omega_1, \omega_{n+1}, \dots).$$

Next, for all $\omega \in \Omega^\infty$ and $x \in (0, 1)$ we define a sequence $(f_n(x, \omega))_{n \in \mathbb{N}}$ by setting

$$f_n(x, \omega) = f^n(x, \sigma_n(\omega))$$

and for any $\psi \in C([0, 1], \mathbb{R})$ we define a sequence $(\xi_n^\psi)_{n \in \mathbb{N}}$ of random variables putting

$$\xi_n^\psi(\omega) = \int_{[0,1]} \psi(f_n(y, \omega)) d\mu_*(y) \quad \text{for every } \omega \in \Omega^\infty. \tag{7.3}$$

Since μ_* is an invariant measure, we easily check that $(\xi_n^\psi)_{n \in \mathbb{N}}$ is a bounded martingale with respect to the natural filtration. From the Martingale Convergence Theorem (see [11, Section XI.14]) it follows that $(\xi_n^\psi)_{n \in \mathbb{N}}$ is convergent P^∞ -a.s. This jointly with the separability of $C([0, 1], \mathbb{R})$ implies that there exists a set $\Omega_0 \subset \Omega^\infty$ with $P^\infty(\Omega_0) = 1$ such that $(\xi_n^\psi(\omega))_{n \in \mathbb{N}}$ is convergent for any $\psi \in C([0, 1], \mathbb{R})$ and $\omega \in \Omega_0$. Therefore, by the Riesz Representation Theorem, there exists a function $\mu: \Omega_0 \rightarrow \mathcal{M}_1$ such that

$$\lim_{n \rightarrow \infty} \xi_n^\psi(\omega) = \int_{[0,1]} \psi(y) d\mu(\omega)(y) \quad \text{for every } \psi \in C([0, 1], \mathbb{R}). \tag{7.4}$$

Arguing as in the proof of [8, Theorem 2] we can easily show that there exist a set $\Omega_0^0 \subset \Omega_0$ and a function $s: \Omega_0^0 \rightarrow [0, 1]$ such that $P^\infty(\Omega_0^0) = 1$ and

$$\mu(\omega) = \delta_{s(\omega)} \quad \text{for every } \omega \in \Omega_0^0. \tag{7.5}$$

Moreover,

$$\mu_* = \int_{\Omega_0^0} \delta_{s(\omega)} dP^\infty(\omega). \tag{7.6}$$



By Theorem 6.2, we have $\text{supp } \mu_* \cap \{0, 1\} \neq \emptyset$. Assume that $0 \in \text{supp } \mu_*$ (the case where $1 \in \text{supp } \mu_*$ can be considered in the same way) and set $\gamma = \sup \text{supp } \mu_*$.

We begin proving (7.2) from the case where $x \in (0, \gamma)$.

Fix $x \in (0, \gamma)$. We first want to show that

$$\lim_{n \rightarrow \infty} f_n(x, \omega) = s(\omega) \quad \text{for every } \omega \in \Omega_0^0. \tag{7.7}$$

For this purpose we fix $\omega \in \Omega_0^0$, $\varepsilon > 0$, and let $\psi_\omega \in C([0, 1], \mathbb{R})$ be such that $0 \leq \psi_\omega \leq 1$, $\psi_\omega(s(\omega)) = 1$ and $\psi_\omega(y) = 0$ if $|y - s(\omega)| > \varepsilon$. Then making use of (7.3), (7.4) and (7.5) we obtain

$$\lim_{n \rightarrow \infty} \int_{[0,1]} \psi_\omega(f_n(y, \omega)) d\mu_*(y) = \int_{[0,1]} \psi_\omega(y) d\delta_{s(\omega)}(y) = \psi_\omega(s(\omega)) = 1.$$

This jointly with the fact that $\mu_*((0, x)) > 0$ and $\mu_*((x, \gamma)) > 0$, and the definition of ψ_ω implies that there exists $n_0 \in \mathbb{N}$ such that for any $n \geq n_0$ there are $y_n \in (0, x)$ and $z_n \in (x, 1)$ with $|f_n(y_n, \omega) - s(\omega)| \leq \varepsilon$ and $|f_n(z_n, \omega) - s(\omega)| \leq \varepsilon$. Since

$$f_n(y_n, \omega) < f_n(x, \omega) < f_n(z_n, \omega) \quad \text{for every } n \geq n_0,$$

we conclude that $|f_n(x, \omega) - s(\omega)| < \varepsilon$, which means that (7.7) holds.

According to the Alexandrov Theorem (see [7, Theorem 2.1]), to prove (7.2) it is enough to show that for any $a \in (0, 1)$ we have

$$\lim_{n \rightarrow \infty} P^\infty(\{\omega \in \Omega^\infty \mid f^n(x, \omega) < a\}) = \mu_*((0, a)).$$

To verify this condition we fix $a \in (0, 1)$ and note that

$$\begin{aligned} P^\infty(\{\omega \in \Omega^\infty \mid f^n(x, \omega) < a\}) &= P^\infty(\{\omega \in \Omega^\infty \mid f^n(x, \sigma_n(\omega)) < a\}) \\ &= P^\infty(\{\omega \in \Omega^\infty \mid f_n(x, \omega) < a\}) \end{aligned}$$

for every $n \in \mathbb{N}$. Then, applying (7.7) and (7.6), we get

$$\begin{aligned} \lim_{n \rightarrow \infty} P^\infty(\{\omega \in \Omega^\infty \mid f^n(x, \omega) < a\}) &= P^\infty(\{\omega \in \Omega_0^0 \mid s(\omega) < a\}) \\ &= \int_{\{\omega \in \Omega_0^0 \mid s(\omega) < a\}} \delta_{s(\omega)}((0, 1)) dP^\infty(\omega) \\ &= \int_{\Omega_0^0} \delta_{s(\omega)}((0, a)) dP^\infty(\omega) = \mu_*((0, a)), \end{aligned}$$

which completes the proof in case where $x \in (0, \gamma)$.

Now, we are going to show that condition (7.2) also holds for any $x \in [\gamma, 1)$.

Fix $x \in [\gamma, 1)$ and observe that to complete the proof, it suffices to show that

$$\lim_{m \rightarrow \infty} P^\infty(\{\omega \in \Omega^\infty \mid f^m(x, \omega) \in (0, \gamma)\}) = 1; \tag{7.8}$$



indeed, having this, we can choose $m \in \mathbb{N}$ such that $f^m(x, \omega) \in (0, \gamma)$ happens P^∞ -a.s., and then, keeping in mind that $f^{n+m} = f^n \circ f^m$ for every $n \in \mathbb{N}$, we are able to apply (7.2), which is valid by the first part of the proof.

To prove (7.8) we fix $\varepsilon > 0$. Observe first that, by Remark 6.3, for every $n \in \mathbb{N}$ we have

$$P^\infty(\{\omega \in \Omega^\infty \mid f^{n+1}(x, \omega) \in (0, \gamma)\}) \geq P^\infty(\{\omega \in \Omega^\infty \mid f^n(x, \omega) \in (0, \gamma)\}). \tag{7.9}$$

Further, by the Krylov–Bogolubov Theorem (see [9, Theorem 7.1]), the sequence

$$\left(\frac{1}{n} \sum_{m=0}^{n-1} \mathbf{P}^m \delta_x \right)_{n \in \mathbb{N}}$$

converges weakly to an invariant measure for (\mathcal{F}, P) . Since μ_* is the unique invariant measure for (\mathcal{F}, P) with $\mu_*((0, 1)) = 1$, it follows that the Alexandrov Theorem implies the existence of $m \in \mathbb{N}$ such that

$$\mathbf{P}^m \delta_x((0, \gamma)) > \mu_*((0, \gamma)) - \varepsilon = 1 - \varepsilon,$$

due to the fact that μ_* is atomless (see Proposition 3.3 and Remark 3.5). Therefore,

$$P^\infty(\{\omega \in \Omega^\infty \mid f^m(x, \omega) \in (0, \gamma)\}) = \mathbf{P}^m \delta_x((0, \gamma)) > 1 - \varepsilon,$$

which jointly with (7.9) yields (7.8). □

8 Strong Law of Large Numbers

From the uniqueness of an invariant measure and Breiman’s law of large numbers we may easily derive the strong law of large numbers.

Theorem 8.1 *Assume that (\mathcal{F}, P) satisfies (A) and (B). If μ_* is the unique invariant measure for (\mathcal{F}, P) with $\mu_*((0, 1)) = 1$, then for all $x \in (0, 1)$ and P^∞ -almost all $\omega \in \Omega^\infty$ we have*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \psi(f^k(x, \omega)) = \int_{[0,1]} \psi(x) d\mu_*(x) \text{ for any } \psi \in C([0, 1], \mathbb{R}).$$

Proof Fix $x \in (0, 1)$. From Breiman’s law of large numbers (see [6, Corollary 3.4]) it follows that for P^∞ -almost all $\omega \in \Omega^\infty$ the sequence

$$\left(\frac{1}{n} \sum_{m=0}^{n-1} \delta_{f^m(x, \omega)} \right)_{n \in \mathbb{N}} \tag{8.1}$$



converges weakly to an invariant measure for (\mathcal{F}, P) . To complete the proof we have to show that this invariant measure is equal to μ_* . This, however, easily follows from (7.8) (or its counterpart $\lim_{m \rightarrow \infty} P^\infty(\{\omega \in \Omega^\infty \mid f^m(x, \omega) \in (\inf \text{supp } \mu_*, 1)\}) = 1$) and the Birkhoff Ergodic Theorem applied to the dynamical system corresponding to the considered Markov process on (Ω, \mathcal{A}, P) . In fact, since

$$\lim_{m \rightarrow \infty} P^\infty(\{\omega \in \Omega^\infty \mid f^m(x, \omega) \in (\beta, \gamma)\}) = 1,$$

where $\beta = \inf \text{supp } \mu_*$ and $\gamma = \sup \text{supp } \mu_*$, with no loss of generality we may (and do) assume that $x \in (\beta, \gamma)$; note that, by Theorem 6.2, $\beta = 0$ or $\gamma = 1$. Choose $x_1, x_2 \in (\beta, \gamma)$ such that $x_1 < x < x_2$. Due to the Birkhoff Ergodic Theorem for all $i \in \{1, 2\}$, $c \in (\beta, \gamma)$ and P^∞ -almost all $\omega \in \Omega^\infty$ we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=0}^{n-1} \chi_{(0,c)}(f^m(x_i, \omega)) = \mu_*((0, c)) \in (0, 1) \tag{8.2}$$

and

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=0}^{n-1} \chi_{(c,1)}(f^m(x_i, \omega)) = \mu_*((c, 1)) = 1 - \mu_*((0, c)), \tag{8.3}$$

since μ_* is atomless. Furthermore, (H_2) yields

$$\frac{1}{n} \sum_{m=0}^{n-1} f^m(x_1, \omega) < \frac{1}{n} \sum_{m=0}^{n-1} f^m(x, \omega) < \frac{1}{n} \sum_{m=0}^{n-1} f^m(x_2, \omega),$$

which jointly with (8.2) and (8.3) implies that the weak limit of the sequence (8.1) is equal to μ_* . □

Acknowledgements The first author was supported by the University of Silesia Mathematics Department (Iterative Functional Equations and Real Analysis program). The research of the second author was supported by the Polish NCN Grant 2021/43/B/ST1/01042.

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.



References

1. Aliprantis, C.D., Border, K.C.: *Infinite Dimensional Analysis*, 3rd ed. Springer, Berlin (2006). A Hitchhiker's Guide
2. Alsedá, L., Misiurewicz, M.: Random interval homeomorphisms. *Publ. Math.* **58**, 15–36 (2014)
3. Barnsley, M.F., Demko, S.G., Elton, J.H., Geronimo, J.S.: Invariant measures for Markov processes arising from iterated function systems with place-dependent probabilities. *Ann. Inst. H. Poincaré Probab. Stat.* **24**(3), 367–394 (1988)
4. Baron, K., Jarczyk, W.: Random-valued functions and iterative functional equations. *Aequationes Math.* **67**(1–2), 140–153 (2004)
5. Baron, K., Kuczma, M.: Iteration of random-valued functions on the unit interval. *Colloq. Math.* **37**(2), 263–269 (1977)
6. Benoist, Y., Quint, J.F.: *Random Walks on Reductive Groups*. Springer, Berlin (2016)
7. Billingsley, P.: *Convergence of Probability Measures*. Wiley Series in Probability and Statistics: Probability and Statistics, 2nd ed. Wiley, New York (1999). A Wiley-Interscience Publication
8. Czudek, K., Szarek, T.: Ergodicity and central limit theorem for random interval homeomorphisms. *Isr. J. Math.* **239**, 75–98 (2020)
9. Da Prato, G.: *An Introduction to Infinite-dimensional Analysis*. Universitext. Springer, Berlin (2006). Revised and extended from the 2001 original by Da Prato
10. Diamond, P.: A stochastic functional equation. *Aequationes Math.* **15**(2–3), 225–233 (1977)
11. Doob, J.L.: *Measure Theory*. Graduate Texts in Mathematics, Vol. 143. Springer, New York (1994)
12. Dudley, R.M.: *Real Analysis and Probability*, volume 74 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge (2002). Revised reprint of the 1989 original
13. Furstenberg, H.: Boundary Theory and Stochastic Processes on Homogenous Spaces. In: *Proceedings of Symposia in Pure Mathematics*, vol. 26, pp. 193–229. American Mathematical Society, Providence (1973)
14. Gharaei, M., Homburg, A.J.: Random interval diffeomorphisms. *Discrete Contin. Dyn. Syst.* **10**, 241–272 (2017)
15. Kapica, R.: Sequences of iterates of random-valued vector functions and continuous solutions of a linear functional equation of infinite order. *Bull. Pol. Acad. Sci. Math.* **50**(4), 447–455 (2002)
16. Kochanek, T., Morawiec, J.: Probability distribution solutions of a general linear equation of infinite order. *Ann. Polon. Math.* **95**(2), 103–114 (2009)
17. Malicet, D.: Random walks on homeo(s^1). *Comm. Math. Phys.* **356**(3), 1083–1116 (2017)
18. Matkowski, J.: Remark on BV-solutions of a functional equation connected with invariant measures. *Aequationes Math.* **29**(2–3), 210–213 (1985)
19. Mihailescu, E., Urbański, M.: Random countable iterated function systems with overlaps and applications. *Adv. Math.* **298**, 726–758 (2016)
20. Mihailescu, E., Urbański, M.: Geometry of measures in random systems with complete connections. *J. Geom. Anal.* **32**(5), 162, 18 (2022)
21. Morawiec, J., Reich, L.: The set of probability distribution solutions of a linear functional equation. *Ann. Polon. Math.* **93**(3), 253–261 (2008)
22. Morawiec, J., Zürcher, T.: A new take on random interval homeomorphisms. *Fund. Math.* **257**(1), 1–17 (2022)
23. Ott, W., Tomforde, M., Willis, P.N.: *One-Sided Shift Spaces Over Infinite Alphabets*. New York Journal of Mathematics, vol. 5. NYJM Monographs. State University of New York, University at Albany, Albany (2014)
24. Szarek, T., Zdunik, A.: Attractors and invariant measures for random interval homeomorphisms. Manuscript

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

