



Full length article

Wave dispersion relations in peridynamics: Influence of kernels and similarities to nonlocal elasticity theories

Victor A. Eremeyev ^{a,b} ,* , Konstantin Naumenko ^c ^a Department of Civil and Environmental Engineering and Architecture (DICAAR), University of Cagliari, Via Marengo, 2, 09123 Cagliari, Italy^b Gdańsk University of Technology, ul. Gabriela Narutowicza 11/12, 80-233 Gdańsk, Poland^c Institute of Materials, Technologies and Mechanics (IWTM), Faculty of Mechanical Engineering (FMB), Otto von Guericke University, Universitätsplatz 2, 39106 Magdeburg, Germany

ARTICLE INFO

Keywords:

Peridynamics
Dispersion relation
Anomalous dispersion
Nonlocal models
Long-range interactions

ABSTRACT

We investigate the wave dispersion relations of an infinite elastic bar within the framework of linear bond-based peridynamics. This nonlocal integral-type model accounts for long-range interactions, which become significant at small scales and in cases of damage and fracture. Since a key element of this material model is the kernel function, we derive dispersion curves for several kernel choices. Notably, for non-singular kernels, we observe negative group velocities, indicating that peridynamics can describe materials with anomalous dispersion. By comparing one-dimensional (1D) peridynamics with the 1D nonlocal elasticity of Eringen's type, we highlight similarities between the two models in terms of dispersion behavior.

1. Introduction

Recently, there has been a significant increase in interest in nonlocal models of continua and structures. These models have found various applications in modeling long-range interactions, which are essential at small scales (see, for example, the discussions in [de Gennes \(1981\)](#), [Kröner \(1967\)](#), [Maugin \(2017\)](#)). They are also particularly relevant in the study of damage and fracture, as exemplified by the nonlocal elasticity theory proposed by [Eringen \(1972\)](#), [Eringen and Edelen \(1972\)](#) and summarized in [Eringen \(2002\)](#). The integral constitutive relations for stresses introduced in these works can, in some cases, be transformed into a differential form, commonly referred to as stress-gradient elasticity—a complementary counterpart to strain-gradient elasticity ([Maugin, 2017](#)). This approach, sometimes with modifications, has been extensively studied in the literature (see, for example, [Aifantis \(1999\)](#), [Askes and Aifantis \(2011\)](#), [Barretta, Caporale, Luciano, and Vaccaro \(2023\)](#), [Barretta, Luciano, de Sciarra, and Vaccaro \(2024\)](#), [Khodabakhshi and Reddy \(2015\)](#), [Penna \(2023\)](#) and the references therein).

In addition to Eringen-type nonlocal elasticity, another nonlocal model, known as peridynamics, has been proposed by [Silling \(2000\)](#) (see also [Madenci and Oterkus \(2013\)](#), [Silling and Lehoucq \(2010\)](#)). Peridynamics has also found applications in modeling long-range interactions at small scales and in describing damage and fracture (see, for example, [Chan and Chen \(2023\)](#), [Izadi, Das, Fantuzzi, and Trovalusci \(2024\)](#), [Naumenko, Pander, and Würkner \(2022\)](#) and the references therein). Wave phenomena, including dispersion relations, have been analyzed in several studies ([Alebrahim, Packo, Zaccariotto, & Galvanetto, 2022](#); [Bažant, Luo, Chau, & Bessa, 2016](#); [Chan & Chen, 2021](#); [Chen, Peng, Jafarzadeh, & Bobaru, 2023](#); [Gu, Zhang, Huang, & Yv, 2016](#); [Mutnuri & Gopalakrishnan, 2018](#); [Silling, Zimmermann, & Abeyaratne, 2003](#); [Wang & Huang, 2019](#); [Weckner & Abeyaratne, 2005](#); [Yang, Ma, Oterkus, Oterkus, & Naumenko, 2023](#)). In the case of state-based peridynamics, dispersion has been discussed in [Alebrahim](#)

* Corresponding author at: Department of Civil and Environmental Engineering and Architecture (DICAAR), University of Cagliari, Via Marengo, 2, 09123 Cagliari, Italy.

E-mail addresses: victor.eremeyev@unica.it (V.A. Eremeyev), konstantin.naumenko@ovgu.de (K. Naumenko).

<https://doi.org/10.1016/j.ijengsci.2025.104256>

Received 22 February 2025; Received in revised form 10 March 2025; Accepted 11 March 2025

Available online 22 March 2025

0020-7225/© 2025 The Authors.

Published by Elsevier Ltd.

This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

(2023), Butt, Timothy, and Meschke (2017). These studies have shown that dispersion behavior strongly depends on the choice of kernel functions. While previous works have typically considered one or two kernels, a comparative analysis of five kernels in the static case was presented by Naumenko, Yang, Ma, and Chen (2023). Dispersion properties play a crucial role in wave propagation and the dynamic response of materials in general (see Achenbach (1973), Whitham (1999)). Moreover, experimental data and/or theoretical estimations of dispersion can be used both for validating nonlocal models and for determining material parameters.

The aim of this paper is to analyze dispersion relations within the one-dimensional (1D) bond-based peridynamics framework, considering different kernel functions. It is worth noting that the 1D model is not only of theoretical interest but also has practical applications, for example in modeling brittle fibers such as glass fibers, which are widely used in modern engineering.

The remainder of this paper is organized as follows. In Section 2, starting from the classical case, we introduce the basic equation of motion and the dispersion relations for peridynamics. In Section 3, we derive the dispersion relations—i.e., the relations between frequency and wavenumber—for several kernel functions and present the corresponding dispersion curves. Finally, in Section 4, we compare the dispersion behavior in peridynamics with that in nonlocal elasticity.

2. Equations of motion

In the following we consider infinitesimal deformations of an infinite homogeneous elastic bar. So we introduce the 1D displacement field u as a smooth enough function of a spatial coordinate x and time t :

$$u = u(x, t).$$

2.1. Simple bar

Within the linear elasticity we have Hooke’s law

$$\sigma = E\varepsilon,$$

where σ is the stress, E is the Young modulus, and $\varepsilon \equiv u'$ is the strain. As a result, we get the equation of motion in the form of the wave equation (Achenbach, 1973)

$$\rho\ddot{u} = Eu'', \tag{1}$$

where ρ is the mass density. Hereinafter the prime stands for the derivative with respect to x , whereas the overdot denotes the derivative with respect to t .

Looking for a solution of (1) in the harmonic form

$$u = U \exp [i(kx - \omega t)], \tag{2}$$

where U is an amplitude, $i = \sqrt{-1}$ is the imaginary unit, k and ω are the wavenumber and frequency, respectively, we get the dispersion relation

$$\omega = c_\infty k, \quad c_\infty = \sqrt{\frac{E}{\rho}}, \tag{3}$$

where c_∞ is the speed of the longitudinal waves. Obviously, Eq. (1) describes non-dispersive waves.

Motivated by experimental data and comparisons with solutions obtained from 3D linear elasticity, various extensions of (1) related to dispersive waves can be found in the literature. These include Rayleigh’s and Bishop’s equations (Bishop, 1952; Strutt, 1945), the 3D Pochhammer–Chree solution presented in Royer and Dieulesaint (2000), and the reviews in Artobolevskiy, Bobrovitskiy, and Genkin (1979), Eremeyev (2025).

2.2. Bond-based peridynamics

Following Silling (2000), Silling and Lehoucq (2010) let us now consider 1D wave propagation within the linear bond-based peridynamics. Instead of (1) we have the integro-differential equation

$$\rho\ddot{u} = \frac{E}{\chi(\delta)} \int_{-\delta}^{\delta} F(\xi) [u(x + \xi, t) - u(x, t)] d\xi, \tag{4}$$

where δ is the horizon size, $F(\xi)$ is an even kernel function such that

$$F(-\xi) = F(\xi), \quad \lim_{\xi \rightarrow 0} \xi^2 F(\xi) < 0,$$

and

$$\chi(\delta) = \frac{1}{2} \int_{-\delta}^{\delta} \xi^2 F(\xi) d\xi.$$

For $\delta \rightarrow 0$ and some reasonable assumptions about $F(\xi)$ Eq. (4) could be transformed to the classical one.

Although the concept of stress does not play an essential role in peridynamics, see, e.g., comments by [Madenci and Oterkus \(2013\)](#), [Silling and Lehoucq \(2008, 2010\)](#), [Silling et al. \(2003\)](#), [Weckner and Abeyaratne \(2005\)](#). Just to have a possibility to make a connection between this model and other local and non-local theories, the effective stress could be introduced similar to [Silling et al. \(2003\)](#), [Weckner and Abeyaratne \(2005\)](#) as follows

$$\sigma(x, t) = \frac{E}{\varkappa(\delta)} \int_0^\delta \int_0^{\delta-\xi} F(\xi + \eta) [u(x + \eta, t) - u(x - \xi, t)] d\eta d\xi, \tag{5}$$

In particular, for homogeneous deformations of the form $u(x, t) = x\varepsilon(t)$ we came to Hooke's law.

Substituting (2) into Eq. (4) we came to

$$\omega^2 = c_\infty^2 C(k), \quad C(k) = \frac{1}{\varkappa} \int_{-\delta}^\delta F(\xi) [1 - \cos k\xi] d\xi. \tag{6}$$

As a result, we get the dispersion relation in the form

$$\omega = c_\infty A(k), \quad A(k) = [C(k)]^{1/2}, \tag{7}$$

see e.g. [Silling et al. \(2003\)](#), [Weckner and Abeyaratne \(2005\)](#) for more details.

Using (7) we obtain the phase c and group v velocities given by

$$c \equiv \frac{\omega}{k} = c_\infty \frac{A(k)}{k}, \quad v \equiv \frac{d\omega}{dk} = c_\infty \frac{dA(k)}{dk}. \tag{8}$$

3. Kernel functions and dispersion relations

The kernel function $F(\xi)$ determines the material response within the bond-based peridynamics. Some useful kernel functions can be found in the literature and are summarized below as in [Naumenko et al. \(2023\)](#)

$$F_1(\xi) = \begin{cases} 1, & |\xi| \leq \delta, \\ 0 & |\xi| > \delta \end{cases}, \tag{9}$$

$$F_2(\xi) = \begin{cases} 1 - |\xi|, & |\xi| \leq \delta, \\ 0 & |\xi| > \delta \end{cases}, \tag{10}$$

$$F_3(\xi) = \begin{cases} \frac{1}{|\xi|}, & |\xi| \leq \delta, \\ 0 & |\xi| > \delta \end{cases}, \tag{11}$$

$$F_4(\xi) = \begin{cases} \exp\left(-4\frac{\xi^2}{\delta^2}\right), & |\xi| \leq \delta, \\ 0 & |\xi| > \delta \end{cases}. \tag{12}$$

In particular, the uniform kernel function (9) was used by [Bažant et al. \(2016\)](#), [Silling et al. \(2003\)](#), [Yang et al. \(2023\)](#). The exponential kernel (12) was also used by [Weckner and Abeyaratne \(2005\)](#) along with three other kernels, where the corresponding dispersion curves were obtained for an infinite horizon size. In the dispersion analysis conducted by [Mikata \(2012, 2019\)](#), several kernel functions were considered, including the uniform (9), triangular (10), and exponential (12) kernels. A static analysis involving the hyperbolic kernel (11) was provided by [Yang et al. \(2023\)](#).

Some other kernel functions could be proposed such as the following

$$F_5(\xi) = \begin{cases} (\delta^2 - \xi^2)^m, & |\xi| \leq \delta, \\ 0 & |\xi| > \delta \end{cases}, \tag{13}$$

$$F_6(\xi) = \begin{cases} \sin^m \frac{\pi\xi}{\delta}, & |\xi| \leq \delta, \\ 0 & |\xi| > \delta \end{cases}, \tag{14}$$

$$F_7(\xi) = \begin{cases} \frac{1}{|\xi|^2}, & |\xi| \leq \delta, \\ 0 & |\xi| > \delta \end{cases}. \tag{15}$$

where m is an even number. Note that the kernel functions F_3 and F_5 are singular. These functions tend to infinity at $\xi \rightarrow 0$.

The introduced kernels lead to the following values of \varkappa and functions

$$D(k) \equiv \int_0^\delta F(\xi) [1 - \cos k\xi] d\xi = \frac{1}{2} \varkappa C(k)$$

used in the dispersion Eqs. (6) and (7)

$$\varkappa_1 = \frac{1}{3} \delta^3, \quad D_1(k) = \delta - \frac{\sin(k\delta)}{k},$$




Fig. 1. Normalized frequency $\bar{\omega} = \frac{\delta\omega}{c_\infty}$ versus normalized wave number $\bar{k} = k\delta$. Dashed line corresponds to (3). Curves 1, 2, ..., 7 relates to kernels F_1, F_2, \dots, F_7 , respectively.




Fig. 2. Normalized phase velocity $\bar{c} = \frac{c}{c_\infty}$ versus normalized wave number $\bar{k} = k\delta$. Dashed line corresponds to the equation $c = c_\infty$. Curves 1, 2, ..., 7 relates to kernels F_1, F_2, \dots, F_7 , respectively.

$$x_2 = \frac{1}{3} \delta^3 - \frac{1}{4} \delta^4,$$

$$x_3 = \frac{1}{2} \delta^2,$$

$$x_4 = \frac{\delta^3}{32} \left(\sqrt{\pi} \operatorname{erf}(2) - \frac{4}{e^4} \right),$$

$$x_5 = \frac{1}{3} \delta^4 - \frac{2}{5} \delta^2 + \frac{1}{7},$$

$$x_6 = \frac{1}{12} \frac{\delta^3 (2\pi^2 - 3)}{\pi^2},$$

$$x_7 = \delta,$$

$$D_2(k) = \frac{1}{2} + \frac{\cos(k) - 1}{k^2},$$

$$D_3(k) = \gamma + \ln(k\delta) - \operatorname{Ci}(k\delta),$$

$$D_4(k),$$

$$D_5(k) = \frac{8}{15} \delta^5 + \frac{\sin(k\delta) \delta^2 k^2 + 3k \cos(k\delta) \delta - 3 \sin(k\delta)}{k^5},$$

$$D_6(k) = \frac{4\pi^2 k \delta - \delta^3 k^3 - 4\pi^2 \sin(k\delta)}{2k(-k^2 \delta^2 + 4\pi^2)},$$

$$D_7(k) = \frac{\operatorname{Si}(k\delta) k \delta + \cos(k\delta) - 1}{\delta},$$

where $\gamma \approx 0.57721$ is Euler's constant, Ci and Si is the cosine and sine integral, see Abramowitz and Stegun (1972). We have used the value $m = 2$, and we omit the awkward expression for $D_4(k)$.

The dispersion curves are shown in Fig. 1. Here $\bar{\omega} = \frac{\delta\omega}{c_\infty}$, $\bar{k} = k\delta$. The dashed line corresponds to the classical elasticity (3), while the i th curve corresponds to the F_i kernel, $i = 1, \dots, 7$. At $k = 0$ all curves have the same tangent given by (3).




Fig. 3. Normalized group velocity $\bar{v} = \frac{v}{c_\infty}$ versus normalized wave number $\bar{k} = k\delta$. Dashed line corresponds to the equation $c = c_\infty$. Curves 1, 2, ..., 7 relates to kernels F_1, F_2, \dots, F_7 , respectively.

In Fig. 2, we show the normalized phase velocities, $\bar{c} = \frac{c}{c_\infty}$. Clearly, for all kernel functions, $c_i < c_\infty$, meaning that waves propagate at lower velocities compared to classical elasticity. At $k = 0$, all curves have the same horizontal tangent, $c = c_\infty$.

Finally, in Fig. 3, we present the normalized group velocities, $\bar{v} = \frac{v}{c_\infty}$. Except for singular kernels, all other curves intersect the horizontal axis, meaning that for certain wavelength ranges, the group velocity becomes negative. This phenomenon is associated with so-called anomalous dispersion. For hyperbolic kernels, the group velocity remains non-negative but reaches zero at certain values of k , similar to other curves for non-singular kernels. In this case, waves with the corresponding wavelengths do not transmit energy.

4. Comparison with the nonlocal elasticity

Let us compare the dispersion relations with those derived within nonlocal elasticity. Following Eringen (2002), Eringen and Edelen (1972), in the 1D case the constitutive relation has the form

$$\sigma = E \int_{-\infty}^{\infty} K(x - \xi) \varepsilon(\xi, t) d\xi, \tag{16}$$

where E is again Young's modulus, K is an even kernel function normalized as follows

$$\int_{-\infty}^{\infty} K(\xi) d\xi = 1.$$

Now the equation of motion is given by

$$\rho \ddot{u} = \sigma' \equiv E \left[\int_{-\infty}^{\infty} K(x - \xi) \varepsilon(\xi, t) d\xi \right]'. \tag{17}$$

Looking for a solution of (17) in the harmonic form we came to the equation

$$\omega^2 = c_\infty^2 k^2 \bar{K}(k), \quad \bar{K}(k) = \int_{-\infty}^{\infty} K(\eta) e^{-ik\eta} d\eta = \int_{-\infty}^{\infty} K(\eta) \cos k\eta d\eta. \tag{18}$$

This gives us the dispersion relation

$$\omega \equiv \omega_{NL}(k) = c_\infty B(k), \quad B(k) = k \left[\bar{K}(k) \right]^{1/2}. \tag{19}$$

So the phase velocity became

$$c_{NL} = c_\infty \frac{B(k)}{k} = c_\infty \left[\bar{K}(k) \right]^{1/2}. \tag{20}$$

Note that if $c_{NL}(k) \in L^2(\mathbb{R})$ we can find the kernel function using (20) and the inverse Fourier transform by the formula

$$K(\eta) = \frac{1}{2\pi c_\infty^2} \int_{-\infty}^{\infty} c_{NL}^2(k) e^{ik\eta} dk. \quad (21)$$

Askes and Aifantis (2011) provided a detailed analysis of the dispersion relations within stress and strain gradient elasticity, where ω^2 takes the form of a rational function of k .

Introducing non-local model Eringen was motivated by the comparison of dispersion with the lattice dynamics, see Eringen (1972, 2002), Eringen and Edelen (1972). In particular, a kernel can be taken to fit the dispersion curves obtained within the lattice dynamics.

A similar idea can be applied to peridynamics. Moreover, discussing the correspondence between the equation and the dispersion relation (Whitham, 1999, pp. 367, 368) mentioned that the equation can be constructed to give any desired phase velocity and consequently any desired dispersion function.

Comparing dispersion relations (7) and (19) we can see that they coincide to each other when

$$A(k) = B(k), \quad (22)$$

or if

$$\frac{1}{x} \int_{-\delta}^{\delta} F(\xi) [1 - \cos k\xi] d\xi = k^2 \int_{-\infty}^{\infty} K(\eta) \cos k\eta d\eta. \quad (23)$$

For example, in the case of the uniform kernel function (9) Eq. (23) transforms to the determination of an inverse Fourier transform of the function

$$\bar{K}(k) = \frac{6}{\delta^2 k^2} \left[1 - \frac{\sin k\delta}{k\delta} \right].$$

This correspondence means that the same dispersion could be described using both non-local theories. Of course, this conclusion is based on the analysis of the 1D case of an infinite bar, i.e. we have not considered here spatial effects and boundary conditions. We have also assumed that all mathematical operations are meaningful.

Conclusions

Within the 1D linear bond-based peridynamics, we have obtained dispersion relations for several kernel functions and plotted the corresponding phase and group velocity curves. Examination of the group velocity shows that it can be negative, which is related to the so-called anomalous dispersion observed in some metamaterials, see e.g. Bigoni, Guenneau, Movchan, and Brun (2013), Bossart and Fleury (2023), Park and Oh (2019) and the references therein.

Notably, all considered non-singular kernels produce qualitatively similar oscillatory dispersion curves for short-wavelength waves ($k\delta \geq 5$), with a horizontal asymptote. In contrast, for long-wavelength waves ($k\delta \leq 2$), all curves nearly coincide.

As already mentioned by Mikata (2012), peridynamics could be useful for modeling materials with anomalous dispersion. Comparing peridynamics with nonlocal elasticity, we emphasize that both theories can exhibit the same dispersion behavior. In particular, depending on the chosen kernel, nonlocal elasticity may also describe anomalous dispersion.

CRedit authorship contribution statement

Victor A. Eremeyev: Writing – review & editing, Writing – original draft, Visualization, Validation, Software, Project administration, Funding acquisition, Formal analysis, Conceptualization. **Konstantin Naumenko:** Writing – review & editing, Writing – original draft, Visualization, Investigation, Funding acquisition, Formal analysis, Conceptualization.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

V.A.E. acknowledges the financial support under the National Recovery and Resilience Plan (NRRP), Mission 4, Component 2, Investment 1.1, Call for tender No. 1409 published on 14.9.2022 by the Italian Ministry of University and Research (MUR), funded by the European Union – NextGenerationEU – Project Title “A Fluid-Structure Interaction tool for the protection of Clean Energy Production sites (FSI-CEP)” – CUP F53D23009640001- Grant Assignment Decree No. P20227CSJ5 adopted on 14 September 2022 by the Italian Ministry of University and Research (MUR).

K.N. acknowledges the support of the German Research Foundation, DFG, Grant Nr. NA 342/7-1.

Data availability

No data was used for the research described in the article.

References

- Abramowitz, M., & Stegun, I. A. (1972). *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, Vol. 55. Washington: US Government printing office.
- Achenbach, J. (1973). *Wave propagation in elastic solids*. Amsterdam: North Holland.
- Aifantis, E. C. (1999). Gradient deformation models at nano, micro, and macro scales. *Journal of Engineering Materials and Technology*, 121(2), 189–202.
- Alebrahim, R. (2023). Modified wave dispersion properties in 1D and 2D state-based peridynamic media. *Computers & Mathematics with Applications*, 151, 21–35.
- Alebrahim, R., Pacco, P., Zaccariotto, M., & Galvanetto, U. (2022). Improved wave dispersion properties in 1D and 2D bond-based peridynamic media. *Computational Particle Mechanics*, 9(4), 597–614.
- Artobolevskiy, I. I., Bobrovitskiy, Y. I., & Genkin, M. D. (1979). *Introduction to acoustical dynamics of machines (in Russian)*. Moscow: Nauka.
- Askes, H., & Aifantis, E. C. (2011). Gradient elasticity in statics and dynamics: An overview of formulations, length scale identification procedures, finite element implementations and new results. *International Journal of Solids and Structures*, 48(13), 1962–1990.
- Barretta, R., Caporale, A., Luciano, R., & Vaccaro, M. S. (2023). Nonlocal gradient mechanics of nanobeams for non-smooth fields. *International Journal of Engineering Science*, 189, Article 103879.
- Barretta, R., Luciano, R., de Sciarra, F. M., & Vaccaro, M. S. (2024). Modelling issues and advances in nonlocal beams mechanics. *International Journal of Engineering Science*, 198, Article 104042.
- Bažant, Z. P., Luo, W., Chau, V. T., & Bessa, M. A. (2016). Wave dispersion and basic concepts of peridynamics compared to classical nonlocal damage models. *Journal of Applied Mechanics*, 83(11), Article 111004.
- Bigoni, D., Guenneau, S., Movchan, A. B., & Brun, M. (2013). Elastic metamaterials with inertial locally resonant structures: Application to lensing and localization. *Physical Review B—Condensed Matter and Materials Physics*, 87(17), Article 174303.
- Bishop, R. E. D. (1952). Longitudinal waves in beams. *Aeronautical Quarterly*, 3(4), 280–293.
- Bossart, A., & Fleury, R. (2023). Extreme spatial dispersion in nonlocally resonant elastic metamaterials. *Physical Review Letters*, 130(20), Article 207201.
- Butt, S. N., Timothy, J. J., & Meschke, G. (2017). Wave dispersion and propagation in state-based peridynamics. *Computational Mechanics*, 60, 725–738.
- Chan, W., & Chen, H. (2021). Peridynamic bond-associated correspondence model: Wave dispersion property. *International Journal for Numerical Methods in Engineering*, 122(18), 4848–4863.
- Chan, W., & Chen, H. (2023). Modeling material length-scale effect using the second-order peridynamic material correspondence model. *International Journal of Engineering Science*, 189, Article 103877.
- Chen, Z., Peng, X., Jafarzadeh, S., & Bobaru, F. (2023). Analytical solutions of peridynamic equations. Part II: elastic wave propagation. *International Journal of Engineering Science*, 188, Article 103866.
- de Gennes, P.-G. (1981). Some effects of long range forces on interfacial phenomena. *Journal de Physique Lettres*, 42(16), 377–379.
- Eremeyev, V. A. (2025). On dispersion relations within 1D nonlocal elasticity. In H. Altenbach, & et al. (Eds.), *Current developments in solid mechanics and their applications*. Cham: Springer.
- Eringen, A. C. (1972). Linear theory of nonlocal elasticity and dispersion of plane waves. *International Journal of Engineering Science*, 10(5), 425–435.
- Eringen, A. C. (2002). *Nonlocal continuum field theories*. New York: Springer.
- Eringen, A. C., & Edelen, D. G. B. (1972). On nonlocal elasticity. *International Journal of Engineering Science*, 10(3), 233–248.
- Gu, X., Zhang, Q., Huang, D., & Yv, Y. (2016). Wave dispersion analysis and simulation method for concrete SHPB test in peridynamics. *Engineering Fracture Mechanics*, 160, 124–137.
- Izadi, R., Das, R., Fantuzzi, N., & Trovalusci, P. (2024). Fracture properties of green nano fibrous network with random and aligned fiber distribution: A hierarchical molecular dynamics and peridynamics approach. *International Journal of Engineering Science*, 204, Article 104136.
- Khodabakhshi, P., & Reddy, J. N. (2015). A unified integro-differential nonlocal model. *International Journal of Engineering Science*, 95, 60–75.
- Kröner, E. (1967). Elasticity theory of materials with long range cohesive forces. *International Journal of Solids and Structures*, 3(5), 731–742.
- Madenci, E., & Oterkus, E. (2013). *Peridynamic theory and its applications*. Springer.
- Maugin, G. A. (2017). *Non-classical continuum mechanics: A dictionary*. Singapore: Springer Singapore.
- Mikata, Y. (2012). Analytical solutions of peristatic and peridynamic problems for a 1D infinite rod. *International Journal of Solids and Structures*, 49(21), 2887–2897.
- Mikata, Y. (2019). Linear peridynamics for isotropic and anisotropic materials. *International Journal of Solids and Structures*, 158, 116–127.
- Mutnuri, V. S., & Gopalakrishnan, S. (2018). A comparative study of wave dispersion between discrete and continuum linear bond-based peridynamics systems: 1D framework. *Mechanics Research Communications*, 94, 40–44.
- Naumenko, K., Pander, M., & Würkner, M. (2022). Damage patterns in float glass plates: Experiments and peridynamics analysis. *Theoretical and Applied Fracture Mechanics*, 118, Article 103264.
- Naumenko, K., Yang, Z., Ma, C.-C., & Chen, Y. (2023). Closed-form series solutions to peridynamic rod equations: Influence of kernel function. *Technische Mechanik-European Journal of Engineering Mechanics*, 43(2), 259–270.
- Park, H. W., & Oh, J. H. (2019). Study of abnormal group velocities in flexural metamaterials. *Scientific Reports*, 9(1), 13973.
- Penna, R. (2023). Bending analysis of functionally graded nanobeams based on stress-driven nonlocal model incorporating surface energy effects. *International Journal of Engineering Science*, 189, Article 103887.
- Royer, D., & Dieulesaint, E. (2000). *Elastic waves in solids I: Free and guided propagation*. Berlin: Springer.
- Silling, S. A. (2000). Reformulation of elasticity theory for discontinuities and long-range forces. *Journal of the Mechanics and Physics of Solids*, 48(1), 175–209.
- Silling, S. A., & Lehoucq, R. B. (2008). Convergence of peridynamics to classical elasticity theory. *Journal of Elasticity*, 93, 13–37.
- Silling, S. A., & Lehoucq, R. B. (2010). Peridynamic theory of solid mechanics. In H. Aref, & E. van der Giessen (Eds.), *Advances in applied mechanics: Vol. 44, Advances in applied mechanics* (pp. 73–168). Elsevier.
- Silling, S. A., Zimmermann, M., & Abeyaratne, R. (2003). Deformation of a peridynamic bar. *Journal of Elasticity*, 73, 173–190.
- Strutt, J. W. (1945). *The theory of sound. In two volumes*. New York: Dover.
- Wang, X., & Huang, Z. (2019). A possible reason about origin of singularity and anomalous dispersion in peridynamics. *Computer Modeling in Engineering & Sciences*, 121(2), 385–398.
- Weckner, O., & Abeyaratne, R. (2005). The effect of long-range forces on the dynamics of a bar. *Journal of the Mechanics and Physics of Solids*, 53(3), 705–728.
- Whitham, G. B. (1999). *Linear and nonlinear waves*. New York: John Wiley & Sons.
- Yang, Z., Ma, C.-C., Oterkus, E., Oterkus, S., & Naumenko, K. (2023). Analytical solution of 1-dimensional peridynamic equation of motion. *Journal of Peridynamics and Nonlocal Modeling*, 5(3), 356–374.