

Arabinoylans: A review on protocols for their recovery, functionalities and roles in food formulations

Fernanda Jimena Hernández-Pinto¹, Juan Daniel Miranda-Medina², Abril Natera-Maldonado³, Óscar Vara-Aldama⁴, Mary Pily Ortueta-Cabranes⁴, Jorge A. Vázquez del Mercado-Pardiño⁴, Mohammad Saeed Kharazmi⁸, Roberto Castro-Muñoz^{9,10*}, Seid Mahdi Jafari^{5,6,7*}

¹Tecnologico de Monterrey, Campus Querétaro. Av. Epigmenio González 500, Tecnológico, 76130, Santiago de Querétaro, Qro., Mexico

²Tecnologico de Monterrey, Campus Guadalajara, Av. General Ramón Corona 2514, Zapopan, 45138, Jalisco, Mexico

³Tecnologico de Monterrey, Campus Chihuahua, Av. H Colegio Militar 4700, Nombre de Dios, Chihuahua, Chih., Mexico

⁴Tecnologico de Monterrey, Campus Monterrey. Av. Eugenio Garza Sada Sur 2501 Sur, Tecnológico, 64849 Monterrey, N.L., Mexico

⁵Department of Food Materials and Process Design Engineering, Gorgan University of Agricultural Science and Natural Resources, Gorgan, Iran

⁶Universidade de Vigo, Nutrition and Bromatology Group, Department of Analytical Chemistry and Food Science, Faculty of Science, E-32004 Ourense, Spain

⁷College of Food Science and Technology, Hebei Agricultural University, Baoding 071001, China

⁸Faculty of Medicine, University of California, Riverside, CA 92679, USA

⁹Tecnologico de Monterrey, Campus Toluca. Av. Eduardo Monroy Cárdenas 2000 San Antonio Buenavista, 50110 Toluca de Lerdo, México

¹⁰Gdansk University of Technology, Faculty of Civil and Environmental Engineering, Department of Sanitary Engineering, 11/12 Narutowicza St., 80-233, Gdansk, Poland

*Corresponding Author:

E-mail: food.biotechnology88@gmail.com; roberto.castro-munoz@pg.edu.pl; castromr@tec.mx

28 Abstract

29 Arabinoxylans (AXs) are compounds with high nutritional value and applicability, including
30 prebiotics or supplementary ingredients, in food manufacturing industries. Unfortunately, the
31 recovery of AXs may require advanced separation and integrated strategies. Here, an analysis of the
32 emerging techniques to extract AXs from cereals and their by-products is discussed. This review
33 covers distinct methods implemented over the last 2-3 years, identifying that the type of method,
34 extraction source, AX physicochemical properties and pre-treatment conditions are the main factors
35 influencing the recovery yield. Alkaline extraction is among the most used methods nowadays, mostly
36 due to its simplicity and high recovery yield. Concurrently, recovered AXs applied in food
37 applications is timely reviewed, such as potential bread ingredient, prebiotic and as a wall material
38 for probiotic encapsulation, in beer and non-alcoholic beverage manufacturing, complementary
39 ingredient in bakery products and cookies, improvers in Chinese noodles, 3D food printing and
40 designing of nanostructures for delivery platforms.

41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
Keywords: Arabinoxylan; Purification; Gelling properties; Cereals; Dietary fiber; Functional foods.

101 Acronyms

102 AXs: Arabinoxylans

103 AXOS: arabinoxylooligosaccharides

104 BSG: brewers' spent grain

105 GI: Glycemic Index

106 MAX: small molecular arabinoxylans

107 NSP: non-starch polysaccharides

108 PWMB: brewers' spent grain

109 SWE: subcritical water extraction

110 SCFAs: short chain fatty acids

111 UAX: unmodified arabinoxylans

58 1. Introduction

59 As of 2022, many global trends lean into the search for new and innovative techniques towards
60 the development of environmentally conscious processes. Ongoing climate change has created a
61 heightened need for re-designing the use and obtaining high-value-added compounds from by-
62 products of the food industry. Many abundantly available by-products end up as waste (Roberto
63 Castro-Muñoz et al. 2018); thus, the present-day demand for emerging applied technologies for their
64 management (Valencia-Arredondo et al. 2020; Roberto Castro-Muñoz 2020a). This not only
65 represents an opportunity in a much-needed favourable time for taking care of the environment, but
66 also **opens a new opportunity** for the food industry to improve and make the most out of its protocols
67 and processes in multiple areas. Recovering by-products not only influences the economic spectrum
68 of the protocol but also gives out a chance for finding, studying, and investigating new benefits from
69 previously discarded materials (Castro-Muñoz, Gontarek-Castro, and Jafari 2022; Castro-Muñoz and
70 García-Depraect 2021) .

71 Particular by-products, like the ones derived from cereal processing industries, are potentially
72 a rich source of high-added value molecules (Díaz-Montes and Castro-Muñoz 2021), which are either
73 difficult to chemically synthesize or produce (Roberto Castro-Muñoz et al. 2021). This is the case of
74 arabinoxylans (AXs), which is a complex fiber providing various health benefits regulated thanks to
75 the chemical composition which are naturally present in specific parts of cell walls of distinct grains
76 including corn, wheat, rice, rye, sorghum, oats, barley and other plants. **AXs are polysaccharides**
77 **presenting various d-Xylose molecules arranged in linear backbone, which are joined due to β (1→4)**
78 **bonds, while α -L-arabinofuranosyl residues molecules are also chemically attached to the specific d-**
79 **Xylose molecules** (Lapierre et al. 2018). Their isolation has been successful, showing that the
80 different molecular weight compositions are affected directly by their source and extraction methods
81 and hence impacting on their functionality. AXs have shown functional properties acting as a
82 nutraceutical agent; as soluble fiber aids in controlling diabetes mellitus (specially type II), and
83 enhancing the metabolism of the three main macronutrients (carbohydrates, proteins and lipids) (Nie

84 et al. 2018). AXs have also exhibited a role as a prebiotic agent since they stimulate bacterial growth,
85 making them a beneficial and important factor on the microbial gut health. Being able to improve
3 86 microbial health has been linked to reducing risk in diseases caused by shifts within the gut. Other
4
5
6 87 health benefits, such as hepatoprotective and immunomodulatory activities, postprandial glucose and
7
8 88 cholesterol control, improving absorption of calcium and magnesium, enhancing colon function and
9
10
11 89 lower risk of colon cancer, have also been found (Fadel et al. 2018). However, these compounds not
12
13 90 only provide benefits to humans, but also to animals. Therefore, they can also be valuable as feed
14
15 91 additives for livestock (Siddiqui et al. 2022).

17
18 92 This review timely identifies potential sources containing AXs such as food by-products, as
19
20 93 well as to elucidate the updated strategies and methods applied for AX recovery. Knowing their
21
22
23 94 advantages and disadvantages including details on the techniques and protocols used, we will declare
24
25 95 the most appropriate processes for AXs separation. Therefore, an explicit **search for understanding** of
26
27
28 96 recent scientific literature (over the last 2-3 years) has been done. Knowing the implications of the
29
30 97 molecules of interest and their potential use in new formulations within the industry, we review the
31
32
33 98 latest research studies focused on AX exploitation for new food formulations or helping to improve
34
35 99 the existing ones, including their importance to the industry.

400 41 42 43 102 **2. Extraction strategies to recover arabinoxylans from food wastes**

44
45 103 To date, different methods have been applied for extracting AXs present in distinct sources,
46
47 104 including sugarcane bagasse, wheat bran, corn, barley, among some others. These extraction methods
48
49
50 105 include but are not limited to enzymatic and alkaline extraction. Initially, **it is obvious that selecting**
51
52
53 106 **the adequate source of recovery is a key variable to obtain a higher yield, however, it is worth**
54
55
56 107 **mentioning that the recovery strategy is planned and strongly depends on the initial physicochemical**
57
58
59 108 **composition of the AXs source.** Maize bran is a primary source of election, mostly due to the
60
61
62 109 quantities produced globally, resulting in an ease of availability (Ayala-Soto et al. 2014).

9 **2.1. Alkaline extraction method with defatted and dried maize**

110 When studying the extraction and recovery of AXs with defatted and dried maize, Herrera-
111 Balandrano et al. (2018) achieved an 8.23% recovery with an alkaline extraction method, meanwhile
112 Paz-Samaniego et al. (2016) obtained a 46% yield but when using maize wastewater or “nejayote” as
113 a primary source with acid hydrolysis extraction method, a percentage that can be interpreted as a
114 good source of recovery for AXs (Roberto Castro-Muñoz 2020b; Ramírez-Jiménez and Castro-
115 Muñoz 2020). **Another alternative source to extract AXs is wheat bran, a by-product of different
116 wheat processes, which has a significant nutrient content, such as dietary fiber and protein, and makes
117 it a product with value-added in the current market.** Concurrently, as being one of the biggest food
118 products in the world, there is nowadays an environmental concern from the under-consumed wheat
119 (Liu et al. 2020).

2.2. Hydrothermal treatment

121 By using hydrothermal treatment and on mesoporous silica catalyst, it has been reported the
122 simultaneous extraction and recovery of AXs from detached wheat bran; in this study, Sánchez-
123 Bastardo et al. (2017) observed that extracting large quantities of small fractions of AXs is promoted
124 in a short extraction time while maintaining high temperature, catalyzed with extreme acidity values.
125 For instance, when extracting at optimum conditions of 180 °C, 10 minutes, the total AX yield
126 extracted was 78% AXs (expressed as the ratio of total AXs contained in liquid extract and raw
127 material), while Li et al. (2020) reported lower recovery rates (approximately 17.98%), representing
128 a more attractive source compared with maize bran.

129 In recent years, the use of brewers’ spent grain (BSG) has been a novel source of AXs due to the
130 large amount of biomass generated by this agro-industrial process. The BSG is a by-product left after
131 barley malting and separation of the wort during the brewing process (Jaguey-Hernández et al. 2022).
132 In their study, Pérez-Flores et al. (2019) obtained a yield near 65% using alkaline extraction method,
133 representing the amount of AXs in the BSG-AX compared to the total AX amount to the original
134 sample, but a yield of 72% of pure AX. The BSG-AX contained glucose (ca. 17.94%), arabinose (ca.
135 33.74%), and xylose (ca. 37.90%).

2.3. Saccharification and fermentation process

On the other hand, Lynch et al. (2021) used a simultaneous saccharification and fermentation process on several BSG samples obtaining an average concentration of 23.5%, where they performed different treatment processes in order to obtain diverse samples containing different content of soluble AX differing in structural characteristics, including polymerization and substitution degree.

A source of recent interest for AXs extraction is the sugarcane bagasse; this source represents a high waste stream arising from sugar and alcohol industries. According to Campbell et al. (2019), bagasse exhibits different functional properties to take into consideration when choosing the method of extraction, since bagasse has a reduced solubility and greater susceptibility to enzyme action. They also evaluated an extraction of AXs from two different sources: sugarcane and wheat bread, obtaining a total recovery of 34% and 11.5%, respectively. On the other hand, Solier et al. (2022) were able to obtain a higher yield of precipitated hemicellulose (ca. 85%), corresponding to the treatment performed at high alkalinity with no use of peroxide.


Preliminarily, by analyzing the current strategies used for AX extraction, alkaline extraction is likely to be the most prominent method, mostly due to its simplicity and equipment needed, with adequate yields and concentrations depending on the primary source used. This method was described for the first time in 1998 by Zilliox and Debeire (1998), who established the use of organic solvents, such as acetic acid (also known as ethanoic acid) and ethanol, to precipitate AXs. Several washing steps are required until reaching a neutral pH; after this, an alkaline treatment is performed with a certain concentration of KOH. However, this protocol has been adapted several times in order to obtain higher concentrations of AXs. The alkaline treatment is generally followed by a liquid fraction (see **Fig. 1**) to obtain AX or even a solid fraction if it is desired to obtain lignocellulose. Alkaline extraction method has a substantial operation time between 2 and 8 h depending on the solution used; unfortunately, among its disadvantages are the hydrolysis of polysaccharides destroying its structure and when there is an improper concentration of alkali, it can break glycosidic bonds, generating many impurities and thus difficult to further process (e.g., filtration). Also, extraction solution viscosity is

162 too high while presenting strong alkali taste which affect the quality (Lihua, Xin, and Jiayang 2018).
163 It has been proved to reduce the bioactivity and antioxidant properties of AXs since the long
164 extraction time, exposes AX to longer periods of constant high temperatures and pressure, leading
165 into depolymerization of the molecules (Jaguey-Hernández et al. 2022; Daniela D. Herrera-
166 Balandrano et al. 2018; Schmidt, Wiege, and Hollmann 2021; Pérez-Flores et al. 2019; Rudjito et al.
167 2019).

168 In view of this problem, different extraction strategies and methods have emerged to reduce such
169 a limitation.

170 2.4. Subcritical water extraction

171 For instance, subcritical water extraction (SWE) is an alternative method that has demonstrated
172 to decrease the extraction time with significant recovery yields depending on the operating extraction
173 conditions. Since this technique uses water instead of a buffer solution for de-starching the source,
174 SWE offers better yields and less environmental impacts (Rudjito et al. 2019).



175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

According to de Figueiredo et al. (2017), the alkalization can be successfully done with a solution composed of KOH (24% m/V) and NaBH₂ (1% m/V), which was subsequently blended with

182 the extraction source (8% m/v) for 3 h at 35 °C. After this, the resulting liquid fraction was then pre-
183 filtered using gauze until obtaining a liquid lacking in visible solids. In a subsequent step, the AXs
184 contained in such liquid fraction were precipitated by mixing with an organic mixture containing
185 ethanol (60% v/v), acetic acid (6.7% v/v) and AXs (33.3% v/v). The precipitated matter, identified
186 as AXs, was recovered using centrifugation (for short time 15 min at 4000 g). Finally, the obtained
187 AX fraction was washed in proportional ratio (1:1 v/v) using ethanolic solution containing ethanol
188 (50%) and EDTA (0.5%) in deionized water, subsequently, the AX fraction was subjected to drying
189 (at 60 °C).

190 Following with the extraction methods of AXs, Lazaridou et al. (2008) described a protocol using
191 water treatment which consist in the solubilization of AXs by immersing the grain fragments in hot
192 water varying between 45-90 °C; observing that the higher the temperature, the shorter the exposure
193 time. This method could preserve the native structure of AX. However, the water-soluble property of
194 AXs was to some extent dependent on the type of grain and its germination degree, location within
195 the grains and chemical nature of the carbohydrate fractions present in the grain. All these aspects
196 greatly determine the overall recovery yield of AXs via water extraction (Zannini et al. 2022). To
197 some extent, Lazaridou´s protocol seem to be feasible enough for AX extraction, however, it depends
198 on the type of grain used. Importantly, the removal of starch and other carbohydrates using specific
199 enzymes may be required. Even though, the nature of the polysaccharides may result as a main
200 limiting factor when chemically crosslinked AXs to other polymer fractions (from grain cell wall)
201 are not efficiently separated and thus inherently limits the extraction efficiency.

202 2.5. Milling

203 Also, there are a variety of emerging technologies based on mechanical treatment, including
204 milling which offer many advantages representing an ecofriendly and economically feasible option
205 to obtain a homogeneous dispersion when blended with biopolymers. Such technologies provide
206 other advantages in terms of less environmental impact, controlled degradation reactions regulated
207 by high temperatures, enhanced compatibility of blends lacking in miscibility, the promotion of

208 mechano-chemical reactions and an appropriate handling of thermolabile active molecules (Gorrasi
209 and Sorrentino 2015).

210 **2.6. Extrusion**

211 For instance, extrusion stands out as a preferred technique for the preparation of extruded
212 foods with high acceptability, while impacting specific properties of the product, including water
213 solubility, water and oil absorption, expansion indices of extruded formulations and their bulk
214 density, as well as the dough viscosity (Alam et al. 2015). Demuth et al. (2020), for example,
215 performed an extraction of AX using a milling and exclusion; their results showed that this
216 mechanical treatment helped to boost the recovery yield due to more extractable AXs.

217 **2.7. Ultrasound and microwave-based technologies**

218 Ultrasound and microwave-based technologies are other successful approaches for valuable
219 solutes extraction, such as AXs. According to their principles, the use of these technologies can not
220 only result in shortened extraction times but also contributes to obtaining high recovery yield towards
221 a targeted compound. Moreover, these technologies exhibit a reduction of solvent usage (if required)
222 as cavitation bubbles pressure may disrupt the cell wall (Görgüç, Bircan, and Yılmaz 2019; Cako et
223 al. 2022) and thus result into enhanced transfer of mass in the extraction process (Jiang et al. 2019).

224 **2.8. Steam-pressure**


225 Steam-pressure application positively affects AX's extractability. Sui et al. (2018), for
226 instance, evaluated the applicability of steam pressure for insoluble fiber extraction, which could be
227 potentially transformed into a soluble one boosting its extraction with water.

228 As discussed previously, the protocols dedicated to the extraction and purification of AX depend
229 on the source of recovery, in which the extraction may imply the combination of two or more different
230 methods. In most of the protocols, chemical or biological factors, such as KOH concentration,
231 enzymatic concentration, extraction temperature, exposure time, solvent for precipitation (e.g., acetic
232 acid, ethanol) and its ratio, and number of washes, varied according to the needs of the investigation.
233 According to **Table 1**, available studies demonstrated acceptable recovery efficiencies toward AX,

234 pointing out that alkaline extraction **can offer** a yield of up to 72%, which is comparable with
235 hydrothermal synthesis with an AX yield = 78%.

236 237 **3. Potential applications of recovered AXs in food formulations**

238 As represented in **Fig. 2**, AXs have been attributed with multiple functionalities for costumers
239 as sugar substitute, probiotic and prebiotic, antioxidant, bioactive compounds for lowering
240 cholesterol, and as a nutraceutical in pharmaceutical formulations. Several studies have shown that
241 AXs greatly promote the nutritional health of the digestive system as a dietary fiber and prebiotic.
242 This molecule is contained in foods, particularly cereals as the main ingredient. Such molecules are
243 targeted for human consumption or animal feed due to their great impact in terms of different
244 biological activities, such as the synergic effect with phenolics, prebiotic activity due to fermentation,
245 cholesterol control because of the generation of short chain fatty acids (SCFAs) (Y. Zhao et al. 2017),
246 regulation of blood sugar for patients with diabetes and as a booster of the intestinal microbiota
247 composition improving the immune response (Ashaolu 2020).



9
0
Fig. 2. Main health benefits of AXs.

Table 1. Arabinoxylan extraction techniques and outcomes.

Molecule	Source	Technique	Operating conditions	Yield (%)	Reference
BSG-AX	Barley	Alkaline extraction	Temperature: 25°C Time: 8 h, complete darkness Stirring: 150 rpm Solvent: NaOH 0.5 mol L ⁻¹ (pH 11-12)	72	(Pérez-Flores et al. 2019)
BSG-AX	Barley	Alkaline extraction	Temperature: 25°C Time: 8h, complete darkness Stirring: Yes, 100 rpm Solvent: NaOH 0.5 mol L ⁻¹ (pH 11-12)	-	(Jaguey-Hernández et al. 2022)
BSG-AX	Barley	Saccharification +water treatment	Temperature: 90°C Time: 6 h Solvent: water	21.9	(Lynch et al. 2021)
Sugarcane AX	Sugarcane bagasse	Alkaline extraction	Temperature: 55°C Time: 4 h Stirring: Yes, 200 rpm Solvent: NaOH 5 mol/L and 5–10 drops of antifoaming agent	-	(Solomou et al. 2022)
Native and acetylated AX	Wheat bran	-Pretreated by α -amylase. -Subcritical water extraction (SWE) of AX from destarched wheat bran	Temperature: 90 °C for Time: 17 h Solvent: Sodium phosphate buffer (pH= 7.0)	FAX: 7.5 NAX: 31.1	(Yilmaz-Turan et al. 2020)
Water un-extractable AX	Corn bran	Ultrasonic-Microwave	Temperature: 70 °C Time: 5-30 min Power: 200-600 W Solvent: NaOH 0.15-0.35 mol/L pH= 4.5 Stirring: 4000 rpm, 20 min	25	(Jiang et al. 2019)
Feruloylated AXs	Wheat bran	Subcritical water treatment. Microwave heating extraction system	Temperature: 120-180 °C Time: 10-120 min Solvent: 10% (w/v) aqueous suspension pH 4-10	17.98	(J. Li and Du 2019)
Feruloylated AXs	Maize bran	Alkaline extraction	Temperature: 37 °C Time: 2-6 h Solvent: 0.5 N NaOH solution	8.23	(Herrera-Balandrano et al. 2019)
Maize waste water AXs	Maize wastewater “nejayote”	Acid hydrolysis extraction	Temperature: 120 °C Time: 2 h Solvent: 2 N trifluoroacetic acid	46	(Paz-Samaniego et al. 2016)
Arabinose/Xylose	Wheat bran	Hydrothermal synthesis assisted by heterogenous catalyst	Temperature: 140-180 °C Time: 10-20 min Mesoporous silica: MCM-48/Al-MCM-48 With or without Ruthenium catalyst (RuCl ₃)	78	(Sánchez-Bastardo, Romero, and Alonso 2017)

Alkali- insoluble cellulose-rich AX (CAX)	Wheat Bran	Alkaline and water extraction	Temperature: 95 °C Time: 6 h Solvent: α-amylase, ethanol	16.97	(Kaur et al. 2021)
AXs	Sugarcane bagasse and wheat bran	Alkaline and enzyme extraction	For alkaline: Temperature: 50 °C Time: 4 h Solvent: 2% H ₂ O For enzymatic: Temperature: 37 °C Time: 2 h	For sugarcane: 34 For wheat bread: 11.5	(Campbell et al. 2019)(Campbell et al. 2019)
AXs	Sugarcane bagasse	Alkaline extraction	Temperature: 35 °C Time: 3 h Solvent: 250 mL of a 24% (m/v) KOH and 1% (m/v) NaBH ₂ solution	17.5	(de Figueiredo et al. 2017)
BSG-AX	Barley	Water extraction	Temperature: 45-90 °C Time: 11 h Solvent: water	28	(Lazaridou et al. 2008)
Water- extractable AX (WEAX)	Wheat bran	Water extraction	Temperature: 60-95 °C Time: 6 h Solvent: water	73.72	(Demuth, Betschart, and Nyström 2020)

253 According to Damen et al. (2011), a combination of partially enzymatic hydrolyzed water
 254 unextractable and extractable AXs and arabinoxylo-oligosaccharides (AXOS) led to an increase of
 255 SCFAs. Likewise, the fermentation of these polysaccharides helped to lessen the intestinal pH making
 256 an inappropriate environment for pathogens while increasing the *Bifidobacteria* activity. In the same
 257 way, different formulations were tested, and an important synergic effect was found with the
 258 replacement of the proportions between the ingredients showing that the combination of the water
 259 unextractable AX (40% purity), water extractable AX (81% purity) and AXOS, together increased
 260 the butyrate levels. In the following subsections, we report some of the findings in literature dealing
 261 with the use of AXs in food formulations, along with the main implication on the resultant product.
 262 In general, **Fig. 3** overviews the main applications in food technology in which AXs are involved.




Fig. 3. Potential sources and uses of recovered AXs.

3.1. Arabinoxylans as a potential bread ingredient

Table 2 enlists the reported implementations and patents of AXs in the food science aimed at improving the nutritional value of existing food items, as well as improving specific physicochemical properties. The dietary fiber obtained from wheat bran can offer several health benefits including disease prevention and antitumor activity. Although the addition of wheat bran to bread can improve

271 the nutritional value, it can also detriment the overall quality of the bread baking causing lower
272 volume dough, bitter taste, hard crumbing and a darker color (Biliaderis et al. 1995). The relevance
273 of AXs on these effects lies in them being the main dietary fiber component of wheat bran, even if
274 water-soluble AXs can be generally beneficial for loaf volume and crumb structure. It is known that
275 the **water insoluble** AXs could be the main source of spoiling, since this type of AX competes for
276 water molecules affecting the gluten formation. According to Biliaderis et al. (1995), AXs exhibit
277 hydrophilic nature translated to high water affinity and thus compete for water molecules against
278 other compounds present in food matrices. Furthermore, AXs present a unique capability to capture
279 water molecules thanks to “oxidative gelation” and considering the influence of these on water
280 relations, based on hydrogen bonding or other specific interactions. As for food processing properties,
281 they depend on the availability and content of water; AXs certainly become relevant in the final
282 physicochemical properties and quality of wheat. Since AXs are able to bind or capture a relatively
283 high quantity of water, this increases the moisture level, together with the molecular chain mobility
284 of starch, resulting in an improved rate of starch retrogradation (Michniewicz et al. 1992; Biliaderis
285 et al. 1995).

286 It has been known for years that water-insoluble AXs can be released via alkaline or xylanase
287 enzyme treatment to give alkali-**solubilized** AXs or enzyme-**solubilized** AXs (Courtin and Delcour
288 2002). The investigation made by Xue et al. (2020) demonstrated that the quality of the bread
289 containing high quantities of wheat bran could be substantially tuned by arranging the AX structure,
290 which can be done by an enzyme treatment enriched with arabinofuranosidase to enhance the
291 efficiency of degradation while eliminating the arabinose units from the AX structure. Indeed, Wang
292 et al. (2003) explained that AXs negatively affect the formation of gluten and consequently may result
293 in a worsened gluten yield because water-unextractable solids, which include water-insoluble AXs.
294 Also, according to their water competition, AXs somehow provoked a diminished gluten yield, since
295 they limited the molecular chain motion and capability of the gluten formation by the gliadin and
296 glutenin.

297 The enzyme cocktail of TmAra and XynB can improve both the nutritional content and
298 processing functionality of wheat bran, exhibiting a synergetic response to the rheologic properties
299 of the dough and bread quality. Additionally, it displays promising applicability within emerging
300 areas for extracting antioxidants, conferring a higher added value in functional foods, nutraceuticals
301 and pharmaceutical applications.

302 Reducing dietary Glycemic Index (GI) while equilibrating high levels of carbohydrate intake
303 could be a great option for the betterment of health issues, such as being overweight, obesity and the
304 risk to acquire diabetes, especially type II. As (Hodge et al. 2004) mentioned, a promising pathway
305 to possibly achieve this relies on the possibility to replace white bread with low-GI bread. In refined
306 cereals, the contained glucose tends to be more available in the metabolic digestion which can
307 provoke insulin resistance, as continuous fluctuations of blood sugar levels may result in responses
308 to high insulin. Interestingly, AXs obtained from trifluoroacetic acid, ultrasonication processing and
309 xylanase-aided treatments could be potentially employed to regulate specific GI values desired in
310 distinct baked products (Y. Liu et al. 2020).

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Downloaded from MOST WIEDZY

Table 2. Up-to-date developments of AXs in the food products.

Application	Targeted product	Description	Reference/Patent number
Significantly improve the added value of wheat sprouts and wolfberry	Wheat malt juice wolf- berry drink	Mix the wort, wolfberry juice and softened water, add auxiliary materials, and sterilize	CN1103549606A
Added value stimulates the growth of beneficial colonic microflora.	Kvass: cereal-based and non-alcoholic beverage	Extruded rye, xylanolytic enzymes and LAB can be used for production of novel, safe and higher value non-alcoholic beverages	(Basinskiene et al. 2016)
Enhance the nutritional and technological functionality of dough rheology and steamed bread quality	Wheat bran, having a synergetic effect on the dough rheology and steamed bread	Enzyme treatment enriched with arabinofuranosidase to enhance the efficiency of degradation and modifying the AX structure	(Xue et al. 2020)
Reducing dietary GI substituting white bread with low-GI breads.	Bread	AXs obtained from trifluoroacetic acid, ultrasonication and xylanase treatments under specific conditions can be used to modulate different baked products for the desired GI	(Y. Liu et al. 2020; Hodge et al. 2004)
Quality bran-enriched cake improver	Bakery products and cookies	Favourable effects on batter stability, cake cell density and consistency. Improves the water binding properties which helps to maintain the desired crumb moist.	(Moza and Gujral 2017)
Relieving intestine symptoms, as well as psychobiotics with a beneficial effect on microbiome-brain axis and anti-inflammatory effect.	Novel food prototypes with prebiotic applications	Water-soluble extractable AXs in vitro and confirmed in vivo, increasing the number of SCFAs in the intestine, growing the beneficial bacteria in vitro and in vivo, while decreasing the pathogens in the profile of intestinal microorganisms	(C. Paesani et al. 2019; Candela Paesani et al. 2020)
Encapsulating probiotics, better texture and addition of nutritional value to the wastewater	Nejayote: wastewater generated from the maize processing	Maize wastewater enriched in AX (MWAX) to encapsulate probiotics.	(Ayala-Soto et al. 2014)
It is rich in biological activity of wheat and has a good flavor	Wheat polysaccharide drink	Treated with organic acid and then hydrolyzed, adjust the flavor, fill and sterilize	CN104522811A
Moisturize the intestines, control blood sugar and blood lipids, regulate human immunity and prevent colon cancer	Wheat bran dietary fiber	Alkaline extraction and alcohol precipitation are combined with steam explosion and compound enzyme hydrolysis	CN104544137A
Increase the storage period of the yellow peach and preserve the good flavor of the yellow peach	Yellow peach fresh- keeping treatment liquid	Dissolve vegetable oil, sucrose, iron sulfate, and kusatrol in water, and add modified AX, clove, and astragalus mixed powder	CN108935654A
Simulate the mouthfeel of sugar to improve the quality of low- calorie beverage	Composition for im- proving the taste of beverages	Add hydrolyzed β -glucan and AX to a low-calorie or zero- calorie beverage	CN109843086A
Improve the storage effect and edible effect of the composite probiotic product	Compound probiotic products	Add heat protection agent, then the probiotics are embedded with AX and soy protein	CN110623066A
The product not only has the rich fragrance of instant tea, but also can effectively assist in lowering blood sugar	Hypoglycemic instant tea	The product is composed of AX, hop extract, orange peel extract, Pu'er tea extract	CN110897023A
Extend the shelf life of bread and steamed	AX to extend the shelf life of	Modified AX is produced by enzymatic hydrolysis and debranching,	CN110938664A

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
Downloaded from mostwiedzy.pl
MOST WIEDZY

bread, and an efficient and healthy flour improver	bread and steamed bread	which is added to the dough	
Inhibit the growth of ice crystals and protect the dough network structure	AX to improve the quality of frozen dough	Modified by enzymes and added to frozen dough	CN110938665A
Increase the plasticity of the dough and reduce the baking time	Cookies	Reduce water absorption of flours with lower WE-AXc and sucrose solvent retention capacity	(Guttieri, Souza, and Sneller 2008)
Adding WE-AX can increase the volume of bread, reduce rejuvenation and extend the shelf life of bread	Bread	Higher WE-AX generally leads to higher loaf volume and quality, whereas higher WU-AXd has the opposite effect	(Courtin and Delcour 2002)
Low-alcohol beer brewed with rye has a higher viscosity and fullness, and a better taste	Mouthfeel contributor in beer	Grind the raw materials, add saaz hops after the saccharification bath; after sterilization, yeast is added for fermentation	(Langenaeken, De Schutter, and Courtin 2020)
Using 0.25-2.0% (w/w) of AX in wheat flour, for improving the quality. Thus, corroborate that using the percentage of AX in wheat flour, both, the nutrition and quality increase in the Chinese noodles	Chinese noodles	The use of UAX in AX from wheat bran flour and bread quality, due to the increment in the water absorption as well as dough formation of noodles	(Fan et al. 2016)
Development of accessible nutritional blends for 3D printing with AX as the basis	3D food printing	Aqueous solutions of starch-mango AX blends to preserve the properties related to the viscoelasticity were prepared with the addition of mineral water to incorporate them in 3D food	(Montoya et al. 2021)

313 **3.2. Arabinoxylans as prebiotics**

314 As a nondigestible fiber, AXs stimulates specific gut microbial species, specially the
315 *Bifidobacterium* spp. and *Lactobacillus* spp., having great potential as a prebiotic with health benefits
316 including the prevention of dysbiosis effect of gut microbiota, countering the proliferation of
317 pathogens as well as mitigating specific gastrointestinal disorders (Duan et al. 2022). As mentioned
318 previously, AX structure can differ depending on cereal type, the part of the grain from which it is
319 extracted, and the method of recovery. The AX structure becomes relevant regarding the way in which
320 it is degraded and fermented by the gut microbiota. Because of the structural diversity, there is still a
321 lot to learn from each function/use that could have. The most studied structures are the wheat-based
322 AXs, which have shown development of microbial species and their substantial proliferation, but
323 there is much more to know about other types of cereal grains, such as rice, oatmeal and corn
324 (Schupfer et al. 2021). However, Paesani et al. (2019) mentioned the potential of novel food
325 formulations and ingredient prototypes such as AXs and probiotics for relieving intestine symptoms,
326 as well as the possible use as psychobiotics exhibiting a positive effect on microbiome-brain axis.
327 Punctually, this study revealed that *Lactobacillus reuteri* ATCC23272 had anti-inflammatory
328 activity, and its synergistic effect with AXs, opens a new window of developing new food
329 formulations. Another development was presented by Paesani et al. (2020), who studied the prebiotic
330 activity of water-soluble extractable AXs *in vitro* and proved *in vivo* test, showing a higher prebiotic
331 capacity by raising the number of SCFAs in the intestine, growing the health-promoting bacteria in
332 both assays, while mitigating the number of pathogens in the total profile of microorganisms present
333 in the intestine.

335 **3.3. Arabinoxylans in beer manufacturing**

6 AXs are able to cause several issues credited to the increment of wheat employed for brewing
7 the wheat beer, which could compromise the quality of the obtained wheat beer, including the high
8 viscosity of the wort, the haze formation, and the sensorial properties and wheat beer foam
9 performance. Langenaeken et al. (2020) confirmed the significance of AXs and its degree of

340 polymerization for a better mouthfeel attribute of produced beverages, such as alcohol-free beer, low-
341 alcohol content beers and light beers as it increases the viscosity values of the beer. Experimentally,
342 those beers were brewed replacing barley malt grits (20%) with weather oat, non-malted barley, or
343 rye in comparison with a control. As for the one brewed with rye, a substantial increase (ca. 53 %) in
344 AXs presence was observed while increasing the degree of polymerization from 29 to 50. In
345 comparison with the control beer presenting a viscosity = 1.48 mm²/s, this property was observed to
346 increase in the rye beer (ca. 1.85 mm²/s). It is worth mentioning that the highest viscosity in beer was
347 shown by the non-malted rye, this latter phenomenon was credited to the presence of high molecular
348 weight AXs in the final beer. During sensory analysis, the beer, containing rye (30%), was found to
349 as the fullest beer, emphasizing the AXs relevance and their related polymerization degree in the
350 fullness and viscosity of low-alcohol beers, presenting this approach as a viable solution to improve
351 mouthfeel characteristics and postulating that the improved viscosity is attributed to the delivery of
352 higher molecular weight AXs from un-malted cereals than the ones that could be obtained from
353 malted cereals (Langenaeken, De Schutter, and Courtin 2020).

354 In recent years, pure wheat malt beer (PWMB) has risen in popularity due to its imaginably
355 functional properties, such as aroma and taste (Schneiderbanger et al., 2016). As non-starch
356 polysaccharides (NSPs), including the AXs, are the main component in wheat beer, it has been stated
357 by Li et al. (2020) that NSPs can certainly enhance the body of beers and can favorably have an
358 impact on the haze and foam retention of beers, being crucial indicators for consumers to determine
359 whether the beer has a good or bad quality.

360 The potential health value and functional properties of NSPs in PWMB were studied by Díaz
361 et al. (2023), who found out that its antioxidant, prebiotic activities, hypoglycemic and hypolipidemic
362 effects arise. Likewise, Michiels et al. (2023) confirmed the impact of AXs on the brewing process
363 contributing to the palate fullness of beers. This concluded that this promising tunable cereal
364 component, depending on the arabinose units where most often, a low AX concentration is sought to
365 reduce wort viscosity and consequently eliminate wort and beer filtration complications (Sadosky et.,
366 2002).

367 **3.4. Arabinoxylans for probiotic encapsulation**

368 Maize is one of the major sources of food in Mexico. To improve its nutritional value, the
369 maize kernel is subjected to nixtamalization processes (Roberto Castro-Muñoz and Yáñez-Fernández
370 2015; R Castro-Muñoz et al. 2015), in which maize is exposed to an alkali treatment generating large
371 volumes of wastewater which contain gelling AX (Roberto Castro-Muñoz, Fíla, and Durán-Páramo
372 2019). Therefore, there is an interest focused in the rheological, structural and microstructural
373 characteristics of this bio-based material. The capability of maize wastewater enriched in AX
374 (MWAX) to encapsulate probiotics has been studied (Paz-Samaniego et al. 2016). Nixtamalization,
375 as it is called the alkali cooking in Mexico, is widely used to have a better texture and for the addition
376 of nutritional value to the maize and the wastewater generated from the maize processing is known
377 as “Nejayote”, which is considered as highly alkaline with high chemical and biological oxygen
378 demand. Throughout the nixtamalization procedure, once maize bran is removed from the kernel,
379 AXs and cell wall components are partially solubilized in Nejayote, formerly described as a source
380 of gelling AX (Ayala-Soto et al. 2014). Among the most significant findings, is the encapsulation of
381 probiotic *Bifidobacterium* in MWAX gels with a 10% (w/v) where the ones containing and
382 maintaining the maximum probiotic content with a 1×10^7 CFU/mL. Thus, because of the pore size
383 of the matrix that surrounds it, the capsule was about 11 times smaller than the probiotic (Paz-
384 Samaniego et al. 2016).

385 **3.5. Arabinoxylans in bakeries and confectioneries**

386 AXs, as soluble fibers, are mostly present in the endosperm of cereals. Most of their
387 physicochemical properties have been potentially exploited, for example as baking additives, to
388 enhance several properties in the final products, such as consistency property of the dough, loaf
389 volume increase, improve crumb structure, or decrease in the bread staling. So far, AXs have been
0 described as palatable ingredients, in which the AX-enriched foods present excellent acceptability by
1 consumers. For example, AXs, especially the ones present in hard wheat varieties, are widely known
2 and chosen to provide better loaf volume in bread, which also enrich the protein intake and gluten
3 strength (Kiszonas et al. 2015). Several studies have reported positive effects of AXs in different

394 cookies, where the use of flours in combination with these soluble fibers results in a product with a
395 low spread ratio, a decrease in diameter and a firm consistency, which are phenotypic characteristics
396 of great importance for a quality product. The non-starch polysaccharide fraction in wheat flour
397 cookie-making was studied by Guttieri, Souza, and Sneller (2008). It was documented that higher
398 arabinose/xylose ratio demonstrated a tendency in the depletion of water absorption contained in
399 entire grain flour leading to a soft cookie. Also, the influence of substituting small quantity of sugar,
400 or flour, by AX oligosaccharides in cookies was analysed. According to Pareyt et al. (2011), when
401 the flour was replaced, the cookie diameter was somewhat decreased while the hardness behaved
402 opposite. Likewise, it was reported that AXs in cakes have potential benefits, although it has not been
403 extensively studied (Oliete et al. 2010). To some extent, it is recommended to select the flours based
404 on the concentration of the soluble pentosan which must be high. However, the use of xylans has
405 been highlighted as a quality bran-enriched cake improver (Lebesi and Tzia 2011). Furthermore,
406 when baking with high content AX flours, the air cell in final products is smaller which is best viewed
407 by consumers (Moza and Gujral 2017). Particularly, they conclude that when AXs are present, there
408 are multiple favourable effects on batter stability, cake cell density and consistency. Thus, it improves
409 the water binding properties which helps to maintain the desired crumb moist.

Very recently, Molina et al. (2021) evaluated the effect of isolated AXs in preparing model
flours. The refined flour was added with both extractable and water unextractable AXs. In this
investigation, AXs showed a meaningful impact on the mechanical properties, especially elasticity,
increasing the starch gelatinization degree from 24 up to 36%.

3.6. Arabinoxylans in non-alcoholic beverages

Nowadays, the producers of non-alcoholic drinks aim to distinguish their products in the
market of beverages with additional health benefits and functionality. This category is already sought
and commercialized on global markets by healthy lifestyle consumers. They choose to buy fortified
drinks, e.g., water enriched with minerals/vitamins and fruits, and more interested on the functionality
features of ready-to-drink food items (Busuricu et al. 2022). Also, non-alcoholic beverage marketing
is pervasive across multiple media predominantly promoting products with high content in fat, sugar,

421 and/or salt. Given the evidence of diet-related noncommunicable disease risk and obesity, the World
422 Health Organization (WHO) and other authoritative bodies have stepped in to shape policies for
423 regulating unhealthy beverages marketing and reduce the impact in children specially (Boyland et al.
424 2022; Ngqangashe et al. 2022). This is why conducting biotechnological research to find new
425 strategies for enhancing the functionality of non-alcoholic drinks is a big opportunity for the industry.
426 In this field, AXs have a potential to increase the value of these drinks, as mentioned on the
427 (Basinskiene et al. 2016) approach for non-alcoholic beverages called “Kvass”, which is a typical
428 cereal-based beverages manufactured from fermented barley malt and rye, or stale rye bread in eastern
429 European countries (e.g., Lithuania). AX-derived arabinoxylooligosaccharides (AXOS), represent an
430 emerging food ingredient as prebiotics thanks to their exceptional stimulation of the growth of
431 beneficial microflora, as mentioned in the previous section. Concluding that extruded rye, along with
432 xylanolytic enzymes and LAB bacteria, could be interestingly employed within the manufacture of
433 new, safe and higher-value-added non-alcoholic beverages (Basinskiene et al. 2016).

3.7. Arabinoxylans for the improvement of Chinese noodles

434 Being dietary fiber as an important element in human diet and AX a main component of both
435 soluble/insoluble roughage, they indeed provide numerous health benefits such as antioxidant
436 properties. Several works have evidenced that AXs improve substantially the quality of flour and
437 bread since they increment the water absorption capacity as well as dough formation (Guttieri, Souza,
438 and Sneller 2008). Reports have proven that noodles account for 30 up to 35%, of the total produced
439 wheat flour, as they are known to be a popular food in China. With this in mind, Fan et al. (2016)
440 used ultrasonication to treat unmodified AX (UAX) for obtaining small molecular AXs (MAX) and
441 later studied the effect on the color, water properties, cooking characteristics, chemical interactions
442 between free sulfhydryl chemical groups and generated disulfide bonds, and texture of this classic
443 meal. When AX content increased, the dough sheet appeared to be darker, while the texture of the
4 prepared noodles improved but it was worsened with higher content than 1% AXs (Buksa, Nowotna,
5 and Ziobro 2016)(Zhang et al. 2011). Apart from that, the presence of free sulfhydryl chemical
6 functionalities varied, e.g., they decreased by 15.4% and 19.3% when incorporated UAX and MAX,
7

448 (0.5%), respectively. Finally, findings on the water absorption improvement and cooking loss rate
449 were observed to decrease up to 1.0% of AX addition. In this study, the authors confirmed that by
450 incorporating lower AXs (between 0.25–1.0%) in the final wheat flour, h to improve the quality and
451 nutritional value of Chinese noodles.

3.8. Starch-arabinoxylans in 3D food printing

453 3D-food printing materials, such as isolated proteins, carbohydrates (alginate), and fats, are emerging
454 alternatives in homemade cookies in recent years. Apart from their advantages for complex
455 geometries, modifications in texture and specified nutritional contents are desired (Baiano 2022). In
456 this emergent field, Montoya et al. (2021) developed accessible nutritional blends for 3D printing
457 with the help of starch, lyophilized fruit as basis and AXs. In the case of AXs, they were proposed
458 since they own specific characteristics, including adhesive properties, thickener stabilizer,
459 emulsification properties, and film-forming properties (Ayala-Soto, Serna-Saldívar, and Welti-
460 Chanes 2016). By adding AXs, the preparation of blends containing small starch concentration was
461 done intentionally while aiming to preserve specific attributes related to the viscoelasticity. In this
462 study, the first batch of starch-mango aqueous mixtures (0/100, 25/75, 50/50, 75/25, 100/0) were
463 prepared, and second batch contained starch-AX mixtures (50/50, 75/25, 100/0) with the addition of
464 mineral water in first place, and then starch portion, and lastly followed by AXs fraction, as shown
465 in **Table 3**. The three starch-AX blends, identified as 100/0, 75/25, 50/50 ratio, were selected for
466 further 3D printing protocol to determine the best blend with the maximum printed height. Especially,
467 the one with 75/25 formulation displayed a pseudoplastic and shear thinning viscous properties
468 similar to pure starch. Thus, they concluded that appropriate starch/AXs combinations could also
469 form hydrogels, or reticulated natural composites, that are dependent on the starch ratio; when having
470 a 50% of this, they are in the form of hydrogels, on the contrary, become complex composites. At
1 this point, it was proven that high contents in starch-AX blends are very suitable and then suggested
2 for 3D food printing due to its nutritional value (Liu et al. 2018).

3 **Table 3.** Composition of the different starch-mango and AX blend formulations.

Formulation	Water %	Starch %	Mango %	AXs%
Starch (100)	80	20	0	0

Starch/Mango (75/25)	80	15	5	0
Starch/Mango (50/50)	80	10	10	0
Starch/Mango (25/75)	80	5	15	0
Starch/Mango (0/100)	80	0	20	0
Starch/AXs (75/25)	80	15	0	5
Starch/AXs (50/50)	80	10	0	10

3.9. Arabinoxylans in nanodelivery platforms

It is known that the food industry is likely to be one of the main fields demanding large quantities of water and energy consumption. As there is an abrupt increase in global population, the future perspective is to meet higher agricultural sources while mitigating any possible negative impact on the natural resources (e.g. water bodies) and environment (Schneider et al. 2021). Also, climate change plays an important role in future agricultural production, while temperature effects become relevant for policy-makers, farmers, agriculturalists, and crop breeders, who need to guarantee worldwide food systems. To some extent, developing sustainable crop systems and adaptation strategies for specific regions are crucial to guarantee the future food supply chain of an increased global population (C. Zhao et al. 2017). In this regard, Sarker et al. (2020) presented a sustainable idea of using chemically modified AXs, which were successfully obtained from agro-food waste and by-products, as portable materials for gene delivery in agrochemical area. Particularly, nanocarriers should offset the three main identified drawbacks of modern delivery methods in plant systems: (i) as a specific nanoparticle size can permeate through the cell wall, (ii) tuning and adjusting the surface properties (such as charge, hydrophilic/hydrophobic) to carry diverse compounds and cargo, and (iii) extended applicability in plant-based systems (Cunningham et al. 2018). It was proven that AXs, especially the ones present in wheat bran, are analogs with similar chemical properties like the ones to obtained from wheat flour and therefore can be used to construct DNA polyplexes with cationically derivatized AXs, as described in **Fig. 4**. These biodegradable nanocarriers with an average size of 150–200 nm can be used as delivery platforms with facilitated transport ability of protein and nucleic acid fractions in plants (Sarker et al. 2020).




Fig. 4. Nanocarriers based on wheat bran obtained from AXs for nucleic acid delivery. Taken from (Sarker et al. 2020).

3.10. Structure-function relationship of AXs and their application within food systems

Throughout the years, cereals grains have been extensively studied due to their wide range of ingredients with a varying degree of structural and functional complexity. The characterization of the proteins and starches, major components in cereals, has allowed to demonstrate its irrefutably functional role. The interest in the study of cereal cell wall polysaccharides stems from the recognition of the role of these minor components which affects to a degree human nutrition, physical and chemical properties, processing behavior and shelf life. Despite the diverse non-starch polysaccharides that form the cell walls of cereal grains, β -glucans and AXs molecular structures have received the most attention as well as oligosaccharides derived from these polymers. Due to it, endo- β -xylases and endo- β -glucanases have permitted structural examination of these cereal cell wall at its finest detail.

Both polysaccharides afterward mentioned display halfway dissolvability in water and thus in the human diet, notwithstanding the physiological impacts related to the consumption of insoluble fiber. In food systems, β -glucans and AXs may likewise modify the texture, water binding properties and tangible characteristics of a food product (Eliasson et al. 1996). From a wholesome and practical perspective, cereal foods rich in non-starch polysaccharides meet the description of “functional foods” as they give some of the ordinary quality characteristics of a food, like sustenance and texture.

As an example, there is a wide range of studies on the role of AXs in the bread-making process that seek to expound on the effects of AXs on properties such as water adsorption of dough, loaf

518 volume and firmness. Once the studies were run, the one in which water-soluble AXs were added to
519 bread formulations, demonstrated an increase in the water adsorption of the dough and improved the
520 loaf volume of wheat and rye bread. Therefore, the conclusions point that the improvement in the loaf
521 volume of bread is dependent on the AX's concentration in the dough system and it was importantly
522 debated that the concentration which ideally upgrades the loaf volume is related to factors such as the
523 quality of the bread flour along with the molecular weight and same structure indicating that the
524 higher the concentration is compared to the optimum of AX can cause viscosity built up in the dough
525 and thereby hinder the loaf expansion (Izydorzyc & Biliaderis, 2000).

3.11. Potential of purified/isolated AX in food systems

526 There are many studies reporting on the functional role of pentoses (the main forming structure of
527 AXs), particularly as it relates to the breadmaking process giving rise to various experimental findings
528 that are attributed to diverse factors, to mention some are the method of isolation, degree of purity,
529 composition of pentosan preparation and baking systems. Nonetheless, with most proof
530 recommending that the functional properties of pentosans are primarily due to the AX components.
531 When AX is added to wheat flour, these last certainly compete with the dough water (Izydorzyc et
532 al., 1991 & Biliaderis).

4. Concluding remarks and perspectives

533 The purpose of this comprehensive review is to reveal the up-to-date findings on the AXs
534 extraction, their health benefits, and applications in food products. This review reports, for the first
535 time, the main efforts of the scientific community on the different strategies and methods for the
536 extraction, separation, and recovery of AXs from distinct sources and by-products (e.g.,
537 nixtamalization wastewaters). From this review, it can be concluded that AX purification is directly
538 affected by the type of source, germinating state, location in the grain, pre-treatment, source
539 conditioning and finally the extraction technique. However, a detailed description of the extraction
540 conditions of AXs may be useful to overcome the main constraints during the purification protocols

544 and to obtain reproducible data from different studies. AXs have potential health benefits, including
545 SCFAs modulation in colon tissue, increased antioxidant activity, as well as reduced glycaemic
546 response through a variety of mechanisms. The chemical structure of AXs has a decisive influence
547 on the physical properties when contained in foods; generally, AXs, exhibiting low molecular weight
548 and high solubility properties, seems to be suitable for better food systems. Unfortunately, the pre-
549 treatment strategies, extraction and purification techniques and used food manufacture parameters
550 greatly dictate the resulting properties of AXs and their performance in food formulations, suggesting
551 that the functionality properties of isolated AX could be tuned to meet the quality needed for the end-
552 products. To date, alkaline extraction has been the most used method for AX extraction, especially
553 due to its simplicity, the easy to implement facilities and the operation time (between 2 to 8 h).
554 Unfortunately, the main inconvenience of this method relies on reducing the bioactivity and
555 antioxidant properties of AXs (Jaguey-Hernández et al. 2022; Pérez-Flores et al. 2019; Daniela D.
556 Herrera-Balandrano et al. 2018; Rudjito et al. 2019). Regarding the probiotic encapsulation, once the
557 *Bifidobacterium* was already studied and its efficacy has been proven, AXs represent an opportunity
558 for future studies proposing another probiotic models, such as *Lactobacillus* or a mixture of both,
559 giving rise to a healthier gut population microbiota in the consumer. Concerning the effect of AXs in
560 bakery products and cookies, findings lead to possible approaches reducing sugar levels while
561 increasing fiber content, obtaining enriched cookies with high plasticity in the dough and more
562 efficient baking (Pareyt et al. 2011).

563 Furthermore, up-to-date developments of AXs in food products were described, in which the
564 modification of AXs by enzymes results in having a positive effect on the shelf life of a variety of
565 cereals, to mention just a few. They act as a protection for the dough network structure, enhancers of
566 viscosity, water absorption, foam stabilizers, among others. Thus, these polysaccharides'
567 considerable importance in cereal-based processes has been widely accepted and studied due to their
568 nutritional benefits and properties giving rise to an increasing demand for arabinoxylan-enriched
569 products.

570 **Acknowledgments**

571 Financial support from Nobelium Joining Gdańsk Tech Research Community (contract number DEC
572 33/2022/IDUB/1.1; NOBELIUM nr 036236) is gratefully acknowledged. R. Castro-Muñoz also
573 acknowledges the School of Engineering and Science and the FEMSA-Biotechnology Center at
574 Tecnológico de Monterrey for their support through the Bioprocess (0020209I13) Focus Group.

576 **Conflict of Interest**

577 The authors declare no conflict of interest.

579 **References**

- 580 Alam, M S, Ms Jasmeen Kaur, Ms Harjot Khaira, and Ms Kalika Gupta. 2015. “Extrusion and
581 Extruded Products : Changes in Quality Attributes as Affected by Extrusion Process
582 Parameters : A Review.” *Critical Reviews in Food Science and Nutrition*, no. February 2015:
583 37–41. <https://doi.org/10.1080/10408398.2013.779568>.
- 584 Ashaolu, Tolulope Joshua. 2020. “Immune Boosting Functional Foods and Their Mechanisms: A
585 Critical Evaluation of Probiotics and Prebiotics.” *Biomedicine and Pharmacotherapy* 130:
586 110625. <https://doi.org/10.1016/j.biopha.2020.110625>.
- 587 Ayala-Soto, Fabiola E., Sergio O. Serna-Saldívar, Silverio García-Lara, and Esther Pérez-Carrillo.
588 2014. “Hydroxycinnamic Acids, Sugar Composition and Antioxidant Capacity of
589 Arabinoxylans Extracted from Different Maize Fiber Sources.” *Food Hydrocolloids* 35
590 (March): 471–75. <https://doi.org/10.1016/j.foodhyd.2013.07.004>.
- 591 Ayala-Soto, Fabiola E., Sergio O. Serna-Saldívar, and Jorge Welti-Chanes. 2016. “Effect of
592 Processing Time, Temperature and Alkali Concentration on Yield Extraction, Structure and
593 Gelling Properties of Corn Fiber Arabinoxylans.” *Food Hydrocolloids* 60: 21–28.
594 <https://doi.org/10.1016/j.foodhyd.2016.03.014>.
- 595 Baiano, Antonietta. 2022. *3D Printed Foods: A Comprehensive Review on Technologies*,

- 596 *Nutritional Value, Safety, Consumer Attitude, Regulatory Framework, and Economic and*
597 *Sustainability Issues. Food Reviews International*. Vol. 38. Taylor & Francis.
598 <https://doi.org/10.1080/87559129.2020.1762091>.
- 599 Basinskiene, Loreta, Grazina Juodeikiene, Daiva Vidmantiene, Maija Tenkanen, Tomas
600 Makaravicius, and Elena Bartkiene. 2016. "Non-Alcoholic Beverages from Fermented Cereals
601 with Increased Oligosaccharide Content." *Food Technology and Biotechnology* 54 (1): 36–44.
602 <https://doi.org/10.17113/ftb.54.01.16.4106>.
- 603 Biliaderis, Costas G, Marta S Izydorczyk, and Onkar Rattan. 1995. "Effect of Arabinoxylans on
604 Bread-Making Quality of Wheat Flours" 53: 165–71.
- 605 Boyland, Emma, Lauren McGale, Michelle Maden, Juliet Hounsome, Angela Boland, and Andrew
606 Jones. 2022. "Systematic Review of the Effect of Policies to Restrict the Marketing of Foods
607 and Non-Alcoholic Beverages to Which Children Are Exposed." *Obesity Reviews* 23 (8): 1–
608 21. <https://doi.org/10.1111/obr.13447>.
- 609 Buksa, Krzysztof, Anna Nowotna, and Rafał Ziobro. 2016. "Application of Cross-Linked and
610 Hydrolyzed Arabinoxylans in Baking of Model Rye Bread." *Food Chemistry* 192: 991–96.
611 <https://doi.org/10.1016/j.foodchem.2015.07.104>.
- 612 Busuricu, Florica, Verginica Schroder, Doina Margaritti, Andreea Hortensa Anghel, and Sirma
613 Tomos. 2022. "Nutritional Quality of Some Non-Alcoholic Beverages from the Romanian
614 Market." *Technium BioChemMed* 3 (1): 1–6. <https://doi.org/10.47577/biochemmed.v3i1.5471>.
- 615 Cako, Elvana, Zhaohui Wang, Roberto Castro-Muñoz, Manoj P. Rayaroth, and Grzegorz Boczkaj.
616 2022. "Cavitation Based Cleaner Technologies for Biodiesel Production and Processing of
617 Hydrocarbon Streams: A Perspective on Key Fundamentals, Missing Process Data and
618 Economic Feasibility – A Review." *Ultrasonics Sonochemistry* 88 (February).
619 <https://doi.org/10.1016/j.ultsonch.2022.106081>.
- 620 Campbell, G. M., N. Čukelj Mustač, M. Alyassin, L. D. Gomez, R. Simister, J. Flint, D. J. Philips,
621 M. J. Gronnow, and N. J. Westwood. 2019. "Integrated Processing of Sugarcane Bagasse:

- 622 Arabinoxylan Extraction Integrated with Ethanol Production.” *Biochemical Engineering*
623 *Journal* 146 (November 2018): 31–40. <https://doi.org/10.1016/j.bej.2019.03.001>.
- 624 Castro-Muñoz, R, G I Cerón-Montes, B E Barragán-Huerta, and J Yáñez-Fernández. 2015.
625 “Recovery of Carbohydrates from Nixtamalization Wastewaters (Nejayote) by Ultrafiltration.”
626 *Revista Mexicana de Ingeniería Química* 14 (3): 735–44.
- 627 Castro-Muñoz, Roberto. 2020a. “Membrane Technologies for the Production of Nonalcoholic
628 Drinks.” In *Trends in Non-Alcoholic Beverages*, edited by Charis M. Galanakis, 53:141–59.
629 Elsevier.
- 630 ———. 2020b. “Retention Profile on the Physicochemical Properties of Maize Cooking By-
631 Product Using a Tight Ultrafiltration Membrane.” *Chemical Engineering Communications* 207
632 (7): 887–95. <https://doi.org/10.1080/00986445.2019.1618844>.
- 633 Castro-Muñoz, Roberto, Blanca E. Barragán-Huerta, Vlastimil Fíla, Pierre Charles Denis, and René
634 Ruby-Figueroa. 2018. “Current Role of Membrane Technology: From the Treatment of Agro-
635 Industrial by-Products up to the Valorization of Valuable Compounds.” *Waste and Biomass*
636 *Valorization* 9 (4): 513–529. <https://doi.org/10.1007/s12649-017-0003-1>.
- 637 Castro-Muñoz, Roberto, Vlastimil Fíla, and Enrique Durán-Páramo. 2019. “A Review of the
638 Primary By-Product (Nejayote) of the Nixtamalization During Maize Processing: Potential
639 Reuses.” *Waste and Biomass Valorization* 10 (1): 13–22. <https://doi.org/10.1007/s12649-017-0029-4>.
- 640
641 Castro-Muñoz, Roberto, and Octavio García-Depraect. 2021. “Membrane-Based Harvesting
642 Processes for Microalgae and Their Valuable-Related Molecules: A Review.” *Membranes* 11
643 (8): 1–21. <https://doi.org/10.3390/membranes11080585>.
- 644
645 Castro-Muñoz, Roberto, Octavio García-Depraect, Elizabeth León-Becerril, Alfredo Cassano,
646 Carmela Conidi, and Vlastimil Fíla. 2021. “Recovery of Protein-Based Compounds from Meat
647 by-Products by Membrane-Assisted Separations: A Review.” *Journal of Chemical Technology*
648 *and Biotechnology* 96 (11): 3025–42. <https://doi.org/10.1002/jctb.6824>.

- 648 Castro-Muñoz, Roberto, Emilia Gontarek-Castro, and Seid Mahdi Jafari. 2022. “Up-to-Date
649 Strategies and Future Trends towards the Extraction and Purification of Capsaicin: A
650 Comprehensive Review.” *Trends in Food Science and Technology* 123 (July 2021): 161–71.
651 <https://doi.org/10.1016/j.tifs.2022.03.014>.
- 652 Castro-Muñoz, Roberto, and Jorge Yáñez-Fernández. 2015. “Valorization of Nixtamalization
653 Wastewaters (Nejayote) by Integrated Membrane Process.” *Food and Bioproducts Processing*
654 95: 7–18. <https://doi.org/10.1016/j.fbp.2015.03.006>.
- 655 Courtin, C. M., and J. A. Delcour. 2002. “Arabinoxylans and Endoxylanases in Wheat Flour Bread-
656 Making.” *Journal of Cereal Science* 35 (3): 225–43. <https://doi.org/10.1006/jcrs.2001.0433>.
- 657 Cunningham, Francis J., Natalie S. Goh, Gozde S. Demirer, Juliana L. Matos, and Markita P.
658 Landry. 2018. “Nanoparticle-Mediated Delivery towards Advancing Plant Genetic
659 Engineering.” *Trends in Biotechnology* 36 (9): 882–97.
660 <https://doi.org/10.1016/j.tibtech.2018.03.009>.
- 661 Damen, Bram, Joran Verspreet, Annick Pollet, Willem F. Broekaert, Jan A. Delcour, and
662 Christophe M. Courtin. 2011. “Prebiotic Effects and Intestinal Fermentation of Cereal
663 Arabinoxylans and Arabinoxylan Oligosaccharides in Rats Depend Strongly on Their
664 Structural Properties and Joint Presence.” *Molecular Nutrition and Food Research* 55 (12):
665 1862–74. <https://doi.org/10.1002/mnfr.201100377>.
- 666 Demuth, Teresa, Julian Betschart, and Laura Nyström. 2020. “Structural Modifications to Water-
667 Soluble Wheat Bran Arabinoxylan through Milling and Extrusion.” *Carbohydrate Polymers*
668 240 (January): 116328. <https://doi.org/10.1016/j.carbpol.2020.116328>.
- 669 Díaz-Montes, Elsa, and Roberto Castro-Muñoz. 2021. “Edible Films and Coatings as Food-Quality
670 Preservers: An Overview.” *Foods* 10 (2): 249. <https://doi.org/10.3390/foods10020249>.
- 671 Díaz, A. B., Durán-Guerrero, E., Valiente, S., Castro, R., & Lasanta, C. (2023). Development and
672 Characterization of Probiotic Beers with *Saccharomyces boulardii* as an Alternative to
673 Conventional Brewer’s Yeast. *Foods*, 12(15), 2912.

- 674 Duan, Hui, Leilei Yu, Fengwei Tian, Qixiao Zhai, Liuping Fan, and Wei Chen. 2022. “Antibiotic-
675 Induced Gut Dysbiosis and Barrier Disruption and the Potential Protective Strategies.” *Critical*
676 *Reviews in Food Science and Nutrition* 62 (6): 1427–52.
677 <https://doi.org/10.1080/10408398.2020.1843396>.
- 678 Eliasson, A. C., & Gudmundsson, M. (1996). *Starch: physicochemical and functional aspects*. Food
679 Science And Technology-New York-Marcel Dekker-, 431-504.
- 680 Fadel, Abdulmannan, Ayman M Mahmoud, Jason J Ashworth, Weili Li, Yu Lam Ng, and Andrew
681 Plunkett. 2018. “Health-Related Effects and Improving Extractability of Cereal
682 Arabinoxylans.” *International Journal of Biological Macromolecules* 109: 819–31.
683 <https://doi.org/10.1016/j.ijbiomac.2017.11.055>.
- 684 Fan, Ling, Sen Ma, Xiaoxi Wang, and Xueling Zheng. 2016. “Improvement of Chinese Noodle
685 Quality by Supplementation with Arabinoxylans from Wheat Bran.” *International Journal of*
686 *Food Science and Technology* 51 (3): 602–8. <https://doi.org/10.1111/ijfs.13042>.
- 687 Figueiredo, Franciane Cristina de, Ana Flavia Azevedo Carvalho, Michel Brienzo, Tania Sila
688 Campioni, and Pedro de Oliva-Neto. 2017. “Chemical Input Reduction in the Arabinoxylan
689 and Lignocellulose Alkaline Extraction and Xylooligosaccharides Production.” *Bioresource*
690 *Technology* 228: 164–70. <https://doi.org/10.1016/j.biortech.2016.12.097>.
- 691 Görgüç, Ahmet, Cavit Bircan, and Fatih Mehmet Yılmaz. 2019. “Sesame Bran as an Unexploited
692 By-Product: Effect of Enzyme and Ultrasound-Assisted Extraction on the Recovery of Protein
693 and Antioxidant Compounds.” *Food Chemistry* 283 (October 2018): 637–45.
694 <https://doi.org/10.1016/j.foodchem.2019.01.077>.
- 695 Gorrasi, Giuliana, and Andrea Sorrentino. 2015. “Mechanical Milling as a Technology to Produce
696 Structural and Functional Bio-Nanocomposites.” <https://doi.org/10.1039/c5gc00029g>.
- 697 Guttieri, Mary J., Edward J. Souza, and Clay Sneller. 2008. “Nonstarch Polysaccharides in Wheat
698 Flour Wire-Cut Cookie Making.” *Journal of Agricultural and Food Chemistry* 56 (22): 10927–
699 32. <https://doi.org/10.1021/jf801677a>.

- 700 Herrera-Balandrano, D.D, J.G Báez-González, E Carvajal-Milán, Gerardo Mendez-Zamora, Vania
701 Urías-Orona, Carlos A Amaya-guerra, and Guillermo Niño-Medina. 2019. “Feruloylated
702 Arabinoxylans from Nixtamalized Maize Frankfurter Sausages.” *Molecules* 24 (2056): 1–11.
703 Herrera-Balandrano, Daniela D., Juan G. Báez-González, Elizabeth Carvajal-Millán, Dolores Mui-
704 Rangel, Vania Urías-Orona, Ana L. Martínez-López, Jorge A. Márquez-Escalante, José B.
705 Heredia, Trust Beta, and Guillermo Niño-Medina. 2018. “Alkali-Extracted Feruloylated
706 Arabinoxylans from Nixtamalized Maize Bran Byproduct: A Synonymous with Soluble
707 Antioxidant Dietary Fiber.” *Waste and Biomass Valorization* 0 (0): 0.
708 <https://doi.org/10.1007/s12649-018-0462-z>.
709 Hodge, Allison M., Dallas R. English, Kerin O’Dea, and Graham G. Giles. 2004. “Glycemic Index
710 and Dietary Fiber and the Risk of Type 2 Diabetes.” *Diabetes Care* 27 (11): 2701–6.
711 <https://doi.org/10.2337/diacare.27.11.2701>.
712 Izydorczyk, M. S., & Biliaderis, C. G. (1995). Cereal arabinoxylans: advances in structure and
713 physicochemical properties. *Carbohydrate polymers*, 28(1), 33-48.
714 Izydorczyk, M. S., & Biliaderis, C. G. (2000). Structural and functional aspects of cereal
715 arabinoxylans and β -glucans. *Developments in Food Science*, 361–384. doi:10.1016/s0167-
716 4501(00)80016-
717 Jaguey-Hernández, Yari, Cecilio Tapia-Ignacio, Karina Aguilar-Arteaga, Luis Guillermo González-
718 Olivares, Evodio Pedro Castañeda-Ovando, Nelly Cruz-Cansino, Deyanira Ojeda-Ramirez,
719 and Araceli Castañeda-Ovando. 2022. “Thermoplastic Biofilms Obtained from an
720 Arabinoxylan-Rich Fraction from Brewers’ Spent Grain: Physicochemical Characterization
721 and Thermal Analysis.” *Biomass Conversion and Biorefinery*, no. 0123456789.
722 <https://doi.org/10.1007/s13399-021-02288-x>.
723 Jiang, Yuying, Xue Bai, Shuangmei Lang, Yuelin Zhao, Chang Liu, and Lei Yu. 2019.
724 “Optimization of Ultrasonic-Microwave Assisted Alkali Extraction of Arabinoxylan from the
725 Corn Bran Using Response Surface Methodology.” *International Journal of Biological*

- 726 *Macromolecules* 128: 452–58. <https://doi.org/10.1016/j.ijbiomac.2019.01.138>.
- 727 Kaur, Amritpal, Balwinder Singh, Madhav P. Yadav, Seerat Bhinder, and Narpinder Singh. 2021.
- 728 “Isolation of Arabinoxylan and Cellulose-Rich Arabinoxylan from Wheat Bran of Different
729 Varieties and Their Functionalities.” *Food Hydrocolloids* 112 (June 2020): 106287.
730 <https://doi.org/10.1016/j.foodhyd.2020.106287>.
- 731 Kiszonas, Alecia M., E. Patrick Fuerst, Devanand Luthria, and Craig F. Morris. 2015.
- 732 “Arabinoxylan Content and Characterisation throughout the Bread-Baking Process.”
733 *International Journal of Food Science and Technology* 50 (8): 1911–21.
734 <https://doi.org/10.1111/ijfs.12829>.
- 735 Langenaeken, Niels A., David P. De Schutter, and Christophe M. Courtin. 2020. “Arabinoxylan
736 from Non-Malted Cereals Can Act as Mouthfeel Contributor in Beer.” *Carbohydrate Polymers*
737 239 (December 2019): 116257. <https://doi.org/10.1016/j.carbpol.2020.116257>.
- 738 Lapierre, Catherine, Aline Voxeur, Steven D Karlen, Richard F Helm, and John Ralph. 2018.
- 739 “Evaluation of Feruloylated and P-Coumaroylated Arabinosyl Units in Grass Arabinoxylans
740 by Acidolysis in Dioxane/Methanol.” *Journal of Agricultural and Food Chemistry* 66 (21):
741 5418–24. <https://doi.org/10.1021/acs.jafc.8b01618>.
- 742 Lazaridou, Athina, Tricia Chornick, Costas G. Biliaderis, and Marta S. Izydorczyk. 2008.
- 743 “Sequential Solvent Extraction and Structural Characterization of Polysaccharides from the
744 Endosperm Cell Walls of Barley Grown in Different Environments.” *Carbohydrate Polymers*
745 73 (4): 621–39. <https://doi.org/10.1016/j.carbpol.2008.01.001>.
- 746 Lebesi, Dimitra M., and Constantina Tzia. 2011. “Effect of the Addition of Different Dietary Fiber
747 and Edible Cereal Bran Sources on the Baking and Sensory Characteristics of Cupcakes.”
748 *Food and Bioprocess Technology* 4 (5): 710–22. <https://doi.org/10.1007/s11947-009-0181-3>.
- 749 Li, Cheng, Li Wang, Zhengxing Chen, Yongfu Li, and Juan Li. 2020. “Facile and Green
750 Preparation of Diverse Arabinoxylan Hydrogels from Wheat Bran by Combining Subcritical
751 Water and Enzymatic Crosslinking.” *Carbohydrate Polymers* 241 (December 2019): 116317.

752 <https://doi.org/10.1016/j.carbpol.2020.116317>.

753 Li, Jie, and Jinhua Du. 2019. "Molecular Characterization of Arabinoxylan from Wheat Beer, Beer
754 Foam and Defoamed Beer." *Molecules* 24 (7). <https://doi.org/10.3390/molecules24071230>.

755 Lihua, C., W. Xin, and C. Jiayang. 2018. "Comparison of Crude Polysaccharides Extraction from
756 Marine Diatom under Different Methods." *Acta Agriculturae Zhejiangensis* 30: 280.

757 Liu, Yan, Shaowei Wang, Ji Kang, Nifei Wang, Meng Xiao, Zhenjing Li, Changlu Wang, Qingbin
758 Guo, and Xinzhong Hu. 2020. "Arabinoxylan from Wheat Bran: Molecular Degradation and
759 Functional Investigation." *Food Hydrocolloids* 107 (March): 105914.
760 <https://doi.org/10.1016/j.foodhyd.2020.105914>.

761 Liu, Zhenbin, Min Zhang, Bhesh Bhandari, and Chaohui Yang. 2018. "Impact of Rheological
762 Properties of Mashed Potatoes on 3D Printing." *Journal of Food Engineering* 220: 76–82.
763 <https://doi.org/10.1016/j.jfoodeng.2017.04.017>.

764 Lynch, Kieran M., Conall R. Strain, Crystal Johnson, Dhrati Patangia, Catherine Stanton, Fatma
765 Koc, Jorge Gil-Martinez, et al. 2021. "Extraction and Characterisation of Arabinoxylan from
766 Brewers Spent Grain and Investigation of Microbiome Modulation Potential." *European
767 Journal of Nutrition* 60 (8): 4393–4411. <https://doi.org/10.1007/s00394-021-02570-8>.

768 Michiels, P., Delputte, N., Debyser, W., Langenaeken, N. A., & Courtin, C. M. (2023). The
769 occurrence and structural heterogeneity of arabinoxylan in commercial pilsner beers and their
770 non-alcoholic counterparts. *Carbohydrate Polymers*, 306, 120597.

771 Michniewicz, J, G G Biliaderis, and W Bushuk. 1992. "Effect of Added Pentosans on Some
772 Properties of Wheat Bread." *Food Chemistry* 43 (4): 251–57. [https://doi.org/10.1016/0308-8146\(92\)90208-J](https://doi.org/10.1016/0308-8146(92)90208-J).

773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

4 Molina, María Teresa, Lisa Lamothe, Deniz Z. Gunes, Sandra M. Vaz, and Pedro Bouchon. 2021.
5 "The Effect of Arabinoxylan and Wheat Bran Incorporation on Dough Rheology and Thermal
6 Processing of Rotary-Moulded Biscuits." *Foods* 10 (10).
7 <https://doi.org/10.3390/foods10102335>.

- 778 Montoya, Juliana, Jorge Medina, Arturo Molina, Janet Gutiérrez, Blanca Rodríguez, and Rodrigo
779 Marín. 2021. “Impact of Viscoelastic and Structural Properties from Starch-Mango and Starch-
780 Arabinoxylans Hydrocolloids in 3D Food Printing.” *Additive Manufacturing* 39 (January).
781 <https://doi.org/10.1016/j.addma.2021.101891>.
782 Moza, Jessy, and Hardeep Singh Gujral. 2017. “Influence of Barley Non-Starchy Polysaccharides
783 on Selected Quality Attributes of Sponge Cakes.” *Lwt* 85: 252–61.
784 <https://doi.org/10.1016/j.lwt.2017.07.024>.
785 Ngqangashe, Y., K. Cullerton, S. Phulkerd, C. Schneider, A. Thow, and S. Friel. 2022. “Discursive
786 Framing in Policies for Restricting the Marketing of Food and Non-Alcoholic Beverages.”
787 *Food Policy* 109: 102270.
788 Nie, Qixing, Haihong Chen, Jielun Hu, He Gao, Linlin Fan, Zhimin Long, and Shaoping Nie. 2018.
789 “Arabinoxylan Attenuates Type 2 Diabetes by Improvement of Carbohydrate , Lipid and
790 Amino Acid Metabolism,” 1–38. <https://doi.org/10.1002/mnfr.201800222>.
791 Oliete, Bonastre, Gabriela T. Pérez, Manuel Gómez, Pablo D. Ribotta, Malena Moiraghi, and
792 Alberto E. León. 2010. “Use of Wheat, Triticale and Rye Flours in Layer Cake Production.”
793 *International Journal of Food Science and Technology* 45 (4): 697–706.
794 <https://doi.org/10.1111/j.1365-2621.2010.02183.x>.
795 Paesani, C., E. Salvucci, M. Moiraghi, L. Fernandez Canigia, and G. T. Pérez. 2019. “Arabinoxylan
796 from Argentinian Whole Wheat Flour Promote the Growth of Lactobacillus Reuteri and
797 Bifidobacterium Breve.” *Letters in Applied Microbiology* 68 (2): 142–48.
798 <https://doi.org/10.1111/lam.13097>.
799 Paesani, Candela, Alicia Laura Degano, Emiliano Salvucci, Maria Ines Zalosnik, João Paulo Fabi,
800 Lorena Susana Sciarini, and Gabriela Teresa Perez. 2020. “Soluble Arabinoxylans Extracted
801 from Soft and Hard Wheat Show a Differential Prebiotic Effect in Vitro and in Vivo.” *Journal*
802 *of Cereal Science* 93 (March). <https://doi.org/10.1016/j.jcs.2020.102956>.
803 Pareyt, Bram, Marijke Goovaerts, Willem F. Broekaert, and Jan A. Delcour. 2011. “Arabinoxylan

- 804 Oligosaccharides (AXOS) as a Potential Sucrose Replacer in Sugar-Snap Cookies.” *Lwt* 44
805 (3): 725–28. <https://doi.org/10.1016/j.lwt.2010.11.008>.
- 806 Paz-Samaniego, Rita, Elizabeth Carvajal-Millan, Norberto Sotelo-Cruz, Francisco Brown, Agustín
807 Rascón-Chu, Yolanda L. López-Franco, and Jaime Lizardi-Mendoza. 2016. “Maize Processing
808 Waste Water Upcycling in Mexico: Recovery of Arabinoxylans for Probiotic Encapsulation.”
809 *Sustainability (Switzerland)* 8 (11). <https://doi.org/10.3390/su8111104>.
- 810 Pérez-Flores, Jesús Guadalupe, Elizabeth Contreras-López, Araceli Castañeda-Ovando, Fidel Pérez-
811 Moreno, Karina Aguilar-Arteaga, Giaan A. Álvarez-Romero, and Alejandro Téllez-Jurado.
812 2019. “Physicochemical Characterization of an Arabinoxylan-Rich Fraction from Brewers’
813 Spent Grain and Its Application as a Release Matrix for Caffeine.” *Food Research
814 International* 116 (September 2018): 1020–30. <https://doi.org/10.1016/j.foodres.2018.09.041>.
- 815 Ramírez-Jiménez, Aurea K., and Roberto Castro-Muñoz. 2020. “Emerging Techniques Assisting
816 Nixtamalization Products and By-Products Processing: An Overview.” *Critical Reviews in
817 Food Science and Nutrition* 0 (0): 1–14. <https://doi.org/10.1080/10408398.2020.1798352>.
- 818 Rudjito, Reskandi C, Andrea C Ruthes, Amparo Jiménez-Quero, and Francisco Vilaplana. 2019.
819 “Feruloylated Arabinoxylans from Wheat Bran: Optimization of Extraction Process and
820 Validation at Pilot Scale.” *ACS Sustainable Chemistry & Engineering* 7 (15): 13167–77.
821 <https://doi.org/10.1021/acssuschemeng.9b02329>.
- 822 Sánchez-Bastardo, Nuria, Alberto Romero, and Esther Alonso. 2017. “Extraction of Arabinoxylans
823 from Wheat Bran Using Hydrothermal Processes Assisted by Heterogeneous Catalysts.”
824 *Carbohydrate Polymers* 160: 143–52. <https://doi.org/10.1016/j.carbpol.2016.12.035>.
- 825 Sadosky, P., Schwarz, P. B., & Horsley, R. D. (2002). Effect of arabinoxylans, β -glucans, and
826 dextrans on the viscosity and membrane filterability of a beer model solution. *Journal of the
827 American Society of Brewing Chemists*, 60(4), 153-162.
- 828 Sarker, Niloy C., Priyanka Ray, Creighton Pfau, Venu Kalavacharla, Khwaja Hossain, and
829 Mohiuddin Quadir. 2020. “Development of Functional Nanomaterials from Wheat Bran

830 Derived Arabinoxylan for Nucleic Acid Delivery.” *Journal of Agricultural and Food*
 831 *Chemistry* 68 (15): 4367–73. <https://doi.org/10.1021/acs.jafc.0c00029>.

832 Schmidt, Marcus, Berthold Wiege, and Jürgen Hollmann. 2021. “Comparison of Alkaline/Oxidative
 833 and Hydrothermal Extraction of Wheat Bran Arabinoxylans.” *Foods*, 826.

834 Schneider, Petra, Vincent Rochell, Kay Plat, and Alexander Jaworski. 2021. “Circular Approaches
 835 in Small-Scale Food Production.” *Circular Economy and Sustainability* 1 (4): 1231–55.
 836 <https://doi.org/10.1007/s43615-021-00129-7>.

837 Schneiderbanger, J., Grammer, M., Jacob, F., Hutzler, M., 2016. Statistical evaluation of beer
 838 spoilage bacteria by real-time PCR analyses from 2010 to 2016. *J. Inst. Brew.* 124 (2), 173–
 839 181. <https://doi.org/10.1002/jib.486>

840 Schupfer, Emily, Sok Cheon Pak, Shaoyu Wang, Peter S. Micalos, Thomas Jeffries, Soo Liang Ooi,
 841 Terry Golombick, Garth Harris, and Emad El-Omar. 2021. “The Effects and Benefits of
 842 Arabinoxylans on Human Gut Microbiota – A Narrative Review.” *Food Bioscience* 43
 843 (October 2020): 101267. <https://doi.org/10.1016/j.fbio.2021.101267>.

844 Siddiqui, Shahida Anusha, Nur Alim Bahmid, Ahmed Taha, Abdel-Moneim Eid Abdel-Moneim,
 845 Abdelrazeq M Shehata, Chen Tan, Mohammad Saeed Kharazmi, et al. 2022. “Bioactive-
 846 Loaded Nanodelivery Systems for the Feed and Drugs of Livestock; Purposes, Techniques and
 847 Applications.” *Advances in Colloid and Interface Science* 308: 102772.
 848 <https://doi.org/10.1016/j.cis.2022.102772>.

849 Solier, Yamil Nahún, Paulina Mocchiutti, María Noel Cabrera, Mario Carlos Nazareno Saparrat,
 850 Miguel Ángel Zanuttini, and María Cristina Inalbon. 2022. “Alkali-Peroxide Treatment of
 851 Sugar Cane Bagasse. Effect of Chemical Charges on the Efficiency of Xylan Isolation and
 2 Susceptibility of Bagasse to Saccharification.” *Biomass Conversion and Biorefinery* 12 (3):
 3 567–76. <https://doi.org/10.1007/s13399-020-00776-0>.

4 Solomou, Konstantina, Mohammad Alyassin, Athanasios Angelis-Dimakis, and Grant M.
 5 Campbell. 2022. “Arabinoxylans: A New Class of Food Ingredients Arising from Synergies

856 with Biorefining, and Illustrating the Nature of Biorefinery Engineering.” *Food and*
857 *Bioproducts Processing* 132: 83–98. <https://doi.org/10.1016/j.fbp.2021.12.007>.

858 Sui, Wenjie, Xuan Xie, Rui Liu, Tao Wu, and Min Zhang. 2018. “Effect of Wheat Bran
859 Modification by Steam Explosion on Structural Characteristics and Rheological Properties of
860 Wheat Flour Dough.” *Food Hydrocolloids* 84: 571–80.
861 <https://doi.org/10.1016/j.foodhyd.2018.06.027>.

862 Valencia-Arredondo, Jesús Antonio, Gloria Ivonne Hernández-Bolio, Genaro Iván Cerón-Montes,
863 Roberto Castro-Muñoz, and Jorge Yáñez-Fernández. 2020. “Enhanced Process Integration for
864 the Extraction, Concentration and Purification of Di-Acylated Cyanidin from Red Cabbage.”
865 *Separation and Purification Technology* 238 (October 2019): 116492.
866 <https://doi.org/10.1016/j.seppur.2019.116492>.

867 Wang, M., R. J. Hamer, T. Van Vliet, H. Gruppen, H. Marseille, and P. L. Weegels. 2003. “Effect
868 of Water Unextractable Solids on Gluten Formation and Properties: Mechanistic
869 Considerations.” *Journal of Cereal Science* 37 (1): 55–64.
870 <https://doi.org/10.1006/jcrs.2002.0478>.

871 Xue, Y., Xibing Cui, Zonghui Zhang, Tao Zhou, Rui Gao, Yaxian Li, and Xiaoxuan Ding. 2020.
872 “Effect of β -Endoxylanase and α -Arabinofuranosidase Enzymatic Hydrolysis on Nutritional
873 and Technological Properties of Wheat Brans.” *Food Chemistry* 302 (August 2019): 125332.
874 <https://doi.org/10.1016/j.foodchem.2019.125332>.

875 Yilmaz-Turan, Secil, Amparo Jiménez-Quero, Carolin Menzel, Danila Morais de Carvalho, Mikael
876 E. Lindström, Olena Sevastyanova, Rosana Moriana, and Francisco Vilaplana. 2020. “Bio-
877 Based Films from Wheat Bran Feruloylated Arabinoxylan: Effect of Extraction Technique,
878 Acetylation and Feruloylation.” *Carbohydrate Polymers* 250 (April).
879 <https://doi.org/10.1016/j.carbpol.2020.116916>.

880 Zannini, Emanuele, Ángela Bravo Núñez, Aylin W. Sahin, and Elke K. Arendt. 2022.
881 “Arabinoxylans as Functional Food Ingredients: A Review.” *Foods* 11 (7): 1–28.

882 <https://doi.org/10.3390/foods11071026>.

883 Zhang, Ying, Leena Pitkänen, Jacques Douglade, Maija Tenkanen, Caroline Remond, and

884 Catherine Joly. 2011. "Wheat Bran Arabinoxylans: Chemical Structure and Film Properties of

885 Three Isolated Fractions." *Carbohydrate Polymers* 86 (2): 852–59.

886 <https://doi.org/10.1016/j.carbpol.2011.05.036>.

887 Zhao, Chuang, Bing Liu, Shilong Piao, Xuhui Wang, David B. Lobell, Yao Huang, Mengtian

888 Huang, et al. 2017. "Temperature Increase Reduces Global Yields of Major Crops in Four

889 Independent Estimates." *Proceedings of the National Academy of Sciences of the United States*

890 *of America* 114 (35): 9326–31. <https://doi.org/10.1073/pnas.1701762114>.

891 Zhao, Yimin, Jianhui Liu, Wangjun Hao, Hanyue Zhu, Ning Liang, Zouyan He, Ka Ying Ma, and

892 Zhen Yu Chen. 2017. "Structure-Specific Effects of Short-Chain Fatty Acids on Plasma

893 Cholesterol Concentration in Male Syrian Hamsters." *Journal of Agricultural and Food*

894 *Chemistry* 65 (50): 10984–92. <https://doi.org/10.1021/acs.jafc.7b04666>.

895 Zilliox, Caroline, and Philippe Debeire. 1998. "Hydrolysis of Wheat Straw by a Thermostable

896 Endoxylanase: Adsorption and Kinetic Studies." *Enzyme and Microbial Technology* 22 (1):

897 58–63. [https://doi.org/10.1016/S0141-0229\(97\)00105-1](https://doi.org/10.1016/S0141-0229(97)00105-1).

898