

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(diethylamido- κ N)(diethylamine- κ N)-bis(2,6-diisopropylphenylamido- κ N)-zirconium(IV)

Mateusz Zauliczny, Rafał Grubba, Łukasz Ponikiewski* and Jerzy Pikies

Chemical Faculty, Gdansk University of Technology, Gabriela Narutowicza Street 11/12, 80-233 Gdansk, Poland

Correspondence e-mail: lukasz.ponikiewski@pg.gda.pl


Received 4 December 2012; accepted 18 December 2012

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.073; wR factor = 0.188; data-to-parameter ratio = 18.4.

In the title compound, $[\text{Zr}(\text{C}_{12}\text{H}_{18}\text{N})_2(\text{C}_4\text{H}_{10}\text{N})_2(\text{C}_4\text{H}_{11}\text{N})]$ or $[\text{Zr}(\text{HNC}_6\text{H}_3^i\text{Pr}_2)_2(\text{NEt}_2)_2(\text{HNEt}_2)]$, which was obtained by the reaction of $\text{Zr}(\text{NEt})_4$ with $^i\text{Pr}_2\text{C}_6\text{H}_3\text{NH}_2$, the Zr^{IV} atom is in a trigonal-bipyramidal geometry in which the N atoms from two $^i\text{Pr}_2\text{C}_6\text{H}_3\text{NH}$ and one NEt_2 ligand occupy the equatorial positions, and the N atoms of an NEt_2 and an Et_2NH ligand occupy the apical positions. An intramolecular $\text{N}-\text{H}\cdots\text{N}$ contact occurs. There are two independent molecules in the asymmetric unit.

Related literature

For related zirconium(IV) structures, see: Profflet *et al.* (1990); Blake *et al.* (1997); Porter & Danopoulos (2004); Ghesner *et al.* (2006). For related syntheses, see: Kempe (2000).



Experimental

Crystal data

$[\text{Zr}(\text{C}_{12}\text{H}_{18}\text{N})_2(\text{C}_4\text{H}_{10}\text{N})_2(\text{C}_4\text{H}_{11}\text{N})]$
 $M_r = 661.17$
 Triclinic, $P\bar{1}$
 $a = 11.2079$ (3) Å
 $b = 13.1612$ (5) Å
 $c = 14.3443$ (6) Å
 $\alpha = 86.578$ (3)°
 $\beta = 70.484$ (3)°
 $\gamma = 71.232$ (3)°
 $V = 1885.61$ (12) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.32$ mm⁻¹
 $T = 293$ K
 $0.58 \times 0.39 \times 0.34$ mm

Data collection

Agilent Xcalibur (Sapphire2) diffractometer
 Absorption correction: analytical (*CrysAlis PRO*; Agilent, 2010)
 $T_{\min} = 0.889$, $T_{\max} = 0.928$
 11493 measured reflections
 7412 independent reflections
 5867 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.035$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.073$
 $wR(F^2) = 0.188$
 $S = 1.12$
 7412 reflections
 402 parameters
 1 restraint
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 1.82$ e Å⁻³
 $\Delta\rho_{\min} = -2.13$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N3}-\text{H3A}\cdots\text{N1}$	0.84 (5)	2.56 (5)	2.983 (5)	112 (4)

Data collection: *CrysAlis PRO* (Agilent, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SUPERFLIP* (Palatinus & Chapuis, 2007); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5310).

References

- Agilent (2010). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
 Blake, A. J., Nikonov, G. I. & Mountford, P. (1997). *Acta Cryst.* **C53**, 874–876.
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
 Ghesner, I., Fenwick, A. & Stephen, D. W. (2006). *Organometallics*, **25**, 4985–4995.
 Kempe, R. (2000). *Angew. Chem. Int. Ed.* **39**, 468–493.
 Palatinus, L. & Chapuis, G. (2007). *J. Appl. Cryst.* **40**, 786–790.
 Porter, R. M. & Danopoulos, A. A. (2004). *Dalton Trans.* pp. 2556–2562.
 Profflet, R. D., Zabrano, C. H., Fanwick, P. E., Nash, J. J. & Rothwell, I. P. (1990). *Inorg. Chem.* **29**, 4362–4364.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2013). E69, m72 [doi:10.1107/S160053681205115X]

Bis(diethylamido- κ N)(diethylamine- κ N)bis(2,6-diisopropylphenylamido- κ N)zirconium(IV)

Mateusz Zauliczny, Rafał Grubba, Łukasz Ponikiewski and Jerzy Pikies

S1. Comment

Complex (I) was synthesized in the course of our studies on amido complexes of zirconium (Kempe, 2000). The compound was obtained in the reaction $\text{Zr}(\text{NEt}_2)_4$ with ${}^i\text{Pr}_2\text{C}_6\text{H}_3\text{NH}_2$ (molar ratio 1:2). Complex (I) contains four amido ligands (two NEt_2 and two ${}^i\text{Pr}_2\text{C}_6\text{H}_3\text{NH}$) and one amino ligand (HNEt_2). Analyzing bond lengths in title compound it is easily spotted that bond $\text{Zr}-\text{N}3$ is much longer than other bonds. It is caused by the fact, that ligand containing $\text{N}1$ is amine ligand. Difference between length of bonds $\text{Zr}-\text{N}$ between NEt_2 and HNEt_2 is about 0.25 Å. Distances between N atoms of ${}^i\text{Pr}_2\text{C}_6\text{H}_3\text{NH}$ and Zr are both about 2.12 Å and very similar to related $\text{Zr}(\text{IV})$ amido complexes (Profilet *et al.*, 1990; Blake *et al.*, 1997). In case of two NEt_2 ligands distances $\text{Zr}-\text{N}$ differ by about 0.12 Å but both are in the range typical for zirconium complexes with diethylamido ligands (Porter *et al.*, 2004; Ghesner *et al.*, 2006). Comparing angles between $\text{N}1-\text{Zr}1-\text{N}2$, $\text{N}2-\text{Zr}1-\text{N}5$ and $\text{N}5-\text{Zr}1-\text{N}1$ [table 1] it can be seen that they are roughly 120°, in addition the angle between $\text{N}3-\text{Zr}1-\text{N}4$ [table 1] indicates that molecular geometry is close to trigonal bipyramidal. Admittedly in perfect trigonal bipyramid first three angles would be equal to 120° and $\text{N}3-\text{Zr}1-\text{N}4$ would be equal to 180°, but actual angles are so close to said values that it is safe to say they resemble trigonal bipyramid.

The crystal packing diagram shows, that the compound crystallizes with two molecules in the unit cell in the triclinic space group. The crystal packing of the title compound is presented in Fig.2.

S2. Experimental

To a 100 ml Schlenk flask, equipped with a magnetic stirrer, charged with a solution of 4,8 g (4,44 ml) $\text{Zr}(\text{NEt}_2)_4$ in 30 ml of pentane, 5 g (4,76 ml) of 2,6-diisopropylaniline in 10 ml of pentane was added dropwise. The reaction was carried on in a room temperature in an argon atmosphere. Solution was left on a magnetic stirrer. Over a week of stirring, the mixture changed colour from yellow to brown. The solvent was removed under vacuum. After evaporating most of the solvent, dark solid was obtained which after keeping it longer under vacuum got darker and became oil. Residue was dissolved in 8 ml of pentane, and recrystallized at 4°C to obtain about 2 g of colorless X-ray-quality crystals. The total yield was 24%. Elemental analysis, found %: C 65.73, H 9.72, N 10.46; calc. % for $\text{C}_{30}\text{H}_{52}\text{N}_5\text{Zr}$: C 65.40, H 10.21, N 10.59.

S3. Refinement

The C—H H atoms were positioned with idealized geometry and were refined isotropically with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic, methylene and methine H atoms (1.5 for methyl H atoms) using a riding model with C—H = 0.93 Å (aromatic H atoms), 0.96 Å (methyl H atoms), 0.97 Å (methylene H atoms) and 0.98 Å (methine H atoms). The amine hydrogen atoms were located in the difference Fourier map and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$. The hydrogen atom H1A is located in the difference map and restrained, $\text{N}1-\text{H}1\text{A} = 0.89$ Å with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$. The

highest residual electron density peaks are located within 1 Å from atom Zr1 and the deepest hole is located 0.78 Å from Zr1.




Figure 1

The molecular structure of the title molecule with the atom-numbering. Displacement ellipsoids are drawn at the 30% probability level H atoms connected to C have been omitted.




Figure 2

A view of the packing of the title compound along the *a* axis.

Bis(diethylamido- κ N)(diethylamine- κ N)bis(2,6-diisopropylphenylamido- κ N)zirconium(IV)

Crystal data

$[\text{Zr}(\text{C}_{12}\text{H}_{18}\text{N})_2(\text{C}_4\text{H}_{10}\text{N})_2(\text{C}_4\text{H}_{11}\text{N})]$

$M_r = 661.17$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 11.2079$ (3) Å

$b = 13.1612$ (5) Å

$c = 14.3443$ (6) Å

$\alpha = 86.578$ (3)°

$\beta = 70.484$ (3)°

$\gamma = 71.232$ (3)°

$V = 1885.61$ (12) Å³

$Z = 2$

$F(000) = 716$

$D_x = 1.164$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 6954 reflections

$\theta = 2.9\text{--}28.4^\circ$

$\mu = 0.32$ mm⁻¹

$T = 293$ K

Block, colourless

$0.58 \times 0.39 \times 0.34$ mm

Data collection

Agilent Xcalibur (Sapphire2
diffractometer

Graphite monochromator

Detector resolution: 8.1883 pixels mm⁻¹

ω scans

Absorption correction: analytical
(*CrysAlis PRO*; Agilent, 2010)

$T_{\min} = 0.889$, $T_{\max} = 0.928$

11493 measured reflections

7412 independent reflections

5867 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.035$

$\theta_{\max} = 26^\circ$, $\theta_{\min} = 2.9^\circ$

$h = -13 \rightarrow 8$

$k = -16 \rightarrow 12$

$l = -17 \rightarrow 12$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.073$
 $wR(F^2) = 0.188$
 $S = 1.12$
 7412 reflections
 402 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0694P)^2 + 9.6553P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 1.82 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -2.13 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zr1	0.73585 (6)	0.79022 (4)	0.21365 (4)	0.03188 (17)
N1	0.7260 (3)	0.8397 (3)	0.3554 (3)	0.0216 (8)
H1A	0.650 (3)	0.838 (4)	0.401 (3)	0.026*
N2	0.7770 (4)	0.6310 (3)	0.1615 (3)	0.0199 (8)
H2	0.856 (5)	0.591 (4)	0.160 (4)	0.024*
N3	0.9717 (4)	0.7334 (3)	0.1861 (3)	0.0244 (8)
H3A	0.963 (5)	0.779 (4)	0.229 (4)	0.029*
N4	0.5246 (4)	0.8112 (3)	0.2736 (3)	0.0277 (9)
N5	0.7378 (3)	0.9016 (3)	0.1088 (3)	0.0197 (7)
C1	0.8144 (4)	0.8339 (3)	0.4082 (3)	0.0196 (9)
C2	0.8987 (4)	0.8990 (4)	0.3821 (3)	0.0207 (9)
C3	0.9889 (5)	0.8903 (4)	0.4321 (4)	0.0285 (10)
H3	1.0441	0.9331	0.4152	0.034*
C4	0.9975 (5)	0.8195 (4)	0.5060 (4)	0.0320 (11)
H4	1.0588	0.8141	0.5381	0.038*
C5	0.9145 (5)	0.7562 (4)	0.5325 (4)	0.0306 (11)
H5	0.9202	0.7093	0.5831	0.037*
C6	0.8225 (4)	0.7616 (4)	0.4847 (3)	0.0248 (10)
C7	0.8854 (4)	0.9830 (4)	0.3060 (3)	0.0242 (9)
H7	0.8631	0.9538	0.2552	0.029*
C8	0.7667 (5)	1.0843 (4)	0.3554 (4)	0.0349 (12)
H8A	0.6871	1.0649	0.3849	0.052*
H8B	0.7539	1.1357	0.3064	0.052*
H8C	0.7855	1.1153	0.4057	0.052*

C9	1.0102 (5)	1.0135 (5)	0.2538 (4)	0.0367 (12)
H9A	1.0316	1.0469	0.301	0.055*
H9B	0.9947	1.0627	0.2037	0.055*
H9C	1.0833	0.95	0.2236	0.055*
C10	0.7317 (5)	0.6931 (4)	0.5149 (4)	0.0277 (10)
H10	0.7206	0.672	0.4547	0.033*
C11	0.5927 (5)	0.7584 (5)	0.5843 (4)	0.0404 (13)
H11A	0.5981	0.7717	0.6475	0.061*
H11B	0.5318	0.7186	0.5925	0.061*
H11C	0.5612	0.8256	0.5561	0.061*
C12	0.7852 (6)	0.5903 (5)	0.5637 (5)	0.0453 (14)
H12A	0.871	0.5482	0.52	0.068*
H12B	0.7244	0.5496	0.5773	0.068*
H12C	0.7937	0.6083	0.6246	0.068*
C13	0.6972 (4)	0.5653 (3)	0.1692 (3)	0.0192 (9)
C14	0.7022 (4)	0.4784 (4)	0.2323 (3)	0.0226 (9)
C15	0.6217 (5)	0.4149 (4)	0.2379 (3)	0.0282 (10)
H15	0.6266	0.357	0.2784	0.034*
C16	0.5353 (5)	0.4361 (4)	0.1847 (4)	0.0306 (11)
H16	0.4813	0.3937	0.1899	0.037*
C17	0.5298 (5)	0.5218 (4)	0.1233 (3)	0.0261 (10)
H17	0.4718	0.5357	0.087	0.031*
C18	0.6073 (4)	0.5871 (3)	0.1142 (3)	0.0193 (9)
C19	0.7952 (5)	0.4533 (4)	0.2924 (3)	0.0267 (10)
H19	0.8076	0.5209	0.3049	0.032*
C20	0.7410 (6)	0.4094 (4)	0.3930 (4)	0.0375 (12)
H20A	0.6539	0.4573	0.4286	0.056*
H20B	0.8	0.4035	0.43	0.056*
H20C	0.7351	0.3398	0.3836	0.056*
C21	0.9322 (5)	0.3769 (5)	0.2345 (4)	0.0390 (12)
H21A	0.9246	0.3085	0.2232	0.058*
H21B	0.9916	0.3682	0.2716	0.058*
H21C	0.9668	0.406	0.172	0.058*
C22	0.6044 (4)	0.6763 (3)	0.0418 (3)	0.0210 (9)
H22	0.6282	0.7316	0.0675	0.025*
C23	0.4647 (5)	0.7303 (4)	0.0324 (4)	0.0306 (11)
H23A	0.4414	0.6797	0.0022	0.046*
H23B	0.4661	0.7911	-0.008	0.046*
H23C	0.3999	0.7538	0.0969	0.046*
C24	0.7096 (5)	0.6349 (4)	-0.0588 (3)	0.0294 (10)
H24A	0.7965	0.6084	-0.0518	0.044*
H24B	0.7071	0.6924	-0.1033	0.044*
H24C	0.6918	0.578	-0.0848	0.044*
C25	1.0286 (4)	0.6345 (4)	0.2313 (3)	0.0251 (10)
H25A	1.0531	0.5737	0.1859	0.03*
H25B	0.9602	0.6261	0.2911	0.03*
C26	1.1508 (5)	0.6316 (5)	0.2573 (4)	0.0380 (12)
H26A	1.2205	0.6373	0.1982	0.057*

H26B	1.1814	0.5652	0.2868	0.057*
H26C	1.1274	0.6907	0.3033	0.057*
C27	1.0646 (4)	0.7550 (4)	0.0928 (3)	0.0239 (9)
H27A	1.1431	0.7593	0.1047	0.029*
H27B	1.0217	0.8239	0.0707	0.029*
C28	1.1072 (7)	0.6691 (5)	0.0121 (4)	0.0509 (16)
H28A	1.1663	0.6043	0.027	0.076*
H28B	1.1527	0.6931	-0.0502	0.076*
H28C	1.0298	0.6555	0.0081	0.076*
C29	0.4813 (5)	0.7500 (4)	0.3598 (3)	0.0282 (10)
H29A	0.4477	0.7966	0.419	0.034*
H29B	0.5585	0.6917	0.3636	0.034*
C30	0.3737 (5)	0.7030 (5)	0.3591 (4)	0.0366 (12)
H30A	0.2926	0.7603	0.364	0.055*
H30B	0.3577	0.6583	0.4143	0.055*
H30C	0.403	0.6608	0.2985	0.055*
C31	0.4177 (4)	0.9023 (4)	0.2580 (4)	0.0292 (10)
H31A	0.3505	0.8756	0.2497	0.035*
H31B	0.4546	0.9342	0.1969	0.035*
C32	0.3507 (8)	0.9887 (5)	0.3406 (5)	0.067 (2)
H32A	0.3025	0.9608	0.3992	0.101*
H32B	0.2898	1.0489	0.3214	0.101*
H32C	0.4173	1.0112	0.3536	0.101*
C33	0.7937 (4)	0.8757 (4)	0.0015 (3)	0.0268 (10)
H33A	0.8264	0.7982	-0.0097	0.032*
H33B	0.8697	0.9015	-0.0258	0.032*
C34	0.6955 (5)	0.9233 (4)	-0.0547 (4)	0.0344 (12)
H34A	0.6159	0.9041	-0.0243	0.052*
H34B	0.736	0.8955	-0.1223	0.052*
H34C	0.6729	1.0002	-0.0527	0.052*
C35	0.6795 (5)	1.0173 (4)	0.1362 (4)	0.0297 (10)
H35A	0.6508	1.027	0.2077	0.036*
H35B	0.6006	1.0457	0.1166	0.036*
C36	0.7729 (6)	1.0832 (4)	0.0903 (4)	0.0405 (13)
H36A	0.8553	1.0516	0.1035	0.061*
H36B	0.7313	1.1555	0.1186	0.061*
H36C	0.7906	1.0839	0.0201	0.061*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zr1	0.0557 (4)	0.0191 (2)	0.0234 (3)	-0.0121 (2)	-0.0168 (2)	0.00391 (17)
N1	0.0115 (17)	0.028 (2)	0.0236 (19)	-0.0066 (15)	-0.0036 (14)	-0.0005 (16)
N2	0.0132 (18)	0.0206 (19)	0.0262 (19)	-0.0045 (15)	-0.0079 (15)	0.0025 (15)
N3	0.0156 (19)	0.0203 (19)	0.035 (2)	-0.0068 (15)	-0.0042 (16)	-0.0001 (16)
N4	0.0171 (19)	0.035 (2)	0.025 (2)	-0.0036 (17)	-0.0046 (15)	0.0012 (17)
N5	0.0151 (18)	0.0180 (18)	0.0237 (19)	-0.0038 (14)	-0.0047 (14)	-0.0006 (14)
C1	0.013 (2)	0.021 (2)	0.020 (2)	-0.0019 (17)	-0.0023 (16)	-0.0057 (17)



C2	0.011 (2)	0.025 (2)	0.022 (2)	-0.0028 (17)	-0.0021 (16)	-0.0082 (17)
C3	0.018 (2)	0.031 (3)	0.036 (3)	-0.0060 (19)	-0.0078 (19)	-0.007 (2)
C4	0.018 (2)	0.041 (3)	0.040 (3)	-0.004 (2)	-0.015 (2)	-0.008 (2)
C5	0.033 (3)	0.035 (3)	0.026 (2)	-0.007 (2)	-0.015 (2)	0.001 (2)
C6	0.023 (2)	0.024 (2)	0.022 (2)	-0.0027 (19)	-0.0049 (18)	-0.0056 (18)
C7	0.023 (2)	0.024 (2)	0.026 (2)	-0.0083 (19)	-0.0070 (18)	-0.0045 (18)
C8	0.035 (3)	0.025 (3)	0.038 (3)	-0.005 (2)	-0.008 (2)	0.000 (2)
C9	0.030 (3)	0.048 (3)	0.035 (3)	-0.023 (2)	-0.004 (2)	0.003 (2)
C10	0.025 (2)	0.029 (3)	0.029 (2)	-0.009 (2)	-0.0093 (19)	0.003 (2)
C11	0.035 (3)	0.048 (3)	0.033 (3)	-0.018 (3)	0.000 (2)	0.004 (2)
C12	0.059 (4)	0.036 (3)	0.053 (4)	-0.019 (3)	-0.031 (3)	0.013 (3)
C13	0.012 (2)	0.022 (2)	0.020 (2)	-0.0049 (17)	-0.0014 (16)	-0.0032 (17)
C14	0.018 (2)	0.025 (2)	0.023 (2)	-0.0074 (18)	-0.0032 (17)	-0.0011 (18)
C15	0.030 (3)	0.028 (2)	0.028 (2)	-0.016 (2)	-0.006 (2)	0.006 (2)
C16	0.027 (3)	0.036 (3)	0.034 (3)	-0.021 (2)	-0.006 (2)	-0.001 (2)
C17	0.023 (2)	0.033 (3)	0.025 (2)	-0.012 (2)	-0.0072 (19)	-0.0018 (19)
C18	0.015 (2)	0.020 (2)	0.019 (2)	-0.0037 (17)	-0.0023 (16)	-0.0050 (17)
C19	0.027 (2)	0.025 (2)	0.031 (3)	-0.011 (2)	-0.012 (2)	0.0074 (19)
C20	0.048 (3)	0.039 (3)	0.030 (3)	-0.017 (3)	-0.016 (2)	0.009 (2)
C21	0.033 (3)	0.042 (3)	0.038 (3)	-0.004 (2)	-0.014 (2)	0.005 (2)
C22	0.021 (2)	0.018 (2)	0.024 (2)	-0.0035 (17)	-0.0099 (18)	-0.0028 (17)
C23	0.021 (2)	0.028 (3)	0.043 (3)	-0.002 (2)	-0.016 (2)	0.001 (2)
C24	0.025 (2)	0.031 (3)	0.028 (2)	-0.007 (2)	-0.0051 (19)	0.003 (2)
C25	0.022 (2)	0.023 (2)	0.029 (2)	-0.0064 (19)	-0.0084 (19)	0.0012 (19)
C26	0.033 (3)	0.042 (3)	0.047 (3)	-0.014 (2)	-0.022 (2)	0.011 (2)
C27	0.017 (2)	0.026 (2)	0.026 (2)	-0.0059 (18)	-0.0043 (18)	0.0023 (19)
C28	0.072 (4)	0.037 (3)	0.028 (3)	-0.005 (3)	-0.007 (3)	-0.003 (2)
C29	0.020 (2)	0.043 (3)	0.020 (2)	-0.011 (2)	-0.0021 (18)	-0.003 (2)
C30	0.029 (3)	0.047 (3)	0.037 (3)	-0.020 (2)	-0.008 (2)	0.003 (2)
C31	0.015 (2)	0.033 (3)	0.037 (3)	-0.0052 (19)	-0.0066 (19)	-0.003 (2)
C32	0.077 (5)	0.040 (4)	0.057 (4)	0.001 (3)	-0.003 (4)	-0.017 (3)
C33	0.018 (2)	0.032 (3)	0.028 (2)	-0.0070 (19)	-0.0059 (18)	0.005 (2)
C34	0.036 (3)	0.038 (3)	0.033 (3)	-0.014 (2)	-0.018 (2)	0.011 (2)
C35	0.031 (3)	0.022 (2)	0.039 (3)	-0.008 (2)	-0.016 (2)	0.003 (2)
C36	0.048 (3)	0.032 (3)	0.053 (3)	-0.021 (3)	-0.026 (3)	0.017 (3)

Geometric parameters (Å, °)

Zr1—N5	2.036 (4)	C18—C22	1.519 (6)
Zr1—N2	2.122 (4)	C19—C21	1.520 (7)
Zr1—N1	2.129 (4)	C19—C20	1.524 (7)
Zr1—N4	2.160 (4)	C19—H19	0.98
Zr1—N3	2.402 (4)	C20—H20A	0.96
N1—C1	1.417 (5)	C20—H20B	0.96
N1—H1A	0.890 (2)	C20—H20C	0.96
N2—C13	1.407 (5)	C21—H21A	0.96
N2—H2	0.87 (5)	C21—H21B	0.96
N3—C25	1.470 (6)	C21—H21C	0.96

N3—C27	1.473 (6)	C22—C24	1.522 (6)
N3—H3A	0.84 (5)	C22—C23	1.546 (6)
N4—C29	1.465 (6)	C22—H22	0.98
N4—C31	1.465 (6)	C23—H23A	0.96
N5—C33	1.469 (6)	C23—H23B	0.96
N5—C35	1.470 (6)	C23—H23C	0.96
C1—C6	1.417 (6)	C24—H24A	0.96
C1—C2	1.418 (6)	C24—H24B	0.96
C2—C3	1.398 (6)	C24—H24C	0.96
C2—C7	1.515 (6)	C25—C26	1.524 (6)
C3—C4	1.376 (7)	C25—H25A	0.97
C3—H3	0.93	C25—H25B	0.97
C4—C5	1.389 (7)	C26—H26A	0.96
C4—H4	0.93	C26—H26B	0.96
C5—C6	1.399 (6)	C26—H26C	0.96
C5—H5	0.93	C27—C28	1.511 (7)
C6—C10	1.512 (7)	C27—H27A	0.97
C7—C9	1.518 (6)	C27—H27B	0.97
C7—C8	1.546 (6)	C28—H28A	0.96
C7—H7	0.98	C28—H28B	0.96
C8—H8A	0.96	C28—H28C	0.96
C8—H8B	0.96	C29—C30	1.525 (7)
C8—H8C	0.96	C29—H29A	0.97
C9—H9A	0.96	C29—H29B	0.97
C9—H9B	0.96	C30—H30A	0.96
C9—H9C	0.96	C30—H30B	0.96
C10—C12	1.525 (7)	C30—H30C	0.96
C10—C11	1.537 (7)	C31—C32	1.508 (8)
C10—H10	0.98	C31—H31A	0.97
C11—H11A	0.96	C31—H31B	0.97
C11—H11B	0.96	C32—H32A	0.96
C11—H11C	0.96	C32—H32B	0.96
C12—H12A	0.96	C32—H32C	0.96
C12—H12B	0.96	C33—C34	1.531 (6)
C12—H12C	0.96	C33—H33A	0.97
C13—C14	1.413 (6)	C33—H33B	0.97
C13—C18	1.427 (6)	C34—H34A	0.96
C14—C15	1.396 (6)	C34—H34B	0.96
C14—C19	1.511 (6)	C34—H34C	0.96
C15—C16	1.375 (7)	C35—C36	1.528 (7)
C15—H15	0.93	C35—H35A	0.97
C16—C17	1.387 (7)	C35—H35B	0.97
C16—H16	0.93	C36—H36A	0.96
C17—C18	1.378 (6)	C36—H36B	0.96
C17—H17	0.93	C36—H36C	0.96
N5—Zr1—N2	115.34 (14)	C21—C19—H19	107
N5—Zr1—N1	116.93 (14)	C20—C19—H19	107

N2—Zr1—N1	126.85 (14)	C19—C20—H20A	109.5
N5—Zr1—N4	100.60 (14)	C19—C20—H20B	109.5
N2—Zr1—N4	90.93 (15)	H20A—C20—H20B	109.5
N1—Zr1—N4	88.60 (14)	C19—C20—H20C	109.5
N5—Zr1—N3	95.87 (13)	H20A—C20—H20C	109.5
N2—Zr1—N3	83.79 (14)	H20B—C20—H20C	109.5
N1—Zr1—N3	82.12 (14)	C19—C21—H21A	109.5
N4—Zr1—N3	163.41 (14)	C19—C21—H21B	109.5
C1—N1—Zr1	138.4 (3)	H21A—C21—H21B	109.5
C1—N1—H1A	105 (3)	C19—C21—H21C	109.5
Zr1—N1—H1A	111 (3)	H21A—C21—H21C	109.5
C13—N2—Zr1	133.5 (3)	H21B—C21—H21C	109.5
C13—N2—H2	109 (3)	C18—C22—C24	110.6 (4)
Zr1—N2—H2	111 (3)	C18—C22—C23	113.1 (4)
C25—N3—C27	114.4 (4)	C24—C22—C23	111.1 (4)
C25—N3—Zr1	117.5 (3)	C18—C22—H22	107.3
C27—N3—Zr1	120.5 (3)	C24—C22—H22	107.3
C25—N3—H3A	99 (4)	C23—C22—H22	107.3
C27—N3—H3A	106 (4)	C22—C23—H23A	109.5
Zr1—N3—H3A	93 (4)	C22—C23—H23B	109.5
C29—N4—C31	114.7 (4)	H23A—C23—H23B	109.5
C29—N4—Zr1	116.7 (3)	C22—C23—H23C	109.5
C31—N4—Zr1	125.7 (3)	H23A—C23—H23C	109.5
C33—N5—C35	114.2 (4)	H23B—C23—H23C	109.5
C33—N5—Zr1	124.4 (3)	C22—C24—H24A	109.5
C35—N5—Zr1	121.4 (3)	C22—C24—H24B	109.5
C6—C1—N1	120.8 (4)	H24A—C24—H24B	109.5
C6—C1—C2	119.6 (4)	C22—C24—H24C	109.5
N1—C1—C2	119.6 (4)	H24A—C24—H24C	109.5
C3—C2—C1	119.4 (4)	H24B—C24—H24C	109.5
C3—C2—C7	119.5 (4)	N3—C25—C26	114.5 (4)
C1—C2—C7	120.9 (4)	N3—C25—H25A	108.6
C4—C3—C2	121.0 (4)	C26—C25—H25A	108.6
C4—C3—H3	119.5	N3—C25—H25B	108.6
C2—C3—H3	119.5	C26—C25—H25B	108.6
C3—C4—C5	119.9 (4)	H25A—C25—H25B	107.6
C3—C4—H4	120.1	C25—C26—H26A	109.5
C5—C4—H4	120.1	C25—C26—H26B	109.5
C4—C5—C6	121.4 (5)	H26A—C26—H26B	109.5
C4—C5—H5	119.3	C25—C26—H26C	109.5
C6—C5—H5	119.3	H26A—C26—H26C	109.5
C5—C6—C1	118.7 (4)	H26B—C26—H26C	109.5
C5—C6—C10	121.0 (4)	N3—C27—C28	112.3 (4)
C1—C6—C10	120.4 (4)	N3—C27—H27A	109.1
C2—C7—C9	115.3 (4)	C28—C27—H27A	109.1
C2—C7—C8	109.3 (4)	N3—C27—H27B	109.1
C9—C7—C8	110.1 (4)	C28—C27—H27B	109.1
C2—C7—H7	107.3	H27A—C27—H27B	107.9

C9—C7—H7	107.3	C27—C28—H28A	109.5
C8—C7—H7	107.3	C27—C28—H28B	109.5
C7—C8—H8A	109.5	H28A—C28—H28B	109.5
C7—C8—H8B	109.5	C27—C28—H28C	109.5
H8A—C8—H8B	109.5	H28A—C28—H28C	109.5
C7—C8—H8C	109.5	H28B—C28—H28C	109.5
H8A—C8—H8C	109.5	N4—C29—C30	114.9 (4)
H8B—C8—H8C	109.5	N4—C29—H29A	108.5
C7—C9—H9A	109.5	C30—C29—H29A	108.5
C7—C9—H9B	109.5	N4—C29—H29B	108.5
H9A—C9—H9B	109.5	C30—C29—H29B	108.5
C7—C9—H9C	109.5	H29A—C29—H29B	107.5
H9A—C9—H9C	109.5	C29—C30—H30A	109.5
H9B—C9—H9C	109.5	C29—C30—H30B	109.5
C6—C10—C12	114.6 (4)	H30A—C30—H30B	109.5
C6—C10—C11	110.6 (4)	C29—C30—H30C	109.5
C12—C10—C11	109.1 (4)	H30A—C30—H30C	109.5
C6—C10—H10	107.4	H30B—C30—H30C	109.5
C12—C10—H10	107.4	N4—C31—C32	114.6 (5)
C11—C10—H10	107.4	N4—C31—H31A	108.6
C10—C11—H11A	109.5	C32—C31—H31A	108.6
C10—C11—H11B	109.5	N4—C31—H31B	108.6
H11A—C11—H11B	109.5	C32—C31—H31B	108.6
C10—C11—H11C	109.5	H31A—C31—H31B	107.6
H11A—C11—H11C	109.5	C31—C32—H32A	109.5
H11B—C11—H11C	109.5	C31—C32—H32B	109.5
C10—C12—H12A	109.5	H32A—C32—H32B	109.5
C10—C12—H12B	109.5	C31—C32—H32C	109.5
H12A—C12—H12B	109.5	H32A—C32—H32C	109.5
C10—C12—H12C	109.5	H32B—C32—H32C	109.5
H12A—C12—H12C	109.5	N5—C33—C34	114.7 (4)
H12B—C12—H12C	109.5	N5—C33—H33A	108.6
N2—C13—C14	121.1 (4)	C34—C33—H33A	108.6
N2—C13—C18	120.0 (4)	N5—C33—H33B	108.6
C14—C13—C18	118.8 (4)	C34—C33—H33B	108.6
C15—C14—C13	119.5 (4)	H33A—C33—H33B	107.6
C15—C14—C19	120.0 (4)	C33—C34—H34A	109.5
C13—C14—C19	120.5 (4)	C33—C34—H34B	109.5
C16—C15—C14	121.5 (4)	H34A—C34—H34B	109.5
C16—C15—H15	119.3	C33—C34—H34C	109.5
C14—C15—H15	119.3	H34A—C34—H34C	109.5
C15—C16—C17	118.9 (4)	H34B—C34—H34C	109.5
C15—C16—H16	120.5	N5—C35—C36	115.0 (4)
C17—C16—H16	120.5	N5—C35—H35A	108.5
C18—C17—C16	122.3 (4)	C36—C35—H35A	108.5
C18—C17—H17	118.8	N5—C35—H35B	108.5
C16—C17—H17	118.8	C36—C35—H35B	108.5
C17—C18—C13	118.9 (4)	H35A—C35—H35B	107.5

C17—C18—C22	120.8 (4)	C35—C36—H36A	109.5
C13—C18—C22	120.2 (4)	C35—C36—H36B	109.5
C14—C19—C21	111.3 (4)	H36A—C36—H36B	109.5
C14—C19—C20	114.2 (4)	C35—C36—H36C	109.5
C21—C19—C20	109.9 (4)	H36A—C36—H36C	109.5
C14—C19—H19	107	H36B—C36—H36C	109.5
N5—Zr1—N1—C1	-96.8 (5)	N1—C1—C6—C10	-3.1 (6)
N2—Zr1—N1—C1	71.9 (5)	C2—C1—C6—C10	178.9 (4)
N4—Zr1—N1—C1	162.0 (5)	C3—C2—C7—C9	28.7 (6)
N3—Zr1—N1—C1	-4.2 (4)	C1—C2—C7—C9	-155.7 (4)
N5—Zr1—N2—C13	-105.4 (4)	C3—C2—C7—C8	-95.9 (5)
N1—Zr1—N2—C13	85.7 (4)	C1—C2—C7—C8	79.7 (5)
N4—Zr1—N2—C13	-3.2 (4)	C5—C6—C10—C12	-25.1 (6)
N3—Zr1—N2—C13	161.0 (4)	C1—C6—C10—C12	155.9 (4)
N5—Zr1—N3—C25	-167.4 (3)	C5—C6—C10—C11	98.6 (5)
N2—Zr1—N3—C25	-52.5 (3)	C1—C6—C10—C11	-80.3 (5)
N1—Zr1—N3—C25	76.1 (3)	Zr1—N2—C13—C14	-106.6 (4)
N4—Zr1—N3—C25	19.6 (7)	Zr1—N2—C13—C18	72.2 (5)
N5—Zr1—N3—C27	-19.4 (3)	N2—C13—C14—C15	-179.9 (4)
N2—Zr1—N3—C27	95.5 (3)	C18—C13—C14—C15	1.3 (6)
N1—Zr1—N3—C27	-135.9 (3)	N2—C13—C14—C19	-0.7 (6)
N4—Zr1—N3—C27	167.5 (4)	C18—C13—C14—C19	-179.5 (4)
N5—Zr1—N4—C29	-174.3 (3)	C13—C14—C15—C16	-1.3 (7)
N2—Zr1—N4—C29	69.7 (3)	C19—C14—C15—C16	179.5 (4)
N1—Zr1—N4—C29	-57.2 (3)	C14—C15—C16—C17	0.9 (7)
N3—Zr1—N4—C29	-1.4 (7)	C15—C16—C17—C18	-0.5 (7)
N5—Zr1—N4—C31	-14.6 (4)	C16—C17—C18—C13	0.5 (7)
N2—Zr1—N4—C31	-130.6 (4)	C16—C17—C18—C22	176.5 (4)
N1—Zr1—N4—C31	102.5 (4)	N2—C13—C18—C17	-179.8 (4)
N3—Zr1—N4—C31	158.3 (4)	C14—C13—C18—C17	-0.9 (6)
N2—Zr1—N5—C33	-14.4 (4)	N2—C13—C18—C22	4.2 (6)
N1—Zr1—N5—C33	155.6 (3)	C14—C13—C18—C22	-176.9 (4)
N4—Zr1—N5—C33	-110.5 (3)	C15—C14—C19—C21	90.7 (5)
N3—Zr1—N5—C33	71.5 (3)	C13—C14—C19—C21	-88.5 (5)
N2—Zr1—N5—C35	164.6 (3)	C15—C14—C19—C20	-34.4 (6)
N1—Zr1—N5—C35	-25.4 (4)	C13—C14—C19—C20	146.4 (4)
N4—Zr1—N5—C35	68.4 (3)	C17—C18—C22—C24	-90.1 (5)
N3—Zr1—N5—C35	-109.5 (3)	C13—C18—C22—C24	85.9 (5)
Zr1—N1—C1—C6	-105.2 (5)	C17—C18—C22—C23	35.2 (6)
Zr1—N1—C1—C2	72.9 (6)	C13—C18—C22—C23	-148.8 (4)
C6—C1—C2—C3	0.2 (6)	C27—N3—C25—C26	59.7 (5)
N1—C1—C2—C3	-177.8 (4)	Zr1—N3—C25—C26	-150.5 (3)
C6—C1—C2—C7	-175.4 (4)	C25—N3—C27—C28	63.2 (5)
N1—C1—C2—C7	6.6 (6)	Zr1—N3—C27—C28	-85.7 (5)
C1—C2—C3—C4	0.2 (7)	C31—N4—C29—C30	58.1 (6)
C7—C2—C3—C4	175.9 (4)	Zr1—N4—C29—C30	-140.0 (4)
C2—C3—C4—C5	-0.7 (7)	C29—N4—C31—C32	62.7 (6)

C3—C4—C5—C6	0.8 (7)	Zr1—N4—C31—C32	-97.4 (5)
C4—C5—C6—C1	-0.4 (7)	C35—N5—C33—C34	-56.0 (5)
C4—C5—C6—C10	-179.4 (4)	Zr1—N5—C33—C34	123.1 (4)
N1—C1—C6—C5	177.9 (4)	C33—N5—C35—C36	-54.4 (5)
C2—C1—C6—C5	-0.1 (6)	Zr1—N5—C35—C36	126.5 (4)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N3—H3A...N1	0.84 (5)	2.56 (5)	2.983 (5)	112 (4)