

PAPER • OPEN ACCESS

Charge transfer and formation of complexes in the He⁺ collisions with the furan molecules

To cite this article: Tomasz J Wasowicz and Boguslaw Pranszke 2015 *J. Phys.: Conf. Ser.* **635** 032055

View the [article online](#) for updates and enhancements.

You may also like

- [Dissociation of furan-2-carboxylic acid by low energy electrons](#)
M Zawadzki, J Koišek and J Fedor
- [Charge dependence of fragmentation process induced by ion collisions with furan molecule](#)
E. Erdmann, M. abuda, S. Díaz-Tendero et al.
- [Fabrication of all glass microfluidic device with superior chemical and mechanical resistances by glass molding with vitreous carbon mold](#)
Muhammad Refatul Haq, Young Kyu Kim, Jun Kim et al.

PRIME
PACIFIC RIM MEETING
ON ELECTROCHEMICAL
AND SOLID STATE SCIENCE

HONOLULU, HI
Oct 6–11, 2024

Abstract submission deadline:
April 12, 2024

Learn more and submit!

Joint Meeting of
The Electrochemical Society
•
The Electrochemical Society of Japan
•
Korea Electrochemical Society

Charge transfer and formation of complexes in the He⁺ collisions with the furan molecules

Tomasz J. Wasowicz*¹ and Boguslaw Pranszke^{†,‡}

* Department of Physics of Electronic Phenomena, Gdańsk University of Technology, 80-233 Gdańsk, Poland

[†] Institute of Experimental Physics, University of Gdańsk, 80-952 Gdańsk, Poland

[‡] Present address: Gdynia Maritime University, 81-225 Gdynia, Poland

Synopsis Charge transfer and formation of the collision complexes have been studied experimentally in fragmentation of the furan molecules in collisions with He⁺ cations. The excited atomic and diatomic fragments of furan have been identified using collision-induced luminescence spectroscopy. Charge transfer ionization of the furan molecules has been observed in production of helium atoms in the excited $1s4d\ ^1D_2$, $^3D_{1,2,3}$ states. The fragmentation yields of the excited fragments have been measured as a function of the projectile energies (velocities) in the range 5-1000 eV. Enhancement of the fragmentation yields occurring at lower velocities may indicate formation of the $[\text{He-C}_4\text{H}_4\text{O}]^+$ collision complexes prior to dissociation.

Damage induced in the living cells by impact of the charged particles, for example in hadrontherapy, is related to the modifications of the structural and chemical properties of their molecular constituents. These modifications originate from the interaction of the primary cation beams and the secondary particles generated along the track such as low energy electrons, radicals and ions [1]. Unlike electrons and radicals, interaction of ions (primary and secondary) with biological media are poorly understood despite their importance in various therapies. To determine the most sensitive part of the DNA or RNA molecular chains to the cation-induced bond rupture, ionization and fragmentation of 2-deoxy-d-ribose, uracil, thymine, adenine, thymidine and amino acids under ionic irradiation have been investigated in the gas phase and layers condensed on the surfaces (see e.g. [2-4]).

Furan, C₄H₄O, has a special significance for biology and chemistry, since its ring structure resembles that of the simple sugars, ribose and deoxyribose, in the backbone of the RNA and the DNA helix.

In this communication we present studies of charge transfer and the collision complexes formation observed in fragmentation of furan molecule

in the collisions with the He⁺ cations applying collision-induced luminescence spectroscopy [5]. In Figure 1 the emission spectrum obtained for the impact of the 1000 eV He⁺ cations is presented. It reveals production of the excited hydrogen H($n=4,5$) and carbon C($2p3s\ ^1P_1$) atoms and the diatomic CH($A^2\Delta$) and C₂($d^3\Pi_g$) fragments. It also shows formation of the excited He ($1s4d\ ^1D_2$) atoms in the electron transfer from the furan molecule. Figure 2 displays the velocity dependences of the fragmentation yields, σ_{rel} , for the excited CH($A^2\Delta$) and He ($1s4d\ ^3D_{1,2,3}$) fragments. The observed rapid increase of the CH($A^2\Delta$) fragmentation yield occurring at lower velocities may be regarded as an indication of the $[\text{He-C}_4\text{H}_4\text{O}]^+$ complex being formed prior to dissociation of furan molecules [5].




Figure 1. Emission spectrum measured at an energy of 1000 eV for He⁺+C₄H₄O collisions.




Figure 2. The CH($A^2\Delta$) and He I($1s4d\ ^3D$) fragmentation yields obtained in the 15-250 km/s velocity range.

References

- [1] M.C. Fuss *et al* 2010 *Chem. Phys. Lett.* **486** 110
- [2] T. Schlathöler *et al* 2006 *Phys. Scr.* **73** C113
- [3] Z. Deng *et al* 2005 *Phys. Rev. Lett.* **95** 153201
- [4] S. Maclot *et al* 2011 *ChemPhysChem* **12** 930
- [5] T. J. Wasowicz *et al.* 2015 *J. Phys. Chem. A* **119** 581

¹ E-mail: twasowicz@mif.pg.gda.pl

