

## FIRST-ORDER DIFFERENTIAL EQUATIONS WITH NONLOCAL BOUNDARY CONDITIONS

TADEUSZ JANKOWSKI

Department of Differential Equations and Applied Mathematics, Gdansk  
University of Technology, 11/12 G.Narutowicz Str., 80-233 Gdansk, Poland

**ABSTRACT.** We study a first-order boundary value problem subject to some boundary conditions given by Riemann-Stieltjes integrals. Using a monotone iterative method, we formulate sufficient conditions which guarantee the existence of extremal or quasi-solutions in the corresponding region bounded by upper and lower solutions of our problems. The case when a unique solution exists is also investigated. Some examples are given to illustrate our results.

**Key words:** Boundary value problems for differential equations, monotone iterative technique, lower and upper solutions, existence of solutions, extremal solutions, quasi-solutions.

**AMS (MOS) Subject Classification.** 34K10, 34A40, 34A45.

### 1. INTRODUCTION

In this paper, we investigate the first-order differential equation of the form:

$$(1.1) \quad x'(t) = f(t, x(t)) \equiv Fx(t), \quad t \in J = [0, T], \quad T < \infty,$$

subject to various nonlocal Boundary Conditions (BCs):

$$(1.2) \quad x(0) = \lambda[x] + d,$$

$$(1.3) \quad x(T) = \lambda[x] + d,$$

$$(1.4) \quad x(0) = -\lambda[x] + d,$$

$$(1.5) \quad x(T) = -\lambda[x] + d,$$

where  $f \in C(J \times \mathbb{R}, \mathbb{R})$ ,  $d \in \mathbb{R}$ . Here,  $\lambda$  denotes a linear functional on  $C(J)$  given by Riemann-Stieltjes integral

$$\lambda[x] = \int_0^T x(t) dA(t)$$

with a suitable function  $A$  of bounded variation. The advantage is that the well-studied multipoint and integral BCs are both included as special cases.

Note that, BCs in (1.2)–(1.5) cover some nonlocal BCs, for example

$$\begin{aligned}\lambda[x] &= \beta x(T) + a, \\ \lambda[x] &= \sum_{i=1}^m \beta_i x(\xi_i) + b, \quad 0 < \xi_1 < \xi_2 < \cdots < \xi_m < T, \\ \lambda[x] &= \int_0^T x(t)g(t)dt + c, \quad g \in C(J, \mathbb{R}_+).\end{aligned}$$

If  $a = 0$  and  $\beta = 1$  or  $\beta = -1$ , then we have periodic or anti-periodic problem, see for example [1], [3], [11], [12], see also [7]; for multipoint problems, see [4], [6], and BC with  $g(t) = k \in \mathbb{R}$ , see [5]. Indeed, we have more papers in which boundary value problems have been discussed with BCs as in the above mentioned special cases for  $\lambda$ .

It is important to indicate that boundary conditions involving Stieltjes integrals appeared in some papers in which the problem of existence of positive solutions to differential equations have been discussed. When we apply fixed point theorems in cones, then we can obtain conditions which guarantee the existence of positive solutions in cones for boundary problems also for cases when the measure  $dA$  changes sign, see for example, [13], [14], [8], [9].

It is well known, that the monotone iterative technique offers an approach for obtaining approximate solutions to boundary value problems of differential equations, see for example [12], [2]. According to our knowledge, using the monotone iterative technique, the existence results are formulated only for special cases of functional  $\lambda$ . In this paper, we study problem (1.1) under quite general boundary conditions given by functional  $\lambda$ . To obtain the existence results we use the monotone iterative method based on inequalities and therefore we are not able to discuss our problems when the measure  $dA$  changes sign. Therefore, we discuss boundary value problems under the assumption that the measure  $dA$  in functional  $\lambda$  is non-negative. Looking on BCs (1.4), (1.5) we see that the measure  $-dA$  is now non-positive to cover the case of anti-periodic solutions too. The monotone iterative method has also been discussed in paper [10] to second order differential equations with Stieltjes integrals.

We establish sufficient conditions under which boundary value problems have solutions: extremal, quasi or a unique solution too. Two examples are added to verify theoretical results. Let us introduce the following assumption:

$H_1 : f \in C(J \times \mathbb{R}, \mathbb{R})$ ,  $A$  is a function of bounded variation and the measure  $dA$  is non-negative.

## 2. LEMMAS

First, we consider two boundary value problems:

$$(2.1) \quad \begin{cases} x'(t) = -M(t)x(t) + h(t), & t \in J, \\ x(0) = \lambda[x] + d, & d \in \mathbb{R}, \end{cases}$$

$$(2.2) \quad \begin{cases} x'(t) = M(t)x(t) - h(t), & t \in J, \\ x(T) = \lambda[x] + d, & d \in \mathbb{R}. \end{cases}$$

**Lemma 2.1.** *Suppose that:*

$H_2 : M, h \in C(J, \mathbb{R})$ , and  $A$  is a function of bounded variation and moreover

$$\int_0^T \exp\left(-\int_0^t M(s)ds\right) dA(t) \neq 1.$$

Then problem (2.1) has a unique solution given by

$$x(t) = P(t) \left\{ \left[ 1 - \int_0^T P(s) dA(s) \right]^{-1} \left[ d + \int_0^T \left( P(s) \int_0^s P^{-1}(u) h(u) du \right) dA(s) \right] \right\} \\ + P(t) \int_0^t P^{-1}(s) h(s) ds$$

with

$$P(t) = \exp\left(-\int_0^t M(\xi) d\xi\right).$$

*Proof.* Note that

$$x(t) = \exp\left(-\int_0^t M(s) ds\right) \left[ x(0) + \int_0^t \exp\left(\int_0^s M(\tau) d\tau\right) h(s) ds \right].$$

Now, using the boundary condition  $x(0) = \lambda[x] + d$  and Assumption  $H_2$ , we have the assertion of this lemma.  $\square$

Similarly as Lemma 2.1, we can prove the following result.

**Lemma 2.2.** *Suppose that:*

$H_3 : M, h \in C(J, \mathbb{R})$ , and  $A$  is a function of bounded variation and moreover

$$\int_0^T \exp\left(-\int_t^T M(s) ds\right) dA(t) \neq 1$$

Then problem (2.2) has a unique solution given by

$$x(t) = Q(t) \left\{ \left[ 1 - \int_0^T Q(s) dA(s) \right]^{-1} \left[ d + \int_0^T \left( Q(s) \int_s^T Q^{-1}(\tau) h(\tau) d\tau \right) dA(s) \right] \right\} \\ + Q(t) \int_t^T Q^{-1}(s) h(s) ds$$

with

$$Q(t) = \exp\left(-\int_t^T M(\xi) d\xi\right).$$

**Lemma 2.3.** *Suppose that:*

$H_4 : M \in C(J, \mathbb{R})$ ,  $A$  is a function of bounded variation, the measure  $dA$  is non-negative and moreover

$$(2.3) \quad \int_0^T \exp\left(-\int_0^t M(s)ds\right) dA(t) < 1.$$

Let  $p \in C^1(J, \mathbb{R})$  and

$$\begin{cases} p'(t) \leq -M(t)p(t), & t \in J, \\ p(0) \leq \lambda[p]. \end{cases}$$

Then  $p(t) \leq 0$  on  $J$ .

*Proof.* Indeed,

$$p(t) \leq \exp\left(-\int_0^t M(s)ds\right) p(0).$$

Now, using the condition  $p(0) \leq \lambda[p]$  and (2.3), we have the assertion.  $\square$

In a similar way, we can prove the following lemma.

**Lemma 2.4.** *Suppose that:*

$H_5 : M \in C(J, \mathbb{R})$ ,  $A$  is a function of bounded variation, the measure  $dA$  is non-negative and moreover

$$(2.4) \quad \int_0^T \exp\left(-\int_t^T M(s)ds\right) dA(t) < 1.$$

Let  $p \in C^1(J, \mathbb{R})$  and

$$\begin{cases} p'(t) \geq M(t)p(t), & t \in J, \\ p(T) \leq \lambda[p]. \end{cases}$$

Then  $p(t) \leq 0$  on  $J$ .

### 3. SOME COMMENTS TO SECTION 2

1. Let

$$\int_0^T x(t)dA(t) = \gamma x(\mu), \quad \gamma \geq 0.$$

Then conditions (2.3) and (2.4) take respectively the form

$$\gamma \exp\left(-\int_0^\mu M(s)ds\right) < 1, \quad \mu \in (0, T],$$

$$\gamma \exp\left(-\int_\mu^T M(s)ds\right) < 1, \quad \mu \in [0, T).$$

2. Let

$$\int_0^T x(t)dA(t) = \sum_{i=1}^m \gamma_i x(\mu_i), \quad \gamma_i > 0.$$

Then conditions (2.3) and (2.4) take respectively the form

$$\sum_{i=1}^m \gamma_i \exp \left( - \int_0^{\mu_i} M(s) ds \right) < 1, \quad 0 < \mu_1 < \mu_2 < \cdots < \mu_m \leq T,$$

$$\sum_{i=1}^m \gamma_i \exp \left( - \int_{\mu_i}^T M(s) ds \right) < 1, \quad 0 \leq \mu_1 < \mu_2 < \cdots < \mu_m < T.$$

3. Let

$$\int_0^T x(t) dA(t) = \int_0^T x(t) g(t) dt, \quad g \in C(J, \mathbb{R}_+).$$

Then conditions (2.3) and (2.4) take respectively the form

$$\int_0^T \exp \left( - \int_0^t M(s) ds \right) g(t) dt < 1,$$

$$\int_0^T \exp \left( - \int_t^T M(s) ds \right) g(t) dt < 1.$$

4. Let

$$\int_0^T x(t) dA(t) = \sum_{i=1}^m \int_{\mu_i}^{\gamma_i} x(s) g(s) ds,$$

where  $g \in C(J, \mathbb{R}_+)$ ,  $0 \leq \mu_1 < \gamma_1 < \mu_2 < \gamma_2 < \cdots < \mu_m < \gamma_m \leq T$ .

Then conditions (2.3) and (2.4) take respectively the form

$$\sum_{i=1}^m \int_{\mu_i}^{\gamma_i} \exp \left( - \int_0^t M(s) ds \right) g(t) dt < 1,$$

$$\sum_{i=1}^m \int_{\mu_i}^{\gamma_i} \exp \left( - \int_t^T M(s) ds \right) g(t) dt < 1.$$

5. Also we can consider the case when the above points **2** and **4** are combined.

#### 4. EXISTENCE OF SOLUTIONS TO PROBLEMS (1.1), (1.2) and (1.1), (1.3)

Now, we derive a fixed point result for nondecreasing mappings in ordered spaces which play a central role in our investigations. We say that  $Q : [a, b] \rightarrow [a, b]$  is nondecreasing if  $Qx \leq Qy$  for  $x, y \in [a, b]$  and  $x \leq y$ . We say that  $x \in [a, b]$  is the least fixed point of  $Q$  in  $[a, b]$  if  $x = Qx$  and if  $x \leq y$  whenever  $y \in [a, b]$  and  $y = Qy$ . The greatest fixed point of  $Q$  in  $[a, b]$  is defined similarly, by reversing the inequality. If both least and greatest fixed point of  $Q$  in  $[a, b]$  exist, we call them extremal fixed points of  $Q$  in  $[a, b]$ .

**Theorem 4.1** (see [2]). *Let  $[a, b]$  be an ordered interval in a subset  $Y$  of an ordered Banach space  $X$  and let  $Q : [a, b] \rightarrow [a, b]$  be a nondecreasing mapping. If each sequence  $\{Qx_n\} \subset Q([a, b])$  converges, whenever  $\{x_n\}$  is a monotone sequence in  $[a, b]$ , then the sequence of  $Q$ -iteration of  $a$  converges to the least fixed point  $x_*$  of*

$Q$  and the sequence of  $Q$ -iteration of  $b$  converges to the greatest fixed point  $x^*$  of  $Q$ . Moreover,

$$x_* = \min\{y \in [a, b] : y \geq Qy\}, \quad \text{and} \quad x^* = \max\{y \in [a, b] : y \leq Qy\}.$$

Let us introduce the following definition.

We say that  $u \in C^1(J, \mathbb{R})$  is a lower solution of (1.1), (1.2) if

$$u'(t) \leq Fu(t), \quad t \in J, \quad u(0) \leq \lambda[u] + d,$$

and it is an upper solution of (1.1), (1.2) if the above inequalities are reversed.

**Theorem 4.2.** *Assume that Assumption  $H_1$  holds. Let  $y_0, z_0 \in C^1(J, \mathbb{R})$  be lower and upper solutions of problem (1.1), (1.2) respectively and  $y_0(t) \leq z_0(t)$ ,  $t \in J$ . In addition, we assume that:*

$H_6$  : *there exists a function  $M \in C(J, \mathbb{R})$  such that condition (2.3) holds and*

$$f(t, u_1) - f(t, v_1) \leq M(t)[v_1 - u_1]$$

*if  $y_0(t) \leq u_1 \leq v_1 \leq z_0(t)$ .*

*Then problem (1.1), (1.2) has, in the sector  $[y_0, z_0]$ , extremal solutions, where*

$$[y_0, z_0] = \{w \in C^1(J, \mathbb{R}) : y_0(t) \leq w(t) \leq z_0(t), \quad t \in J\}.$$

*Proof.* For each  $h \in C(J, \mathbb{R})$ , problem (2.1) has a unique solution  $x$  given in Lemma 2.1. Indeed,  $x$  is also a unique fixed point of operator  $S_h$ , so  $x = S_h x$ . Choose  $h_1, h_2 \in C(J, \mathbb{R})$  such that  $h_1(t) \leq h_2(t)$  on  $J$ . Let  $x_1, x_2$  denote the solutions of problem (2.1) with  $h_1, h_2$  instead of  $h$ , respectively. Put  $p = x_1 - x_2$ . Then,

$$\begin{cases} p'(t) &= -M(t)p(t) + h_1(t) - h_2(t) \leq -M(t)p(t), \quad t \in J, \\ p(0) &= \lambda[p]. \end{cases}$$

In view of Lemma 2.3, we see that  $x_1(t) \leq x_2(t)$  on  $J$ ; so the operator  $S_h$  is nondecreasing. It is also continuous.

For  $u \in [y_0, z_0]$ , we put

$$\mathcal{F}u(t) = Fu(t) + M(t)u(t),$$

where the operator  $F$  is defined as in problem (1.1). We define the operator  $S = S_{\mathcal{F}}$ .

Let  $x_1 = Sy_0$ ,  $x_2 = Sz_0$ , so

$$\begin{cases} x_1'(t) &= -M(t)x_1(t) + \mathcal{F}y_0(t), \\ x_1(0) &= \lambda[x_1] + d, \end{cases}$$

and

$$\begin{cases} x_2'(t) &= -M(t)x_2(t) + \mathcal{F}z_0(t), \\ x_2(0) &= \lambda[x_2] + d. \end{cases}$$

Now, apply Lemma 2.3 with  $p = y_0 - x_1$ ; so it is easy to show, using the definition of the lower solution  $y_0$ , that  $y_0 \leq x_1 = Sy_0$ . Similarly, we can show  $Sz_0 = x_2 \leq z_0$ . Put  $x = x_1 - x_2$ . Then

$$\begin{cases} x'(t) &= -M(t)x(t) + Fy_0(t) - Fz_0(t) + M(t)[y_0(t) - z_0(t)] \leq -M(t)x(t), \\ x(0) &= \lambda[x]. \end{cases}$$

Using again Lemma 2.3, we see that  $x_1 \leq x_2$ ; so the operator  $S$  is nondecreasing. It means that  $y_0 \leq Su \leq z_0$  for  $u \in [y_0, z_0]$ . Hence  $S : [y_0, z_0] \rightarrow [y_0, z_0]$  and operator  $S$  is bounded because  $\|Su\| \leq \max(\|y_0\|, \|z_0\|) = B$ .

Let  $\{y_n\}$  be a monotone sequence in  $[y_0, z_0]$ ; so  $y_0 \leq Sy_n \leq z_0$ . Hence  $\|Sy_n\| \leq B$ . It is easy to show that  $\{Sy_n\}$  is equicontinuous. By Arzeli-Ascoli theorem,  $\{Ay_n\}$  is compact. It proves that  $\{Sy_n\}$  converges in  $S([y_0, z_0])$ . Finally, operator  $S$  has a least and a greatest fixed point in  $[y_0, z_0]$ , by Theorem 4.1. It results that problem (1.1), (1.2) has minimal and maximal solutions in  $[y_0, z_0]$ . This ends the proof.  $\square$

**Theorem 4.3.** *Assume that all assumptions of Theorem 4.2 hold. In addition, we assume that the following assumption  $H_7$  holds with:*

$H_7$  : *there exists a function  $L \in C(J, \mathbb{R})$  such that  $M(t) + L(t) \geq 0$ ,  $t \in J$ , condition (2.3) holds with  $L$  instead of  $-M$  and*

$$f(t, v_1) - f(t, u_1) \leq L(t)[v_1 - u_1]$$

*if  $y_0(t) \leq u_1 \leq v_1 \leq z_0(t)$ .*

*Then problem (1.1), (1.2) has, in the sector  $[y_0, z_0]$ , a unique solution.*

*Proof.* In view of Theorem 4.2,  $y_0 \leq y \leq z \leq z_0$ , where  $y, z$  are corresponding minimal and maximal solutions of problem (1.1), (1.2) in  $[y_0, z_0]$ . Put  $p = z - y$ . Hence,

$$\begin{cases} p'(t) &= Fz(t) - Fy(t) \leq L(t)p(t), \\ p(0) &= \lambda[p]. \end{cases}$$

By Lemma 2.3,  $z \leq y$ , so the assertion holds.  $\square$

Now, we will discuss problem (1.1), (1.3). We say that  $u \in C^1(J, \mathbb{R})$  is a lower solution of (1.1), (1.3) if

$$u'(t) \leq Fu(t), \quad t \in J, \quad u(T) \geq \lambda[u] + d,$$

and it is an upper solution of (1.1), (1.3) if the above inequalities are reversed.

**Theorem 4.4.** *Assume that Assumption  $H_1$  holds. Let  $y_0, z_0 \in C^1(J, \mathbb{R})$  be lower and upper solutions of (1.1), (1.3), respectively and  $z_0(t) \leq y_0(t)$ ,  $t \in J$ . In addition, we assume that the following assumption  $H_8$  holds with:*

$H_8$  : *there exists a function  $M \in C(J, \mathbb{R})$  such that condition (2.4) holds and*

$$f(t, u_1) - f(t, v_1) \geq -M(t)[v_1 - u_1]$$

if  $z_0(t) \leq u_1 \leq v_1 \leq y_0(t)$ .

Then problem (1.1), (1.3) has, in the sector  $[z_0, y_0]$ , extremal solutions.

*Proof.* For each  $h \in C(J, \mathbb{R})$ , problem (2.2) has a unique solution  $x$  given in Lemma 2.2. Indeed,  $x$  is a unique fixed point of operator  $D_h$ , so  $x = D_h x$ . Choose  $h_1, h_2 \in C(J, \mathbb{R})$  such that  $h_1(t) \leq h_2(t)$  on  $J$ . Let  $x_1, x_2$  denote the solutions of problem (2.2) with  $h_1, h_2$  instead of  $h$ , respectively. Put  $p = x_1 - x_2$ . Then,

$$\begin{cases} p'(t) = M(t)p(t) - h_1(t) + h_2(t) \geq M(t)p(t), & t \in J, \\ p(T) = \lambda[p]. \end{cases}$$

In view of Lemma 2.4, we see that  $x_1(t) \leq x_2(t)$  on  $J$ ; so the operator  $D_h$  is nondecreasing. It is also continuous.

For  $u \in [z_0, y_0]$ , we put

$$\mathcal{F}u(t) = Fu(t) - M(t)u(t),$$

where the operator  $F$  is defined as in problem (1.1). We define the operator  $D = D_{\mathcal{F}}$ . Let  $x_1 = Dz_0$ ,  $x_2 = Dy_0$ , so

$$\begin{cases} x_1'(t) = M(t)x_1(t) + \mathcal{F}z_0(t), \\ x_1(T) = \lambda[x_1] + d, \end{cases}$$

and

$$\begin{cases} x_2'(t) = M(t)x_2(t) + \mathcal{F}y_0(t), \\ x_2(T) = \lambda[x_2] + d. \end{cases}$$

Now, apply Lemma 2.4 with  $p = x_2 - y_0$ ; so it is easy to show, using the definition of the lower solution  $y_0$ , that  $y_0 \geq x_2 = Dy_0$ . Similarly, we can show  $Dz_0 = x_1 \geq z_0$ . Put  $x = x_1 - x_2$ . Then

$$\begin{cases} x'(t) = M(t)x(t) + Fz_0(t) - Fy_0(t) + M(t)[y_0(t) - z_0(t)] \geq M(t)x(t), \\ x(T) = \lambda[x]. \end{cases}$$

Using again Lemma 2.4, we see that  $x_1 \leq x_2$ ; so the operator  $D$  is nondecreasing. It means that  $z_0 \leq Du \leq y_0$  for  $u \in [z_0, y_0]$ . Hence  $D : [z_0, y_0] \rightarrow [z_0, y_0]$  and operator  $D$  is bounded because  $\|Du\| \leq \max(\|y_0\|, \|z_0\|) = B$ . Similarly as in the proof of Theorem 4.2, we can show that problem (1.1), (1.3) has minimal and maximal solutions in  $[z_0, y_0]$ . This ends the proof.  $\square$

**Theorem 4.5.** *Assume that all assumptions of Theorem 4.4 hold. In addition, we assume that the following assumption  $H_9$  holds with:*

$H_9$  : *there exists a function  $L \in C(J, \mathbb{R})$  such that  $M(t) + L(t) \geq 0$ ,  $t \in J$ , condition (2.4) holds with  $-L$  instead of  $M$  and*

$$f(t, v_1) - f(t, u_1) \geq -L(t)[v_1 - u_1]$$

if  $z_0(t) \leq u_1 \leq v_1 \leq y_0(t)$ .



Then problem (1.1), (1.3) has, in the sector  $[z_0, y_0]$ , a unique solution.

*Proof.* In view of Theorem 4.4,  $z_0 \leq z \leq y \leq y_0$ , where  $z, y$  are corresponding minimal and maximal solutions of problem (1.1), (1.3) in  $[z_0, y_0]$ . Put  $p = y - z$ . Hence,

$$\begin{cases} p'(t) = Fy(t) - Fz(t) \geq -L(t)p(t), \\ p(T) = \lambda[p]. \end{cases}$$

By Lemma 2.4,  $y \leq z$ , so the assertion holds.  $\square$

## 5. EXISTENCE OF SOLUTIONS TO PROBLEMS (1.1), (1.4) and (1.1), (1.5)

Note that BCs (1.4) and (1.5) have minus before the functional  $\lambda$ . It means that in our considerations will appear the notations of coupled lower and upper solutions. As the consequence of it, we have to discuss systems of equations or inequalities giving corresponding lemmas.

First, we consider the linear system of the form:

$$(5.1) \quad \begin{cases} y'(t) = -M(t)y(t) + h_1(t), & t \in J, & y(0) = -\lambda[z] + d, \\ z'(t) = -M(t)z(t) + h_2(t), & t \in J, & z(0) = -\lambda[y] + d. \end{cases}$$

**Lemma 5.1.** *Suppose that Assumption  $H_2$  holds with  $h_1, h_2$  instead of  $h$ .*

*Then problem (5.1) has a unique solution given by*

$$\begin{aligned} y(t) &= P(t) \left\{ \frac{1}{\Delta} \left[ d - \int_0^T P(s)H_2(s)dA(s) \right. \right. \\ &\quad \left. \left. - \left( d - \int_0^T P(s)H_1(s)dA(s) \right) \int_0^T P(s)dA(s) \right] + H_1(t) \right\}, \\ z(t) &= P(t) \left\{ \frac{1}{\Delta} \left[ d - \int_0^T P(s)H_1(s)dA(s) \right. \right. \\ &\quad \left. \left. - \left( d - \int_0^T P(s)H_2(s)dA(s) \right) \int_0^T P(s)dA(s) \right] + H_2(t) \right\} \end{aligned}$$

with

$$\begin{aligned} P(t) &= \exp \left( - \int_0^t M(\xi)d\xi \right), & \Delta &= 1 - \left( \int_0^T P(s)dA(s) \right)^2, \\ H_1(t) &= \int_0^t P^{-1}(s)h_1(s)ds, & H_2(t) &= \int_0^t P^{-1}(s)h_2(s)ds. \end{aligned}$$

*Proof.* Note that

$$\begin{aligned} y(t) &= P(t)[y(0) + H_1(t)], \\ z(t) &= P(t)[z(0) + H_2(t)]. \end{aligned}$$

Now, using the boundary conditions to eliminate  $y(0), z(0)$  and Assumption  $H_2$ , we have the assertion of this lemma.  $\square$

**Lemma 5.2.** *Suppose that Assumption  $H_3$  holds with  $h_1, h_2$  instead of  $h$ .*

*Then problem*

$$(5.2) \quad \begin{cases} y'(t) = M(t)y(t) - h_1(t), & t \in J, & y(T) = -\lambda[z] + d, \\ z'(t) = M(t)z(t) - h_2(t), & t \in J, & z(T) = -\lambda[y] + d \end{cases}$$

*has a unique solution  $(y, z)$  given as in Lemma 5.1 with*

$$\begin{aligned} P(t) &= \exp\left(-\int_t^T M(\xi)d\xi\right), & \Delta &= 1 - \left(\int_0^T P(s)dA(s)\right)^2, \\ H_1(t) &= \int_t^T P^{-1}(s)h_1(s)ds, & H_2(t) &= \int_t^T P^{-1}(s)h_2(s)ds. \end{aligned}$$

**Lemma 5.3.** *Suppose that Assumption  $H_4$  holds.*

*Let  $p, q \in C^1(J, \mathbb{R})$  and*

$$\begin{cases} p'(t) \leq M(t)p(t), & t \in J, & p(0) \leq \lambda[q], \\ q'(t) \leq M(t)q(t), & t \in J, & q(0) \leq \lambda[p]. \end{cases}$$

*Then  $p(t) \leq 0, q(t) \leq 0$  on  $J$ .*

*Proof.* Indeed,

$$\begin{aligned} p(t) &\leq \exp\left(-\int_0^t M(s)ds\right)p(0), \\ q(t) &\leq \exp\left(-\int_0^t M(s)ds\right)q(0). \end{aligned}$$

Now, using the conditions for  $p(0), q(0)$  and (2.3), we have the assertion.  $\square$

**Lemma 5.4.** *Suppose that Assumption  $H_5$  holds.*

*Let  $p, q \in C^1(J, \mathbb{R})$  and*

$$\begin{cases} p'(t) \geq M(t)p(t), & t \in J, & p(T) \leq \lambda[q], \\ q'(t) \geq M(t)q(t), & t \in J, & q(T) \leq \lambda[p]. \end{cases}$$

*Then  $p(t) \leq 0, q(t) \leq 0$  on  $J$ .*

*Proof.* Indeed,

$$\begin{aligned} p(t) &\leq \exp\left(-\int_t^T M(s)ds\right)p(T), \\ q(t) &\leq \exp\left(-\int_t^T M(s)ds\right)q(T). \end{aligned}$$

Now, using the conditions for  $p(0), q(0)$  and (2.4), we have the assertion.  $\square$

We say that  $u, v \in C^1(J, \mathbb{R})$  are coupled lower and upper solutions of problem (1.1), (1.4) if

$$\begin{cases} u'(t) \leq Fu(t), & t \in J, & u(0) \leq -\lambda[v] + d, \\ v'(t) \geq Fv(t), & t \in J, & v(0) \geq -\lambda[u] + d. \end{cases}$$

We say that  $U, V \in C^1(J, \mathbb{R})$  are coupled quasi-solutions of problem (1.1), (1.4) if

$$\begin{cases} U'(t) = FU(t), & t \in J, & U(0) = -\lambda[V] + d, \\ V'(t) = FV(t), & t \in J, & V(0) = -\lambda[U] + d. \end{cases}$$

**Theorem 5.5.** *Assume that Assumptions  $H_1, H_6$  hold. Let  $y_0, z_0 \in C^1(J, \mathbb{R})$  be coupled lower and upper solutions of (1.1), (1.4) and  $y_0(t) \leq z_0(t)$ ,  $t \in J$ .*

*Then problem (1.1), (1.4) has, in the sector  $[y_0, z_0]$ , coupled quasi-solutions  $y, z$  and  $y \leq z$ .*

*Proof.* Let  $\eta, \xi \in [y_0, z_0]$ . Put  $\varphi(t) = \sup[\eta(t), \xi(t)]$ ,  $\Phi(t) = \inf[\eta(t), \xi(t)]$ . Consider the following system:

$$(5.3) \quad \begin{cases} v'(t) = F\Phi(t) - M(t)[v(t) - \Phi(t)], & t \in J, & v(0) = -\lambda[w] + d, \\ w'(t) = F\varphi(t) - M(t)[w(t) - \varphi(t)], & t \in J, & w(0) = -\lambda[v] + d. \end{cases}$$

By Lemma 5.1, system (5.3) has a unique solution. Therefore, we can define the operator

$$(5.4) \quad B : \bar{\Omega} \rightarrow C^1(J) \times C^1(J), \quad B(\eta, \xi) = (v, w),$$

where  $(v, w)$  is the solution of (5.3),  $\bar{\Omega} = [y_0, z_0] \times [y_0, z_0]$ .

Now, we want to show that

$$(5.5) \quad y_0(t) \leq v(t) \leq w(t) \leq z_0(t), \quad t \in J.$$

Put  $p = y_0 - v$ ,  $q = w - z_0$ . Then

$$\begin{aligned} p'(t) &\leq Fy_0(t) - F\Phi(t) + M(t)[v(t) - \Phi(t)] \leq M(t)[\Phi(t) - y_0(t)] = -M(t)p(t), \\ q'(t) &\leq F\varphi(t) - M(t)[w(t) - \varphi(t)] - Fz_0(t) \leq -M(t)q(t). \end{aligned}$$

Moreover

$$\begin{aligned} p(0) &\leq -\lambda[z_0] + \lambda[w] = \lambda[q], \\ q(0) &\leq -\lambda[v] + \lambda[y_0] = \lambda[p]. \end{aligned}$$

This and Lemma 5.3 show that  $y_0(t) \leq v(t)$ ,  $w(t) \leq z_0(t)$ ,  $t \in J$ . To show that  $v(t) \leq w(t)$ ,  $t \in J$ , we put  $p = v - w$ . Then

$$\begin{aligned} p'(t) &= F\Phi(t) - F\varphi(t) - M(t)[v(t) - \Phi(t) - w(t) + \varphi(t)] \leq -M(t)p(t), \\ p(0) &= -\lambda[w] + \lambda[v] = \lambda[p]. \end{aligned}$$

This and Lemma 2.3 show that  $v(t) \leq w(t)$ ,  $t \in J$  so (5.5) holds.

Hence  $B : \bar{\Omega} \rightarrow \bar{\Omega}$ . Using (5.3), we can define two sequences  $\{y_n, z_n\}$  by relations

$$\begin{cases} y'_{n+1}(t) = Fy_n(t) - M(t)[y_{n+1}(t) - y_n(t)], & t \in J, & y_{n+1}(0) = -\lambda[z_{n+1}] + d, \\ z'_{n+1}(t) = Fz_n(t) - M(t)[z_{n+1}(t) - z_n(t)], & t \in J, & z_{n+1}(0) = -\lambda[y_{n+1}] + d \end{cases}$$

for  $n = 0, 1, \dots$ . In view of (5.5), we have

$$y_0(t) \leq y_1(t) \leq \dots \leq y_n(t) \leq y_{n+1}(t) \leq z_{n+1}(t) \leq z_n(t) \leq \dots \leq z_1(t) \leq z_0(t).$$

The sequence  $\{y_n\}$  is nondecreasing while  $\{z_n\}$  is nonincreasing. Note that the sequences  $\{y_n, z_n\}$  are uniformly bounded. Indeed,  $y_n, z_n$  are equicontinuous too.

The Arzela-Ascoli theorem guarantees the existence of subsequences  $\{y_{n_k}, z_{n_k}\}$  of  $\{y_n, z_n\}$ , respectively, and continuous functions  $y, z$  with  $y_{n_k}, z_{n_k}$  converging uniformly on  $J$  to  $y$  and  $z$ , respectively. Note that  $y_{n_k}, z_{n_k}$  satisfy the integral equations

$$\begin{cases} y_{n_k+1}(t) = y_{n_k+1}(0) + \int_0^t [Fy_{n_k}(s) - M(t)(y_{n_k+1}(s) - y_{n_k}(s))] ds, & t \in J, \\ z_{n_k+1}(t) = z_{n_k+1}(0) + \int_0^t [Fz_{n_k}(s) - M(t)(z_{n_k+1}(s) - z_{n_k}(s))] ds, & t \in J, \end{cases}$$

and

$$\begin{cases} y_{n_k+1}(0) = -\lambda[z_{n_k+1}] + d, \\ z_{n_k+1}(0) = -\lambda[y_{n_k+1}] + d. \end{cases}$$

If  $n_k \rightarrow \infty$ , then from the above relations, we have

$$\begin{cases} y(t) = y(0) + \int_0^t Fy(s)ds, & t \in J, & y(0) = -\lambda[z] + d, \\ z(t) = z(0) + \int_0^t Fz(s)ds, & t \in J, & z(0) = -\lambda[y] + d \end{cases}$$

because  $f$  is continuous. Thus  $y, z \in C^1(J)$  and

$$y'(t) = Fy(t), \quad z'(t) = Fz(t), \quad t \in J.$$

It proves that  $y, z$  are coupled quasi-solutions of problem (1.1), (1.4) and  $y \leq z$ . This ends the proof.  $\square$

Our next theorem concerns the case when problem (1.1), (1.4) has a unique solution.

**Theorem 5.6.** *Assume that all assumptions of Theorem 5.5 are satisfied. In addition assume that Assumption  $H_7$  holds.*

*Then problem (1.1), (1.4) has, in the sector  $[y_0, z_0]$ , a unique solution.*

*Proof.* Theorem 5.5 guarantees that functions  $y, z$  are coupled quasi-solutions of problem (1.1), (1.4) and  $y_0(t) \leq y(t) \leq z(t) \leq z_0(t)$ ,  $t \in J$ . We first show that  $y(t) = z(t)$ ,  $t \in J$ . Put  $p = z - y$ . Then

$$\begin{aligned} p'(t) &= Fz(t) - Fy(t) \leq L(t)p(t), & t \in J, \\ p(0) &= \lambda[p]. \end{aligned}$$

In view of Lemma 2.3,  $y(t) \geq z(t)$ ,  $t \in J$ . It proves that  $y = z$ , so problem (1.1), (1.4) has a solution.

It remains to show that  $y = z$  is a unique solution of (1.1), (1.4) in the sector  $[y_0, z_0]$ . Let  $w \in [y_0, z_0]$  be any solution of (1.1), (1.4). We assume that  $y_m(t) \leq$

$w(t) \leq z_m(t)$ ,  $t \in J$  for some  $m$ . Let  $p = y_{m+1} - w$ ,  $q = w - z_{m+1}$ , where  $y_m, z_m$  are defined as in Theorem 5.5. Then,

$$\begin{aligned} p'(t) &= Fy_m(t) - M(t)[y_{m+1}(t) - y_m(t)] - Fw(t) \leq -M(t)p(t), \\ q'(t) &= Fw(t) - Fz_m(t) + M(t)[z_{m+1}(t) - z_m(t)] \leq -M(t)q(t), \end{aligned}$$

and

$$p(0) = \lambda[q], \quad q(0) = \lambda[p].$$

It gives  $y_{m+1}(t) \leq w(t) \leq z_{m+1}(t)$ ,  $t \in J$ . By induction,  $y_n(t) \leq w(t) \leq z_n(t)$ ,  $t \in J$ ,  $n = 0, 1, \dots$ . If  $n \rightarrow \infty$ , then  $y = z = w$  which proves the assertion of our theorem.  $\square$

Now, we will discuss problem (1.1), (1.5). We say that  $u, v \in C^1(J, \mathbb{R})$  are coupled lower and upper solutions of problem (1.1), (1.5) if

$$\begin{cases} u'(t) \leq Fu(t), & t \in J, & u(T) \geq -\lambda[v] + d, \\ v'(t) \geq Fv(t), & t \in J, & v(T) \leq -\lambda[u] + d. \end{cases}$$

We say that  $U, V \in C^1(J, \mathbb{R})$  are coupled quasi-solutions of problem (1.1), (1.4) if

$$\begin{cases} U'(t) = FU(t), & t \in J, & U(T) = -\lambda[V] + d, \\ V'(t) = FV(t), & t \in J, & V(T) = -\lambda[U] + d. \end{cases}$$

**Theorem 5.7.** *Assume that Assumptions  $H_1, H_8$  hold. Let  $y_0, z_0 \in C^1(J, \mathbb{R})$  be coupled lower and upper solutions of (1.1), (1.5) and  $z_0(t) \leq y_0(t)$ ,  $t \in J$ .*

*Then problem (1.1), (1.5) has, in the sector  $[z_0, y_0]$ , coupled quasi-solutions  $y, z$  and  $z \leq y$ .*

*Proof.* We introduce only the definitions of sequences:

$$\begin{aligned} y'_{n+1}(t) &= Fy_n(t) + M(t)[y_{n+1}(t) - y_n(t)], & y_{n+1}(T) &= -\lambda[z_{n+1}] + d, \\ z'_{n+1}(t) &= Fz_n(t) + M(t)[z_{n+1}(t) - z_n(t)], & z_{n+1}(T) &= -\lambda[y_{n+1}] + d. \end{aligned}$$

Similarly as before, we can prove the assertion of this theorem.  $\square$

Our next theorem concerns the case when problem (1.1), (1.5) has a unique solution.

**Theorem 5.8.** *Assume that all assumptions of Theorem 5.7 are satisfied. In addition assume that Assumption  $H_7$  holds.*

*Then problem (1.1), (1.5) has, in the sector  $[z_0, y_0]$ , a unique solution.*



## 6. EXAMPLES

**Example 6.1.** Consider the following differential equation:

$$(6.1) \quad x'(t) = 2[\exp(x(t)) - \exp(-1)] \equiv Fx(t), \quad t \in J = [0, T].$$

Note that  $M(t) = 2\exp(t)$ , see the condition for  $f$  in Assumption  $H_8$ . Now we consider equation (6.1) with the boundary condition defined in points **1.** or **2.**

**1.** Let  $\lambda[x] = \int_0^T x(s)g(s)ds$ ,  $g \in C(J, \mathbb{R}_+)$ , so

$$(6.2) \quad x(T) = \int_0^T x(s)g(s)ds, \quad g \in C(J, \mathbb{R}_+).$$

Let us assume that

$$(6.3) \quad \int_0^T \exp[2\exp(s)]g(s)ds < \exp[2\exp(T)],$$

$$(6.4) \quad \int_0^T g(s)ds \leq 1, \quad \int_0^T sg(s)ds \leq T.$$

Take  $y_0(t) = t$ ,  $z_0(t) = -1$ ,  $t \in J$ . Then

$$\begin{aligned} Fy_0(t) &= 2[\exp(t) - \exp(-1)] > 1 = y_0'(t), \\ Fz_0(t) &= 0 = z_0'(t) \end{aligned}$$

and

$$\begin{aligned} \int_0^T y_0(s)g(s)ds &= \int_0^T sg(s)ds \leq T = y_0(T), \\ \int_0^T z_0(s)g(s)ds &= -\int_0^T g(s)ds \geq -1 = z_0(T), \end{aligned}$$

by (6.4). It proves that  $y_0, z_0$  are lower and upper solutions of problem (6.1), (6.2). By Theorem 4.4, this problem has extremal solutions in the sector  $[-1, t]$ .

For example, if we take  $g(t) = t$ ,  $T < \sqrt{2}$ , then conditions (6.3) and (6.4) are satisfied.

**2.** Let  $\lambda[x] = -\int_0^T x(s)g(s)ds - 1$ ,  $g \in C(J, \mathbb{R}_+)$ , so

$$(6.5) \quad x(T) = -\int_0^T x(s)g(s)ds - 1, \quad g \in C(J, \mathbb{R}_+).$$

Let us assume that conditions (6.3) and (6.6) hold with

$$(6.6) \quad \int_0^T g(s)ds \leq 1.$$

Put  $y_0(t) = 0$ ,  $z_0(t) = -1$ ,  $t \in J$ . Then

$$Fy_0(t) > 0 = y_0'(t), \quad Fz_0(t) = 0 = z_0'(t)$$



and

$$\begin{aligned} - \int_0^T z_0(s)g(s)ds - 1 &= \int_0^T g(s)ds - 1 \leq 0 = y_0(T), \\ - \int_0^T y_0(s)g(s)ds - 1 &= -1 = z_0(T), \end{aligned}$$

by (6.6). It proves that  $y_0, z_0$  are coupled lower and upper solutions of problem (6.1), (6.5). By Theorem 5.7, this problem has quasi-solutions in the sector  $[-1, 0]$ .

**Example 6.2** (see [5]). Consider the following differential equation:

$$(6.7) \quad x'(t) = \exp(t \sin^2 x(t)) \equiv Fx(t), \quad t \in J = [0, T] \text{ with } T = \ln 2.$$

We consider equation (6.7) with the condition defined in points **1** or **2** (the results are taken from paper [5]),

**1.** Let  $\lambda[x] = \int_0^T x(s)ds$ , so

$$(6.8) \quad x(0) = \int_0^T x(s)ds.$$

Note that  $y_0(t) = 0$ ,  $z_0(t) = \exp(t)$ ,  $t \in J$  are lower and upper solutions of (6.7), respectively. Problem (6.7), (6.8) has extremal solutions in the sector  $[y_0, z_0]$ , by Theorem 4.2.

**2.** Let  $\lambda[x] = - \int_0^T x(s)ds$ , so

$$(6.9) \quad x(0) = - \int_0^T x(s)ds.$$

Indeed,  $y_0(t) = -1$ ,  $z_0(t) = \exp(t)$ ,  $t \in J$  are coupled lower and upper solutions of (6.7). Problem (6.7), (6.9) has quasi-solutions in the sector  $[y_0, z_0]$ , by Theorem 5.5.

## REFERENCES

- [1] D. Franco, J. Nieto and D. O'Regan, Existence of solutions for first order differential equations with nonlinear boundary conditions, *Appl. Math. Comput.*, 153:793–802, 2004.
- [2] S. Heikkilä and V. Lakshmikantham, *Monotone Iterative Technique for Discontinuous Nonlinear Differential Equations*, Dekker, New York, 1994.
- [3] T. Jankowski, BVPs for functional differential equations, *Dynam. Systems Appl.*, 9:445–455, 2000.
- [4] T. Jankowski, Multipoint boundary value problems for ODEs. I, *Appl. Anal.*, 80:395–407, 2001.
- [5] T. Jankowski, Differential equations with integral boundary conditions, *J. Comput. Appl. Math.*, 147:1–8, 2002.
- [6] T. Jankowski, Multipoint boundary value problems for ODEs. II, *Czechoslovak Math. J.*, 54:843–854, 2004.
- [7] T. Jankowski, Existence of solutions of boundary value problems for differential equations in which deviated arguments depend on the unknown solution, *Comput. Math. Appl.*, 54:357–363, 2007.

- [8] T. Jankowski, Positive solutions for second order impulsive differential equations involving Stieltjes integral conditions, *Nonlinear Anal.*, 74:3775–3785, 2011.
- [9] T. Jankowski, Existence of positive solutions to third order differential equations with advanced arguments and nonlocal boundary conditions, *Nonlinear Anal.*, 75:913–923, 2012.
- [10] T. Jankowski and R. Jankowski, Monotone iterative method to second order differential equations with deviating arguments involving Stieltjes integral boundary conditions, *Dynam. Systems Appl.*, 21:17–32, 2012.
- [11] D. Jiang and J. Wei, Monotone method for first- and second-order periodic boundary value problems and periodic solutions of functional differential equations, *Nonlinear Anal.*, 50:885–898, 2002.
- [12] G. S. Ladde, V. Lakshmikantham and A. S. Vatsala, *Monotone Iterative Techniques for Nonlinear Differential Equations*, Pitman, Boston, 1985.
- [13] J. R. L. Webb and G. Infante, Positive solutions of nonlocal boundary value problems: a unified approach, *J. London Math. Soc.*, 74:673–693, 2006.
- [14] J. R. L. Webb and G. Infante, Positive solutions of nonlocal boundary value problems involving integral conditions, *NoDEA Nonlinear Differential Equations Appl.*, 15:45–67, 2008.