



Neutral pH, multioxidants Fenton oxidation of dimethyl sulfoxide and acetamidophenol as water pollutant models

Aswin Kottapurath Vijay^{1,2} · Gifty Sara Rolly^{3,4} · Vered Marks⁵ · Virender K. Sharma⁶ · Dan Meyerstein^{1,3}

Received: 18 October 2024 / Accepted: 15 February 2025
© The Author(s) 2025

Abstract

Organic water pollution is calling for advanced remediation methods such as the Fenton process, yet actual procedures involve transition metals at acidic pH, and generate only one oxidant, the hydroxyl radical. Here we used a solution of magnesium ions, bicarbonate ions, and hydrogen peroxide at pH 7.4 to generate reactive oxygen species for degrading dimethyl sulfoxide and acetamidophenol, as models of water pollutants. The performance and the identification of degradation products were assessed by nuclear magnetic resonance and high-performance liquid chromatography. Results show the generation of several oxidizing agents such as hydroxyl radical, carbonate anion radical, and superoxide. The novel aspect is that the Fenton-like process can be achieved with Mg^{2+} serving only as a template to facilitate redox reactions rather than participating directly. The mechanisms for generating oxidizing radicals suggests potential applications in both environmental cleanup and biological processes.

Keywords Advanced oxidation processes · Biology · Carbonate-anion-radical · Environmental Fenton-like-reactions · Magnesium

Introduction

The growing accumulation of waste in freshwater systems is increasingly threatening both planetary health and societal sustainability [1]. Currently, there is a sharp increase in waste and pollution due to shifts in human activities following the pandemic. There are growing concerns about the disposal of unused pharmaceuticals and their metabolites, which are often inadequately treated by wastewater treatment plants and could potentially lead to a global health crisis [2]. Researchers are continually developing newer and greener strategies to address the challenge of cleaning water contaminated with pharmaceuticals [3, 4]. The major focus has been on innovating existing technologies to address the current challenges in water purification.

The Fenton and Fenton-like reactions, e.g., $Fe^{2+} + H_2O_2 \rightarrow Fe^{3+} + OH^- + OH^\cdot$, have been investigated for several decades due to their potential to generate highly reactive hydroxyl radical (OH^\cdot) in biological systems [5–7], in environmental processes [8, 9] and in advanced oxidation processes/technologies, AOPs [10–12]. Fenton and Fenton-like reactions are primarily conducted using Fe^{2+} and occasionally involve other transition metal complexes. Transition-metal-based Fenton reactions, often involve the formation

✉ Dan Meyerstein
danm@ariel.ac.il

Virender K. Sharma
vks38@miami.edu

¹ Chemistry Department, Ben-Gurion University, Beer-Sheva, Israel

² Department of Sanitary Engineering, Faculty of Civil and Environmental Engineering, Gdańsk University of Technology, Gdańsk, Poland

³ Department of Chemical Sciences and The Radical Research Center, Ariel University, Ariel, Israel


⁴ Department of Energy Conversion and Storage, Faculty of Chemistry, Gdańsk University of Technology, 80-233 Gdańsk, Poland

⁵ Department of Chemical Sciences, Ariel University, Ariel, Israel


⁶ Department of Chemical, Environmental and Material Engineering, University of Miami, 1251 Memorial Drive, Coral Gables, FL 33146, USA

of metal complexes and the precipitation of metal oxides/hydroxides, e.g., $\text{Fe}(\text{OH})_3$ [13], which poses treatment efficiency, water quality, and sludge generation challenges.

Researchers have been exploring replacing transition metal complexes with other metals (M) like beryllium and aluminum to carry out Fenton-like processes to produce OH^\bullet through reaction (1) [14–17]. However, reaction (1) typically occurs only in organic solvents with minimal water and requires very high concentrations of hydrogen peroxide (H_2O_2) to proceed. Additionally, an acidic environment must be maintained for the reaction. These stringent conditions make reaction (1) impractical for treating polluted water directly.



We stride out to achieve reaction (1) possible for treating water under neutral conditions. This stems from consideration of the similarity in the redox potentials of the couples ($(\text{OOH}^\bullet + \text{e}^- + \text{H}^+)/(\text{H}_2\text{O}_2) = 1.46 \text{ V}$ versus NHE [18] and $(\text{CO}_3^{\bullet-} + \text{e}^-)/\text{CO}_3^{2-} = 1.57 \text{ V}$ versus NHE [18], and thus it seemed reasonable to expect the feasibility of the occurrence of reaction (2).



Significantly, herein we provide evidence of the occurrence of a Fenton-like reaction in an aqueous environment by using magnesium ion (Mg^{2+}), i.e., without the use of a transition metal complex. Magnesium ion (Mg^{2+}) was intuitively chosen as the cation due to the following reasons: (i) it is a small cation and therefore expected to complex with H_2O_2 better than Fe^{2+} , (ii) magnesium salts are soluble in neutral solutions even in the presence of bicarbonate, (iii) magnesium ion plays several important roles in the environment due to its abundance and unique chemical properties [19, 20] and (iv) magnesium ions are present in most of the water bodies including tap water and seawater [21] and thus address the drawbacks of secondary contamination of conventional Fenton reaction processes to treat polluted water.

Importantly, reaction (2) directly forms two strong oxidizing radicals: OH^\bullet and carbonate anion radical ($\text{CO}_3^{\bullet-}$), resulting in a multi-oxidant system. The $\text{CO}_3^{\bullet-}$ as an oxidant is of tremendous interest due to its involvement in various catalytic oxidation processes [22, 23], some of which are major biological [24, 25] and water remediation importance [22, 26]. Significantly, $\text{CO}_3^{\bullet-}$ anion radicals, despite their lower reactivity compared to OH^\bullet radicals, exhibit a specific reactivity towards active moieties of pollutants, establishing their significant role in AOPs [27].

This study thus aims to provide a greener solution for depolluting water. The solutions containing $\text{Mg}(\text{H}_2\text{O})_6^{2+}$, HCO_3^- , and H_2O_2 could carry the Fenton-like processes effectively. Experiments focused on the oxidation of dimethyl sulfoxide (DMSO) and acetamidophenol were performed to demonstrate a new Fenton-like greener process to de-contaminate water. These chosen model reactions serve as representative systems to investigate the intricate mechanisms and dynamics involved in Fenton-like processes in the studied solutions. The implications of these findings in biological processes and the potential for alternative advanced oxidation processes without transition-metal complexes are briefly discussed.

Results and discussion

The experimental procedures are detailed in the supplemental material (Text S1). In an initial experiment, the nature of the formed intermediate oxidizing agents in the reaction mixtures of Mg^{2+} and H_2O_2 in the absence and presence of HCO_3^- was studied. Dimethylsulfoxide ($(\text{CH}_3)_2\text{SO}$) was

introduced into the reaction mixtures, and the identification of dimethylsulfone (DMSO_2) as a product was established through the analysis of ^1H NMR spectra. The results are presented in Fig. 1. The formation of DMSO_2 was not seen in any solution mixtures except in which both Mg^{2+} and HCO_3^- were present together in the solution.

The results in Fig. 1 indicate the oxidation of DMSO by OH^\bullet and $\text{CO}_3^{\bullet-}$ and by $\text{HO}_2^\bullet/\text{O}_2^{\bullet-}$. These reactions have been reported to yield CH_3^\bullet radicals [28]. In this system, the production of radicals is very slow, and therefore $\text{CO}_3^{\bullet-}$ and $\text{O}_2^{\bullet-}$ are long-lived and enable their reactions with DMSO [28] and with H_2O_2 , $k(\text{CO}_3^{\bullet-} + \text{H}_2\text{O}_2) \sim 4.0 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$ [29]. The CH_3^\bullet radicals formed react with H_2O_2 , $k(\text{CH}_3^\bullet + \text{H}_2\text{O}_2 \rightarrow \text{CH}_4 + \text{HO}_2^\bullet/(\text{O}_2^{\bullet-} + \text{H}^+)) = 2.70 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$ [30]. Thus all the radicals formed in reaction (2) are transformed into $\text{HO}_2^\bullet/\text{O}_2^{\bullet-}$ radicals that are expected to react with DMSO to yield CH_3^\bullet and $\text{CH}_3\text{S}(\text{O})\text{OOH}$ [28]. This explains the observed oxidation products in Fig. 1.

The oxidizing properties of a solution containing a mixture of Mg^{2+} , HCO_3^- , and H_2O_2 were further tested to oxidize a well-known micropollutant, acetamidophenol (ACP). The results are presented in Fig. 2, which clearly shows the efficient oxidation of ACP when all

Fig. 1 ^1H nuclear magnetic resonance (NMR) spectra of the products of the reactions of $\text{Mg}^{2+} + \text{H}_2\text{O}_2$ with dimethylsulfoxide (DMSO) at pH 7.40 with and without HCO_3^- . Measured one week after preparation of samples, de-aerated solutions. Experimental conditions: $[\text{H}_2\text{O}_2] = 5.0 \text{ mM}$, $[\text{HCO}_3^-] = 5.0 \text{ mM}$, $[\text{Mg}^{2+}] = 1.0 \text{ mM}$, and $[\text{DMSO}] = 25.0 \text{ mM}$. The results show that DMSO is oxidized by both OH^\bullet and $\text{CO}_3^{\bullet-}$ radicals, as well as by $\text{HO}_2^\bullet/\text{O}_2^{\bullet-}$ species. These reactive species contribute to the oxidation processes observed





Fig. 2 Removal efficiency of acetamidophenol (ACP) was obtained for the systems: $\text{HCO}_3^- + \text{ACP}$, $\text{Mg}^{2+} + \text{ACP}$, $\text{H}_2\text{O}_2 + \text{ACP}$, $\text{H}_2\text{O}_2 + \text{HCO}_3^- + \text{ACP}$, $\text{Mg}^{2+} + \text{H}_2\text{O}_2 + \text{ACP}$, and $\text{Mg}^{2+} + \text{H}_2\text{O}_2 + \text{HCO}_3^- + \text{ACP}$. ($[\text{HCO}_3^-]$, $[\text{H}_2\text{O}_2]$, and $[\text{MgCl}_2 \cdot 6\text{H}_2\text{O}] = 10.0 \text{ mM}$, $[\text{ACP} = \text{acetamidophenol}] = 0.25 \text{ mM}$). All samples were at pH 7.40, de-aerated, and analyzed by HPLC after one week of preparation. The results indicate that the removal efficiency of ACP was evaluated in various systems, including HCO_3^- , Mg^{2+} , H_2O_2 , and their combinations with ACP. HPLC analysis revealed varying levels of ACP removal for each system, with the most complex system ($\text{Mg}^{2+} + \text{H}_2\text{O}_2 + \text{HCO}_3^- + \text{ACP}$) showing significant removal efficiency

components of the mixture are present. Experiments using four mixtures, $\text{HCO}_3^- + \text{ACP}$, $\text{Mg}^{2+} + \text{ACP}$, $\text{H}_2\text{O}_2 + \text{ACP}$, and $\text{Mg}^{2+} + \text{H}_2\text{O}_2 + \text{ACP}$ have no oxidation capacity to degrade ACP. The results are adequately explained by the lack of formation of radicals in these reaction media. The observation that the mixture of Mg^{2+} and H_2O_2 does not oxidize the ACP points out that the Shul'pin mechanism [14–17] is not operating in aqueous solutions and at low








Fig. 3 High performance liquid chromatography-ultraviolet detection (HPLC-UV) chromatograms were obtained for the acetamidophenol (ACP) degradation in the systems: ACP (green), $\text{H}_2\text{O}_2 + \text{HCO}_3^- + \text{ACP}$ (purple), and $\text{Mg}^{2+} + \text{H}_2\text{O}_2 + \text{HCO}_3^- + \text{ACP}$ (orange). The lines are shifted horizontally for better vision. The figure indicates the HPLC chromatogram of the major oxidized product (OP) of ACP. ACP eluted at a retention time of 5.50 min, while the oxidized product was observed at 6.70 min, indicating successful oxidation

H_2O_2 concentrations. Whereas $\text{H}_2\text{O}_2 + \text{HCO}_3^- + \text{ACP}$ and $\text{Mg}^{2+} + \text{H}_2\text{O}_2 + \text{HCO}_3^- + \text{ACP}$ systems showed 25 and 60 percent acetamidophenol degradation respectively. The chromatogram of the major oxidized product (OP) analyzed by HPLC is shown in Fig. 3. The acetamidophenol was eluted at a retention time of 5.50 min, while the oxidized product at 6.70. The pH values mentioned throughout the

manuscript refer to the final pH measured after the completion of the reaction.

The results point out that oxidizing reactive species are formed in a neutral aqueous solution containing Mg^{2+} , H_2O_2 , and HCO_3^- . This suggests that OH^\bullet and $\text{CO}_3^{\bullet-}$ anion radicals are indeed formed via reaction (2) or $\text{CO}_3^{\bullet-}$ and $\text{HO}_2^\bullet/\text{O}_2^{\bullet-}$ via reactions (3)–(6) that are analogous to the proposed mechanism for an analogous system containing $\text{Co}^{\text{II}}_{\text{aq}}$ [31].



The results demonstrate that indeed $\text{CO}_3^{\bullet-}$ are formed via one of these mechanisms. It should be noted that the observed process is considerably slower than the Fenton reaction.

Conclusion

The results clearly show that the solutions of Mg^{2+} , H_2O_2 , and HCO_3^- generate reactive oxygen species, OH^\bullet and $\text{CO}_3^{\bullet-}$, or $\text{CO}_3^{\bullet-}$ and $\text{HO}_2^\bullet/\text{O}_2^{\bullet-}$ under neutral pH conditions. In these systems, the central cation acts only as a template for a redox process between its ligands. Hence, the presented Fenton-like reaction system without transition metal ions opens the door to new advantageous AOPs. The results point out that carbonate anion radicals are formed also, via the mechanism reported herein, in oceans and other natural waters that contain high concentrations of Mg^{2+} and HCO_3^- (Mg^{2+} is present in high concentrations in oceans). Furthermore, the presence of Mg^{2+} in biological systems, where it typically acts as an antioxidant due to complex biological mechanisms, raises intriguing questions. The possibility that the Fenton-like processes reported herein may have been overlooked in biological systems is a noteworthy point. This opens avenues for further research into the role of Mg^{2+} in biological processes and whether it contributes to radical-induced degradation in living organisms. MgCO_3 is a common additive in pharmacological preparations and foods. Thus it has to be investigated whether the new Fenton-like process reported herein plays a role in the radical-induced degradation of pharmaceuticals and foods.

Further studies are crucial to gain a better understanding of the phenomena and their implications. Overall, the findings lead to the development of new advanced oxidation processes that do not require transition metal complexes as catalysts. The advantage of bicarbonate as a co-catalyst in the process is that it does not decompose in the catalytic oxidation processes.

Acknowledgments This study was enabled by grants from the Pazy Foundation and Ariel University.

Funding Open access funding provided by Ariel University.

Declarations

Conflict of interest The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Kuang X, Liu J, Scanlon BR, Jiao JJ, Jasechko S, Lancia M, Biskaborn BK, Wada Y, Li H, Zeng Z, Guo Z, Yao Y, Gleeson T, Nicot J-P, Luo X, Zou Y, Zheng C (2024) The changing nature of groundwater in the global water cycle. *Science*. <https://doi.org/10.1126/science.adf0630>
- Han J, He S, Lichtfouse E (2023) Waves of pharmaceutical waste. *Environ Chem Lett* 21:1251–1255. <https://doi.org/10.1007/s10311-022-01491-0>
- Sharma VK, Ma X, Zboril R (2023) Single atom catalyst-mediated generation of reactive species in water treatment. *Chem Soc Rev* 52:7673–7686. <https://doi.org/10.1039/D3CS00627A>
- Parvulescu VI, Epron F, Garcia H, Granger P (2022) Recent progress and prospects in catalytic water treatment. *Chem Rev* 122:2981–3121. <https://doi.org/10.1021/acs.chemrev.1c00527>
- Nidheesh PV, Zhou M, Oturan MA (2018) An overview on the removal of synthetic dyes from water by electrochemical advanced oxidation processes. *Chemosphere* 197:210–227. <https://doi.org/10.1016/j.chemosphere.2017.12.195>
- Wang X, Liu H, Xue Y, Cui L, Chen L, Ho K, Huang Y (2024) Formation of environmentally persistent free radicals and their risks for human health: a review. *Environ Chem Lett* 22:1327–1343. <https://doi.org/10.1007/s10311-024-01701-x>
- Wang F, Zhang Y, Ming H, Wang L, Zhao Z, Wang Y, Liang J, Qin Y (2020) Degradation of the ciprofloxacin antibiotic by photo-Fenton reaction using a Nafion/iron membrane: role of hydroxyl radicals. *Environ Chem Lett* 18:1745–1752. <https://doi.org/10.1007/s10311-020-01018-5>

8. Wang L, Rao L, Ran M, Shentu Q, Wu Z, Song W, Zhang Z, Li H, Yao Y, Lv W, Xing M (2023) A polymer tethering strategy to achieve high metal loading on catalysts for Fenton reactions. *Nat Commun* 14:7841. <https://doi.org/10.1038/s41467-023-43678-1>
9. Sacchetto JL, Medina LF, Toledo KI, Plem SC, Jalit Y, Gatica EA, Miskoski S, Natera J, Lépori CMO, Massad WA (2024) Epoxiconazole degradation in water samples: a comparative study of Fenton, photo-Fenton, solar photo-Fenton, and solar photolysis processes. *Photochem Photobiol Sci* 23:1143–1153. <https://doi.org/10.1007/s43630-024-00582-x>
10. Farinelli G, Gil AG, Marugan J, Minella M, Fabbri D, Laurenti E, Tiraferri A, Vione D (2024) The dominant role of the peroxymonosulfate radical for removing contaminants in a Fenton process with metabisulfite. *Environ Chem Lett* 22:43–48. <https://doi.org/10.1007/s10311-023-01645-8>
11. Sözen S, Olmez-Hanci T, Hooshmand M, Orhon D (2020) Fenton oxidation for effective removal of color and organic matter from denim cotton wastewater without biological treatment. *Environ Chem Lett* 18:207–213. <https://doi.org/10.1007/s10311-019-00923-8>
12. Morin-Crini N, Lichtfouse E, Fourmentin M, Ribeiro ARL, Noutsopoulos C, Mapelli F, Fenyvesi É, Vieira MGA, Picos-Corrales LA, Moreno-Piraján JC, Giraldo L, Sohajda T, Huq MM, Soltan J, Torri G, Magureanu M, Bradu C, Crini G (2022) Removal of emerging contaminants from wastewater using advanced treatments. A review. *Environ Chem Lett* 20:1333–1375. <https://doi.org/10.1007/s10311-021-01379-5>
13. Jain B, Singh AK, Kim H, Lichtfouse E, Sharma VK (2018) Treatment of organic pollutants by homogeneous and heterogeneous Fenton reaction processes. *Environ Chem Lett* 16:947–967. <https://doi.org/10.1007/s10311-018-0738-3>
14. Kuznetsov ML, Kozlov YN, Mandelli D, Pombeiro AJL, Shul'pin GB (2011) Mechanism of Al^{3+} -catalyzed oxidations of hydrocarbons: dramatic activation of H_2O_2 toward O–O homolysis in complex $[\text{Al}(\text{H}_2\text{O})_4(\text{OOH})(\text{H}_2\text{O}_2)]^{2+}$ explains the formation of HO^\bullet radicals. *Inorg Chem* 50:3996–4005. <https://doi.org/10.1021/ic102476x>
15. Mandelli D, Chiacchio KC, Kozlov YN, Shul'pin GB (2008) Hydroperoxidation of alkanes with hydrogen peroxide catalyzed by aluminium nitrate in acetonitrile. *Tetrahedron Lett* 49:6693–6697. <https://doi.org/10.1016/j.tetlet.2008.09.058>
16. Mandelli D, Kozlov YN, da Silva CAR, Carvalho WA, Pescarmona PP, de Cella DA, de Paiva PT, Shul'pin GB (2016) Oxidation of olefins with H_2O_2 catalyzed by gallium(III) nitrate and aluminum(III) nitrate in solution. *J Mol Catal A Chem* 422:216–220. <https://doi.org/10.1016/j.molcata.2016.03.004>
17. Novikov AS, Kuznetsov ML, Rocha BGM, Pombeiro AJL, Shul'pin GB (2016) Oxidation of olefins with H_2O_2 catalysed by salts of group III metals (Ga, In, Sc, Y and La): epoxidation versus hydroperoxidation. *Catal Sci Technol* 6:1343–1356. <https://doi.org/10.1039/C5CY01367D>
18. Armstrong DA, Huie RE, Koppenol WH, Lyman SV, Merényi G, Neta P, Ruscic B, Stanbury DM, Steenken S, Wardman P (2015) Standard electrode potentials involving radicals in aqueous solution: inorganic radicals (IUPAC Technical Report). *Pure Appl Chem* 87:1139–1150. <https://doi.org/10.1515/pac-2014-0502>
19. Karst J, Sterl F, Linnenbank H, Weiss T, Hentschel M, Giessen H (2020) Watching in situ the hydrogen diffusion dynamics in magnesium on the nanoscale. *Sci Adv*. <https://doi.org/10.1126/sciadv.aaz0566>
20. Chen J, Wang Y, Su A (2024) Composition characteristics of magnesium isotopes in groundwater and their application prospects in water cycle processes. *Environ Earth Sci* 83:407. <https://doi.org/10.1007/s12665-024-11718-8>
21. Werner JJ, Arnold WA, McNeill K (2006) Water hardness as a photochemical parameter: tetracycline photolysis as a function of calcium concentration, magnesium concentration, and pH. *Environ Sci Technol* 40:7236–7241. <https://doi.org/10.1021/es060337m>
22. Patra SG, Mizrahi A, Meyerstein D (2020) The role of carbonate in catalytic oxidations. *Acc Chem Res* 53:2189–2200. <https://doi.org/10.1021/acs.accounts.0c00344>
23. Kottapurath Vijay A, Sharma VK, Meyerstein D (2023) Overlooked formation of carbonate radical anions in the oxidation of iron(II) by oxygen in the presence of bicarbonate. *Angew Chemie Int Ed*. <https://doi.org/10.1002/anie.202309472>
24. Illés E, Mizrahi A, Marks V, Meyerstein D (2019) Carbonate-radical-anions, and not hydroxyl radicals, are the products of the Fenton reaction in neutral solutions containing bicarbonate. *Free Radic Biol Med* 131:1–6. <https://doi.org/10.1016/j.freeradbiomed.2018.11.015>
25. Meyerstein D (2021) Re-examining Fenton and Fenton-like reactions. *Nat Rev Chem* 5:595–597. <https://doi.org/10.1038/s41570-021-00310-4>
26. Kottapurath Vijay A, Marks V, Mizrahi A, Wen Y, Ma X, Sharma VK, Meyerstein D (2023) Reaction of $\text{Fe}_{\text{aq}}^{\text{II}}$ with peroxymonosulfate and peroxydisulfate in the presence of bicarbonate: formation of $\text{Fe}_{\text{aq}}^{\text{IV}}$ and carbonate radical anions. *Environ Sci Technol* 57:6743–6753. <https://doi.org/10.1021/acs.est.3c00182>
27. Xiao R, Meng Y, Fu Y, Waclawek S, Wei Z, Spinney R, Dionysiou DD, Zeng W, Hu W (2023) The overlooked carbonate radical in micropollutant degradation: an insight into hydration interaction. *Chem Eng J* 474:145245. <https://doi.org/10.1016/j.cej.2023.145245>
28. Herscu-Kluska R, Masarwa A, Saphier M, Cohen H, Meyerstein D (2008) Mechanism of the reaction of radicals with peroxides and dimethyl sulfoxide in aqueous solution. *Chem—A Eur J* 14:5880–5889
29. Huie RE (2003) NDRL/NIST Solution Kinetics Database on the web, (n.d.), 108 No. 2 (2003)
30. Ulanski P, Merényi G, Lind J, Wagner R, von Sonntag C (1999) The reaction of methyl radicals with hydrogen peroxide. *J Chem Soc Perkin Trans 2*:673–676. <https://doi.org/10.1039/a900686i>
31. Burg A, Shamir D, Shusterman I, Kornweitz H, Meyerstein D (2014) The role of carbonate as a catalyst of Fenton-like reactions in AOP processes: $\text{CO}_3^{\cdot-}$ as the active intermediate. *Chem Commun* 50:13096–13099. <https://doi.org/10.1039/C4CC05852F>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.