



Some new soliton solutions to the higher dimensional Burger–Huxley and Shallow water waves equation with couple of integration architectonic

Farrah Ashraf^a, Tehsina Javeed^a, Romana Ashraf^a, Amina Rana^a, Ali Akgül^{b,c}, Shahram Rezapour^{d,e,*}, Muhammad Bilal Hafeez^{f,*}, Marek krawczuk^f

^a Department of Mathematics and Statistics, The University of Lahore, Lahore, Pakistan

^b Siirt University, Art and Science Faculty, Department of Mathematics, 56100 Siirt, Turkey

^c Near East University, Mathematics Research Center, Department of Mathematics, Near East Boulevard PC: 99138, Nicosia/Mersin 10, Turkey

^d Department of Mathematics, Azarbaijan Shahid Madani University, Tabriz, Iran

^e Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, Taiwan, China

^f Gdansk university of technology, Faculty of Mechanical Engineering and Ship Technology, Institute of Mechanic and Machine Design, Narutowicza 11/12, 80-233 Gdansk, Poland

ARTICLE INFO

Keywords:

Integrability
F-expansion scheme
Jacobi elliptic function

ABSTRACT

In this paper, we retrieve some traveling wave, periodic solutions, bell shaped, rational, kink and anti-kink type and Jacobi elliptic functions of Burger's equation and Shallow water wave equation with the aid of various integration schemes like improved F-expansion scheme and Jacobi elliptic function method respectively. We also present our solutions graphically in various dimensions.

Introduction

Nonlinear evolution equations (NLEEs) arising in nonlinear sciences play an important role in understanding nonlinear phenomenon. Solitons consist of huge applications in physics, communication systems, optical science, applied mathematics and engineering problems. The most arising and fast growing area of research is the optical solitons. A soliton or solitary wave is a self reinforcing wave packet that maintains its shape while it propagates at constant velocity. Solitons are caused by a cancellation of nonlinear and dispersive effects in the medium. Solitons are the solutions of a well-known class of weakly nonlinear dispersive partial differential equations describing physical systems. Soliton solutions have very interesting properties of some nonlinear partial differential equations. A soliton is a localized traveling wave solution of a nonlinear PDE that is remarkably stable [1–8]. In optical fibers, the solitons are the attractive field of research in applied mathematics, telecommunication systems and in distinctive branches of physics. A lot of research has been done on solutions of the solitons in nonlinear optics [9–14]. Many high-dimensional nonlinear evolution equations also contain some nonlinear structures such as the breath wave and lump solutions. In low-dimensional models, nonlinear waves can be classified from different aspects [15–17]. It is well known that the nonlinear Schrödinger (NLS) equation is a fundamental model in nonlinear physical systems, which plays a prominent role in a wide range of physical subjects such as plasma physics. This work not only

reveals the characteristics of single bright–bright soliton by theoretical analysis, but also describes a wealth of new phenomena of two, three, and four bright–bright solitons, including elastic collision, soliton reflection, parallel propagation, time-periodic propagation, and (space, time)-periodic propagation [18–20].

The objective of this article is to execute the Improved F-expansion method in accomplishing the traveling wave arrangements to NLEEs within the mathematical physics through the DSW condition and the Burgers condition in terms of functions that fulfill the Riccati condition $F(\xi) = k + F^2(\xi)$.

The Burgers condition is the least order estimation for the one-dimensional proliferation of weak waves in a fluid. It is additionally utilized in vehicle frequency in high way activity. It is one of the fundamental PDEs in fluid mechanics. Burgers condition is totally fundamental. The wave arrangements of Burgers equation are single and multiple-front arrangements (Wazwaz, 2009). The DSW condition is an imperative wave demonstrate in material science (Inc, 2006).

Burgers condition was to begin with presented at Bateman within 1915 and afterward analyzed at Burgers within 1948. The condition is utilized as a demonstrate in numerous areas such as, ceaseless stochastic forms [21] dissipate water [22] stun waves, warm [23]. Burgers condition can onto considered to a rearranged frame of the N-S condition onto frame item or the event consistency term [24]. Unused correct arrangements of the common Burgers condition are

* Corresponding authors.

E-mail addresses: rezapourshahram@yahoo.ca (S. Rezapour), muhammad.bilal.hafeez@pg.edu.pl (M.B. Hafeez).

independently determined at a coordinate strategy or only condition to the Bernoulli condition come the only condition [25] individually. The couple correct arrangements are appeared will be equal [26]. In expansion, the recently determined arrangement can be effectively decreased to straight [27] whereas the nonlinear term coefficient equal to zero [28]. At last, inferred arrangement is compared with the irritation arrangement and some existing correct arrangements [29]. A few numerical comes about are displayed and outlined [30,31].

The generalized Burger–Huxley (gBH) equation is a first-order nonlinear partial differential equation (NPDE) in time t that allows for the propagation of a single-wave (single-mode) solution. The generalized Burger–Huxley condition may be a non-linear fractional differential condition of first arrange in time t and peruses [32]:

$$U_t - U_{xx} - \mu U U_x - \nu U(1 - U)(U - \gamma) = 0 \tag{1}$$

By wave transformation

$$U = U(\xi), \quad \xi = x + ct \tag{2}$$

where $\phi = \phi(x, t)$ and μ, ν, γ are common genuine scalars.

For $\mu = 0$ and $\gamma = 1$, gBH is reduced to the Huxley model, representing nerve pulse propagation in nerve fibers and liquid crystal wall motion (Wang et al. 1990). Eq. (1) is the Burgers' equation, which describes the field of wave propagation in nonlinear dissipative models for $\nu = 0$ and $\gamma = 1$. Burger–Huxley Eq. (1) describes how convection terms interact with diffusion transmission (Wang et al. 1990).

We will look at a partial differential equation with non-linearity, a weakly nonlinear shallow water wave equation used to study wave propagation in dispersive and weakly nonlinear media.

The model under consideration is:

$$U_{tt} - \lambda \sigma U_{xx} + \frac{1}{2\sigma} (U_t^2)_x - \frac{\sigma^2}{3} U_{xxx} = 0, \tag{3}$$

By wave transformation

$$U = U(\xi), \quad \xi = x + ct \tag{4}$$

where $U = U(x, t)$ represents the unification of the displacement and velocity of the water particles, the gravitational force is represented by λ . The wave height is represented by σ . The Euler equation of motion, the equation of dynamics and mass conservation, and kinematics constraints on the boundary of the domain consisting of the free surface have essentially illustrated the water problem for gravity waves.

We applied two integration schemes, namely IFE scheme [33] and JEF method [34] in this paper. IEF method is applicable to get some solitary wave, periodic wave and rational function solutions and JEF method provide Jacobi elliptic function solutions.

The paper is written in the following sequence: in Section “Contents of Integration Schemes”, we summarize IFE and JEF scheme. In Section “Mathematical analysis”, we use IFE and JEF schemes to obtain some solitary, periodic wave, bell shaped, kink and anti kink type solutions, rational solutions and Jacobi elliptic solutions in terms of hyperbolic and trigonometric functions with their graphical representations. in Section “Results and discussion” we discuss our results and lastly in Section “Conclusion”, we conclude our results.

Contents of integration schemes

Analysis of IFE method

Consider NLEE in the form [33].

$$G(U, U_x, U_y, U_t, U_{xx}, U_{xy}, U_{xt}, \dots) = 0,$$

where

$$U(x, y, t) = U(\xi), \quad \xi = \mu x + \nu y + \epsilon t.$$

Replace in above equation to get ODE.

$$H(U, U', U'', U''', \dots) = 0, \quad U = U(\xi).$$

Now we use following transformation,

$$U(\xi) = \sum_{i=0}^m a_i F^i + \sum_{i=1}^n b_i F^{-i},$$

where m can be find through homogeneous balance and a_i' and b_i' are to be determined. We know the Riccati equation $F'(\xi) = k + F^2(\xi)$. Substitute these transformations into first equation to get periodic wave and solitary wave solutions. Now we represent the general solution of Riccati equation [35].

When $k < 0$,

$$F_1 = -\sqrt{-k} \tanh(-\sqrt{-k}\xi),$$

$$F_2 = -\sqrt{-k} \coth(-\sqrt{-k}\xi).$$

When $k > 0$,

$$F_3 = \sqrt{k} \tan(\sqrt{k}\xi),$$

$$F_4 = \sqrt{k} \cot(\sqrt{k}\xi).$$

For $k = 0$,

$$F_5 = \frac{-1}{\xi}.$$

Analysis of JEF method

We have nonlinear evolution equation [14].

$$G(U, U_t, U_x, U_{xt}, \dots) = 0.$$

We use following transformations.

$$U(x, y, t) = U(\xi), \quad \xi = \alpha x + \beta y + \epsilon t,$$

this can be transform into nonlinear ODE.

$$G_0(U, U_\xi, U_{\xi\xi}, \dots) = 0.$$

Now use these transformations.

$$U(\xi) = \sum_{i=0}^m d_i F^i,$$

in which d_i 's are to be determined where $i = 0, 1, 2, \dots, m$. By homogeneous balance we find “ m ” and $F(\xi)$ express in following substitution. $F' = \sqrt{r + dF^2 + \frac{eF^4}{2} + \frac{cF^6}{3}}$ where r, c, d , and e are real parameters.

We insert F' in previous transformations to obtain system of equations then obtain the values of d_i 's.

If we assume $F(\xi) = \text{sn}\xi, \text{cn}\xi, \text{cs}\xi$, its called JEF method.

Mathematical analysis

Putting Eq. (2) along its derivatives in Eq. (1) we get,

$$CU' - U'' - \mu U U' - \nu u^2 + \nu U \gamma + \nu U^3 - \nu U^2 \gamma = 0, \tag{5}$$

By comparing highest nonlinear term with highest derivative, we get $m = 1$ through homogeneous balance.

IFE method

We use following transformation [33],

$$U(\xi) = \sum_{i=0}^m a_i F^i + \sum_{i=1}^n b_i F^{-i},$$

$$U(\xi) = a_0 + a_1 F + b_1 F^{-1}. \tag{6}$$

Putting Eq. (5) along its derivatives in Eq. (4), we get

$$C(a_1K + a_1F^2 - \frac{b_1K}{F^2} - b_1) - \mu(a_1K + a_1F^2 - \frac{b_1K}{F^2} - b_1) \\ \cdot (a_0 + a_1F + b_1F^{-1}) - (2a_1FK - 2a_1F^3 - \frac{2b_1K^2}{F^3} - \frac{2b_1K}{F}) \\ - v(a_0 + a_1F + b_1F^{-1})^2 + v\gamma(a_0 + a_1F + b_1F^{-1}) + v(a_0 + a_1F + b_1F^{-1})^3 \\ - v\gamma(a_0 + a_1F + b_1F^{-1})^2 = 0 \tag{7}$$

where $F' = k + F^2$ (see Figs. 1–20). Setting the coefficients of each power of $F(\xi)$ to zero as follows.

$$-2a_1 - \mu a_1^2 + v a_1^3, \tag{8}$$

$$c a_1 + 3v a_0 a_1^2 - \mu a_1 a_0 - v a_1^2 - v \gamma a_1^2, \tag{9}$$

$$v \gamma a_1 - 2a_1 K - \mu(a_1 K - b_1) a_1 - \mu a_1 b_1 \\ + v(b_1 a_1^2 + 2a_0^2 a_1 + a_1(2a_1 b_1 + a_0^2)) - 2v \gamma a_0 a_1 - 2v a_0 a_1, \tag{10}$$

$$v \gamma a_0 + c(a_1 K - b_1) - \mu(a_1 K - b_1) a_0 - v \gamma(2a_1 b_1 + a_0^2) \\ + v(4a_0 a_1 b_1 + a_0(2a_1 b_1 + a_0^2)) - v(2a_1 b_1 + a_0^2), \tag{11}$$

$$-2v b_1 a_0 + v(b_1(2a_1 b_1 + a_0^2) + 2a_0^2 b_1 + a_1 b_1^2) + \mu b_1 K a_1 \\ - \mu(a_1 K - b_1) b_1 + v \gamma b_1 - 2b_1 K - 2v \gamma b_1 a_0, \tag{12}$$

$$3v b_1^2 a_0 - c b_1 K - \beta b_1^2 + \mu b_1 K a_0 - v \gamma b_1^2, \tag{13}$$

$$-2b_1 K^2 + v b_1^3 + \mu b_1^2 K. \tag{14}$$

We get,

Set 1:

$$K = \frac{-1}{16a_1^2}, \quad c = \frac{-1 + v \gamma a_1^2}{a_1}, \quad \mu = \frac{-2 + \beta a_1^2}{a_1} \quad a_1 = a_1, \quad a_0 = \frac{1}{2},$$

$$b_1 = \frac{1}{16a_1}.$$

we get

$$U_1(\xi) = \frac{1}{2} + a_1 F + \frac{1}{16a_1} F^{-1}.$$

When $k < 0$,

$$U_1 = \frac{1}{2} - a_1 \sqrt{-k} \tanh(\sqrt{-k}\xi) - \frac{1}{16a_1(\sqrt{-k} \tanh(\sqrt{-k}\xi))}, \tag{15}$$

and

$$U_1 = \frac{1}{2} - a_1 \sqrt{-k} \coth(\sqrt{-k}\xi) - \frac{1}{16a_1(\sqrt{-k} \coth(\sqrt{-k}\xi))}. \tag{16}$$

When $k > 0$,

$$U_1 = \frac{1}{2} + a_1(\sqrt{k} \tan \sqrt{k}\xi) + \frac{1}{16a_1(\sqrt{k} \tan(\sqrt{k}\xi))}, \tag{17}$$

and

$$U_1 = \frac{1}{2} - a_1(\sqrt{k} \cot \sqrt{k}\xi) - \frac{1}{16a_1(\sqrt{k} \cot(\sqrt{k}\xi))}. \tag{18}$$

When $k = 0$,

$$U_1 = \frac{1}{2} - a_1\left(\frac{1}{\xi}\right) + \frac{1}{16a_1}\left(-\frac{1}{\xi}\right)^{-1}. \tag{19}$$

Set 2:

$$k = k, \quad a_0 = (\mu^2 - v^2)k, \quad a_1 = 0, \quad a_2 = 0, \quad b_1 = 0, \quad b_2 = (\mu^2 - v^2)k^2.$$

we get

$$U_2(\xi) = (\mu^2 - v^2)k + (\mu^2 - v^2)k^2 F^{-2}. \tag{20}$$

When $k < 0$,

$$U_2 = (\mu^2 - v^2)k - (\mu^2 - v^2)k^3(\coth(-\sqrt{-k}\xi))^2, \tag{21}$$

and

$$U_2 = (\mu^2 - v^2)k - (\mu^2 - v^2)k^3(\tanh(-\sqrt{-k}\xi))^2. \tag{22}$$

When $k > 0$,

$$U_2 = (\mu^2 - v^2)k + (\mu^2 - v^2)k^3(\cot(\sqrt{k}\xi))^2, \tag{23}$$

and

$$U_2 = (\mu^2 - v^2)k + (\mu^2 - v^2)k^3(\tan(\sqrt{k}\xi))^2. \tag{24}$$

When $k = 0$,

$$U_2 = (\mu^2 - v^2)k + (\mu^2 - v^2)k^2 \frac{1}{\xi^2}. \tag{25}$$

Set 3:

$$k = k, \quad a_0 = 2(\mu^2 - \nu^2)k, \quad a_1 = 0, \quad a_2 = \mu^2 - \nu^2, \quad b_1 = 0, \quad b_2 = (\mu^2 - \nu^2)k^2.$$

we get

$$U_3(\xi) = 2(\mu^2 - \nu^2)k + (\alpha^2 - \beta^2)F^2 + (\mu^2 - \nu^2)k^2 F^{-2}. \tag{26}$$

When $k < 0$,

$$U_3 = 2(\mu^2 - \nu^2)k - (\mu^2 - \nu^2)k(\tanh(-\sqrt{-k}\xi))^2 - (\mu^2 - \nu^2)k^3(\coth(-\sqrt{-k}\xi))^2, \tag{27}$$

and

$$U_3 = 2(\mu^2 - \nu^2)k - (\mu^2 - \nu^2)k(\coth(-\sqrt{-k}\xi))^2 - (\mu^2 - \nu^2)k^3(\tanh(-\sqrt{-k}\xi))^2. \tag{28}$$

When $k > 0$,

$$U_3 = 2(\mu^2 - \nu^2)k + (\mu^2 - \nu^2)k(\tan(\sqrt{k}\xi))^2 - (\mu^2 - \nu^2)k^3(\cot(\sqrt{k}\xi))^2, \tag{29}$$

and

$$U_3 = 2(\mu^2 - \nu^2)k + (\mu^2 - \nu^2)k(\cot(\sqrt{k}\xi))^2 - (\mu^2 - \nu^2)k^3(\tan(\sqrt{k}\xi))^2. \tag{30}$$

When $k = 0$,

$$U_3 = 2(\mu^2 - \nu^2)k + (\mu^2 - \nu^2)k \frac{1}{\xi^2} + (\mu^2 - \nu^2)k^2 \xi^2. \tag{31}$$

Set 4:

$$k = k, \quad a_0 = \frac{-2}{3}(\mu^2 - \nu^2)k, \quad a_1 = 0, \quad a_2 = (\mu^2 - \nu^2) \quad b_1 = 0, \quad b_2 = (\mu^2 - \nu^2)k^2.$$

we get

$$U_4(\xi) = \frac{-2}{3}(\mu^2 - \nu^2)k + (\mu^2 - \nu^2)F^2 + (\mu^2 - \nu^2)k^2 F^{-2}. \tag{32}$$

When $k < 0$,

$$U_4 = \frac{-2}{3}(\mu^2 - \nu^2)k - (\mu^2 - \nu^2)k(\tanh(-\sqrt{-k}\xi))^2 - (\mu^2 - \nu^2)k^3(\coth(-\sqrt{-k}\xi))^2, \tag{33}$$

and

$$U_4 = \frac{-2}{3}(\mu^2 - \nu^2)k - (\mu^2 - \nu^2)k(\coth(-\sqrt{-k}\xi))^2 - (\mu^2 - \nu^2)k^3(\tanh(-\sqrt{-k}\xi))^2. \tag{34}$$

When $k > 0$,

$$U_4 = \frac{-2}{3}(\mu^2 - \nu^2)k - (\mu^2 - \nu^2)k(\tan(\sqrt{k}\xi))^2 - (\mu^2 - \nu^2)k^3(\cot(\sqrt{k}\xi))^2, \tag{35}$$

and

$$U_4 = \frac{-2}{3}(\mu^2 - \nu^2)k - (\mu^2 - \nu^2)k(\cot(\sqrt{k}\xi))^2 - (\mu^2 - \nu^2)k^3(\tan(\sqrt{k}\xi))^2. \tag{36}$$




Fig. 1. The graphical presentation of $U_1(\xi)$ given by Eq. (13).




Fig. 2. The dynamical behavior of the solution $U_1(\xi)$ given by Eq. (14).




Fig. 3. The shape profile of $U_1(\xi)$ given by Eq. (15).




Fig. 4. The graphical presentation of $U_1(\xi)$ given by Eq. (16).




Fig. 5. The shape profile of $U_2(\xi)$ given by Eq. (17).




Fig. 6. The shape profile of $U_2(\xi)$ given by Eq. (19).




Fig. 7. The graphical presentation of $U_2(\xi)$ given by Eq. (20).




Fig. 8. The shape profile of $U_2(\xi)$ given by Eq. (21).




Fig. 9. The graphical presentation of $U_2(\xi)$ given by Eq. (22).




Fig. 10. The graphical presentation of $U_2(\xi)$ given by Eq. (23).




Fig. 11. The shape profile of $U_3(\xi)$ given by Eq. (25).




Fig. 12. The graphical presentation of $U_2(\xi)$ given by Eq. (26).




Fig. 13. The graphical presentation of $U_2(\xi)$ given by Eq. (27).




Fig. 14. The graphical presentation of $U_2(\xi)$ given by Eq. (28).




Fig. 15. The graphical presentation of $U_2(\xi)$ given by Eq. (29).




Fig. 16. The shape profile of $U_4(\xi)$ given by Eq. (31).




Fig. 17. The graphical presentation of $U_2(\xi)$ given by Eq. (32).




Fig. 18. The graphical presentation of $U_2(\xi)$ given by Eq. (33).




Fig. 19. The graphical presentation of $U_2(\xi)$ given by Eq. (34).




Fig. 20. The graphical presentation of $U_2(\xi)$ given by Eq. (35).




Fig. 21. The graphical presentation of $U(\xi)$ given by Eq. (44).

When $k = 0$,

$$U_4 = \frac{-2}{3}(\mu^2 - v^2)k - (\mu^2 - v^2)\frac{1}{\xi^2} + (\mu^2 - v^2)k^2\xi^2. \tag{37}$$

JEF method

For JEF method, we use this transformation [36].

$$U(\xi) = \sum_{i=0}^m a_i F^i$$

$$U(\xi) = a_0 + a_1 F + a_2 F^2 \tag{38}$$

Putting Eq. (4) along its derivatives in Eq. (3) and set $F' = \sqrt{r + aF^2 + \frac{bF^4}{2} + \frac{cF^6}{3}}$ in Eq. (3), then we get,

$$a_1 a F + a_1 b F^3 + a_1 c F^5 + 2a_2 r + 4a_2 a F^2 + 3a_2 b F^4 + \frac{8}{3} a_2 c F^6 - b_1 a F^{-1} - b_1 F^3 c - b_1 F b$$

$$-\tau(a_0 + a_1 F + a_2 F^2) - \xi(a_0 + a_1 F + a_2 F^2)^2 = 0. \tag{39}$$

Setting the coefficients of each power of $F(\xi)$ to zero as follows.

$$3a_2 b - \xi a_2^2, \tag{40}$$

$$-b_1 + a_1 b - 2\xi a_1 a_2, \tag{41}$$

$$-\xi(2a_0 a_2 + a_1^2) + 4a_2 a - \tau a_2, \tag{42}$$

$$-2\xi a_0 a_1 - b_1 b + a_1 a - \tau a_1, \tag{43}$$

$$-\xi a_0^2 - \tau a_0 + 2a_2 r. \tag{44}$$

By solving simultaneously above equations we get,

$$c = \frac{b^2}{r}, \quad r = r, \quad \tau = 0, \quad \xi = 0, \quad a_0 = a_0, \quad a_1 = \frac{b_1 b}{a}, \quad a_2 = 0.$$

So,

$$U(\xi) = U = a_0 + \frac{b_1 b}{a} F. \tag{45}$$

Depending on a, b, c, r in Eq. (F'), we obtained different type of traveling wave solutions.

Case 1:

$$a = -(1 + m^2), \quad b = 2m^2, \quad r = 1, \quad c = 0.$$

We get,

$$U = a_0 - \frac{2b_1 m^2}{1 + m^2} sn(\xi, m),$$

As $m \rightarrow 1$ this generate shock wave (see Fig. 21).

$$U = a_0 - \frac{2b_1 m^2}{1 + m^2} tanh(\xi). \tag{46}$$

Case 2:

$$a = 2m^2 - 1, \quad b = 2, \quad r = -m(1 - m^2), \quad c = 0.$$

We obtained,

$$U = a_0 + \frac{2b_1}{2m^2 - 1} ds(\xi, m)$$

As $m \rightarrow 1$ this generate (see Fig. 22).

$$U = a_0 + \frac{2b_1}{2m^2 - 1} cosech(\xi) \tag{47}$$

Case 3:

$$a = 2 - m^2, \quad b = 2, \quad r = 1 - m^2, \quad c = 0.$$

We get,

$$U = a_0 + \frac{2b_1}{2 - m^2} cs(\xi, m),$$

As $m \rightarrow 1$ this generate (see Fig. 23).

$$U = a_0 + \frac{2b_1}{2 - m^2} csch(\xi). \tag{48}$$

Case 4:

$$d = 2m^2 - 1, \quad e = -2m^2, \quad r = (1 - m^2), \quad c = 0.$$

We obtained,

$$U(\xi) = \frac{-2(2m^2 - 1)\beta^3 + 2(2m^2 - 1)\alpha^2\beta - 2e}{6} - \frac{-2m^2(\alpha^2 - \beta^2)}{2} cn^2(\xi, m).$$

As $m \rightarrow 1$ this generate (see Fig. 24).

$$U(\xi) = \frac{-2\beta^3 + 2\alpha^2\beta - 2e}{6} + (\alpha^2 - \beta^2) sech^2(\xi). \tag{49}$$

Case 5:

$$d = 2 - m^2, \quad e = -2, \quad r = (m^2 - 1), \quad c = 0.$$

We get,

$$U(\xi) = \frac{-2(2 - m^2)\beta^3 + 2(2 - m^2)\alpha^2\beta - 2e}{6} - \frac{-2(\alpha^2 - \beta^2)}{2} dn^2(\xi, m).$$

As $m \rightarrow 1$ this generate (see Fig. 25).

$$U(\xi) = \frac{-2\beta^3 + 2\alpha^2\beta - 2e}{6} + (\alpha^2 - \beta^2) sech^2(\xi). \tag{50}$$

Case 6:

$$d = \frac{m^2 - 2}{2}, \quad e = \frac{m^2}{2}, \quad r = \frac{1}{4}, \quad c = 0.$$

We get,

$$U(\xi) = \frac{-2(\frac{m^2 - 2}{2})\beta^3 + 2(\frac{m^2 - 2}{2})\alpha^2\beta - 2e}{6} - \frac{\frac{m^2}{2}(\alpha^2 - \beta^2)}{2} \frac{sn^2(\xi, m)}{(1 \pm dn(\xi, m))^2}.$$

As $m \rightarrow 1$ this generate (see Figs. 26 and 27).

$$U(\xi) = \frac{\beta^3 - \alpha^2\beta - 2e}{6} - \frac{(\alpha^2 - \beta^2)}{4} \frac{tanh^2(\xi)}{(1 \pm sech(\xi))^2}. \tag{51}$$




Fig. 22. Graphical presentation of $U(\xi)$ given by Eq. (45).




Fig. 23. The shape profile of $U(\xi)$ given by Eq. (46).




Fig. 24. The graphical presentation of $U(\xi)$ given by Eq. (47).




Fig. 25. The shock wave of $U(\xi)$ given by Eq. (48).

Case 7:

$$d = \frac{m^2-2}{2}, \quad e = \frac{m^2}{2}, \quad r = \frac{m^2}{4}, \quad c = 0.$$

We get,

$$U(\xi) = \frac{-2(\frac{m^2-2}{2})\beta^3 + 2(\frac{m^2-2}{2})\alpha^2\beta - 2e}{6} - \frac{\frac{m^2}{2}(\alpha^2 - \beta^2)}{2}$$

$$\times \frac{dn^2(\xi, m)}{(m^2 + 1)^2(sn^2(\xi, m) \pm dn(\xi, m))^2}.$$

As $m \rightarrow 1$ this generate (see Figs. 28 and 29).

$$U(\xi) = \frac{\beta^3 - \alpha^2\beta - 2e}{6} - \frac{(\alpha^2 - \beta^2)}{4} \frac{\text{sech}^2(\xi)}{(\tanh^2(\xi) \pm \text{sech}(\xi))^2}. \tag{52}$$




Fig. 26. The shock wave of $U(\xi)$ given by Eq. (49).




Fig. 27. The shock wave of $U(\xi)$ given by Eq. (49).




Fig. 28. The shock wave of $U(\xi)$ given by Eq. (50).




Fig. 29. The shock wave of $U(\xi)$ given by Eq. (50).




Fig. 30. The shock wave of $U(\xi)$ given by Eq. (51).




Fig. 31. The shock wave of $U(\xi)$ given by Eq. (51).

Case 8:

$$d = \frac{m^2+1}{2}, \quad e = \frac{-1}{2}, \quad r = \frac{-(1-m^2)^2}{4}, \quad c = 0.$$

We get,

$$U(\xi) = \frac{-2(\frac{m^2+1}{2})\beta^3 + 2(\frac{m^2+1}{2})\alpha^2\beta - 2\epsilon}{6} - \frac{\frac{-1}{2}(\alpha^2 - \beta^2)}{2} (m\text{cn}(\xi, m) \pm \text{dn}(\xi, m))^2.$$

As $m \rightarrow 1$ this generate (see Figs. 30 and 31).

$$U(\xi) = \frac{-2\beta^3 + 2\alpha^2\beta - 2\epsilon}{6} + \frac{(\alpha^2 - \beta^2)}{4} (\text{sech}(\xi) \pm \text{sech}(\xi))^2. \tag{53}$$

Case 9:

$$d = \frac{m^2+1}{2}, \quad e = \frac{m^2+1}{2}, \quad r = \frac{m^2-1}{4}, \quad c = 0.$$

We get,

$$U(\xi) = \frac{-2(\frac{m^2+1}{2})\beta^3 + 2(\frac{m^2+1}{2})\alpha^2\beta - 2\epsilon}{6} - \frac{(\frac{m^2+1}{2})(\alpha^2 - \beta^2)}{2} \frac{\text{dn}^2(\xi, m)}{(1 \pm m\text{sn}(\xi, m))^2}.$$

As $m \rightarrow 1$ this generate (see Figs. 32 and 33).

$$U(\xi) = \frac{-2\beta^3 + 2\alpha^2\beta - 2\epsilon}{6} - \frac{(\alpha^2 - \beta^2)}{2} \frac{\text{sech}^2(\xi, m)}{(1 \pm \tanh(\xi))^2}. \tag{54}$$

Case 10:

$$d = \frac{m^2+1}{2}, \quad e = \frac{1-m^2}{2}, \quad r = \frac{1-m^2}{4}, \quad c = 0.$$

We obtain,

$$U(\xi) = \frac{-2(\frac{m^2+1}{2})\beta^3 + 2(\frac{m^2+1}{2})\alpha^2\beta - 2\epsilon}{6} - \frac{(\frac{1+m^2}{2})(\alpha^2 - \beta^2)}{2} \frac{\text{cn}^2(\xi, m)}{(1 \pm m\text{sn}(\xi, m))^2}.$$

As $m \rightarrow 1$ this generate (see Figs. 34 and 35).

$$U(\xi) = \frac{-2\beta^3 + 2\alpha^2\beta - 2\epsilon}{6} - \frac{(\alpha^2 - \beta^2)}{2} \frac{\text{sech}^2(\xi, m)}{(1 \pm \tanh(\xi))^2}. \tag{55}$$

Case 11:

$$d = \frac{m^2+1}{2}, \quad e = \frac{(1-m^2)^2}{2}, \quad r = \frac{1}{4}, \quad c = 0.$$

We get,

$$U(\xi) = \frac{-2(\frac{m^2+1}{2})\beta^3 + 2(\frac{m^2+1}{2})\alpha^2\beta - 2\epsilon}{6} - \frac{(\frac{1+m^2}{2})(\alpha^2 - \beta^2)}{2} \times \frac{\text{sn}^2(\xi, m)}{(\text{dn}(\xi, m) \pm m\text{sn}(\xi, m))^2}.$$

As $m \rightarrow 1$ this generate (see Fig. 36).

$$U(\xi) = \frac{-2\beta^3 + 2\alpha^2\beta - 2\epsilon}{6} - \frac{(\alpha^2 - \beta^2)}{2} \frac{\tanh^2(\xi)}{(\text{sech}(\xi) \pm \text{sech}(\xi))^2}. \tag{56}$$

Results and discussion

In this manuscript we use IFE and JEF to get solitary wave, periodic wave, bell-shaped, kink and anti-kink type, rational and Jacobi elliptic solutions in terms of hyperbolic and trigonometric functions and plot their 3D and contours graphs.

Conclusion

We obtained various types of solutions i.e solitary wave, periodic wave, bell shaped, kink and anti-kink type, rational solutions and Jacobi elliptic functions of Burger's and weakly nonlinear Shallow water wave differential equation. In this paper, we used IFE and JEF architectonic. First method gives solitary wave solutions, bell shaped, kink and anti-kink type solutions and second one gives Jacobi elliptic functions in terms of hyperbolic and trigonometric functions and rational solutions with their graphical representations like 3-D and contours graphs. Furthermore, our constructed solutions illustrate how simple, reliable, and consistent this method's solution process is existing solitary wave solutions, single soliton solutions, periodic solutions, and kink solutions are obtained if the parameters adopt particular values. The finding appear that the improved F-expansion approach could be a promising device since it can offer a number of solutions with different one of a kind physical contour. To the best of our knowledge, these results were obtained first time for these models.




Fig. 32. The graphical presentation of $U(\xi)$ given by Eq. (52).




Fig. 33. The graphical presentation of $U(\xi)$ given by Eq. (52).




Fig. 34. The graphical presentation of $U(\xi)$ given by Eq. (53).




Fig. 35. The graphical presentation of $U(\xi)$ given by Eq. (53).




Fig. 36. The graphical presentation of $U(\xi)$ given by Eq. (54).

CRedit authorship contribution statement

Farrah Ashraf: Conceptualization. **Tehsina Javeed:** Supervision. **Romana Ashraf:** Validation. **Amina Rana:** Writing first draft. **Ali Akgül:** Investigation.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

Acknowledgment

All authors read and approved the final manuscript.

References

- [1] Rizvi STR, Seadawy AR, Ashraf F, Younis M, Iqbal H, Baleanu D. Lump and interaction solutions of a geophysical Korteweg–de Vries equation. *Results Phys* 2020;19:103661.
- [2] Ashraf Farrah, Seadawy Aly R, Rizvi Syed TR, Ali Kashif, Aamir Ashraf M. Multi-wave, M-shaped rational and interaction solutions for fractional nonlinear electrical transmission line equation. *J Geom Phys* 2022;177:104503.
- [3] Rizvi STR, Seadawy AR, Ashraf F, Younis M, Iqbal H, Baleanu D. Kinky breathers, W-shaped and multi-peak solitons interaction in (2+ 1)-dimensional nonlinear Schrödinger equation with Kerr law of nonlinearity. *Eur Phys J Plus* 2019;134:1–10.
- [4] Rizvi STR, Seadawy Aly R, Ashraf R. Propagation of chirped periodic and solitary waves for the coupled nonlinear Schrödinger equation in two core optical fibers with parabolic law with weak non-local nonlinearity. *Opt Quantum Electron* 2022;54:545.
- [5] Ahmed S, Ashraf R, Seadawy Aly R, Rizvi STR, Younis M, Althobaiti Ali, El-Shehawi Ahmed M. Lump, multi-wave, kinky breathers, interactional solutions, and stability analysis for general (2+1)-th dispersionless Dym equation. *Results Phys* 2021;25:104160.
- [6] Ashraf R, Ahmad MO, Younis M, Tariq KU, Ali K, Rizvi STR. Dipole and combo solitons in DWDM systems. *Optik* 2018;158:1073–9.
- [7] Ashraf R, Ahmad MO, Younis M, Ali K, Rizvi STR. Dipole and Gausson soliton for ultrashort laser pulse with high order dispersion. *Superlattices Microstruct* 2017;109:504–10.
- [8] Rizvi STR, Bashir S, Younis M, Ashraf R, Ahmad MO, Ahmad. Exact soliton of (2 + 1)-dimensional fractional Schrödinger equation. *Superlattices Microstruct* 2017;107:234–9.
- [9] Ablowitz MJ, Clarkson PA. *Solitons, nonlinear evolution equations and inverse scattering*, Vol. 149. Cambridge University Press; 1991.
- [10] El-Wakil SA, Abdou MA. New exact travelling wave solutions using modified extended tanh-function method. *Chaos Solitons Fractals* 2007;840–52.
- [11] Wazwaz AM. Multiple-soliton solutions of two extended model equations for shallow water waves. *Appl Math Comput* 2008;790–9.
- [12] Bekir A. New solitons and periodic wave solutions for some nonlinear physical models by using the Sine-Cosine method. *Phys Scr* 2008;501.
- [13] He JH, Wu XH. Exp-function method for nonlinear wave equations. *Chaos Solitons Fractals* 2006;700.
- [14] Elboree MK. The Jacobi elliptic function method and its application for two component BKP hierarchy equations. *Comput Math Appl* 2011;4402–14.
- [15] Yin Zhe-Yong, Tian Shou-Fu. Nonlinear wave transitions and their mechanisms of (2+1)-dimensional Sawada–Kotera equation. *Physica D* 2021;427:133002.
- [16] Li Tian, Yang. Soliton resolution for the Wadati–Konno–Ichikawa equation with weighted Sobolev initial data. *Henri Poincaré* 2022;23:2611–55.
- [17] Shou-Fu Tian, Mei-Juan Xu, Tian-Tian Zhang. A symmetry-preserving difference scheme and analytical solutions of a generalized higher-order beam equation. *Proc R Soc Lond Ser A Math Phys Eng Sci* 2021;477:0455.
- [18] Li Yuan, Tian Shou-Fu, Yang Jin-Jie, Riemann. Hilbert problem and interactions of solitons in the n-component nonlinear Schrödinger equations. *Stud Appl Math* 2022;148:577–605.
- [19] Fatih Ozbag, Mahmut Modanli. On the stability estimates and numerical solution of fractional order telegraph integro-differential equation. *Phys Scr* 2021;96(9).
- [20] Modanli Mahmut, Ozbag Fatih, Akgul Ali. Finite difference method for the fractional order Pseudo telegraph integro-differential equation. *J Appl Math Comput Mech* 2022;21(1):41–54.
- [21] Sugimoto N. Burgers equation with a fractional derivative; hereditary effects on nonlinear acoustic waves. *J Fluid Mech* 1991;225:631–53.
- [22] Weinan E, Khanin K, Mazel A, Sinai Y. Invariant measures for Burgers equation with stochastic forcing. *Ann of Math* 2000;151(3):877–960.
- [23] Burgers JM. A mathematical model illustrating the theory of turbulence. *Adv Appl Mech* 1948;1:171–99.
- [24] Wang XY, Zhu ZS, Lu YK. Solitary wave solutions of the generalized Burgers Huxley equation. *J Phys A: Math Gen* 1990;23:271–4.
- [25] Wang GW, Liu XQ, Zhang YY. New explicit solutions of the generalized Burgers Huxley equation. *J Vietnam Math* 2013;41:161–6.
- [26] Yemova OY, Kudryashov NA. Exact solutions of the Burgers-Huxley equation. *J Appl Math Mech* 2004;68(3):413–20.
- [27] Wazwaz AM. Analytic study on Burgers, Fisher, Huxley equations and combined forms of these equations. *Appl Math Comput* 2008;195:754–61.
- [28] Wazwaz AM. Travelling wave solutions of generalized forms of Burgers, Burgers-KdV and Burgers-Huxley equations. *Appl Math Comput* 2005;169:639656.
- [29] Korsunsky SV. Soliton solutions for a second-order KdV equation. *Phys Lett A* 1994;185:174–6.
- [30] Wazwaz AM. Two-mode fifth-order KdV equations: necessary conditions for multiple soliton solutions to exist. *Nonlinear Dynam* 2017;87(3):1685–91.
- [31] Biazar J, Mohammadi F. Application of differential transform method to the generalized Burgers-Huxley equation. *Appl Appl Math* 2010;5:1726–40.
- [32] Dehghan M, Saray BN, Lakestani M. Three methods based on the interpolation scaling functions and the mixed collocation finite difference schemes for the numerical solution of the nonlinear generalized Burgers-Huxley equation. *Math Comput Model* 2012;55:1129–42.
- [33] Akbar MA, Ali NHM. The improved F-expansion method with Riccati equation and its applications in mathematical physics. *Cogent Math* 2017;1282577.
- [34] Muhiameed ZIAA, Salam Emad ABA. Generalized Jacobi elliptic function solution to a class of nonlinear Schrödinger-type equations. *Math Probl Eng* 2011.
- [35] Karaman B. The use of improved F-expansion method for the time-fractional Benjamin–Ono equation. *RACSAM* 2021;115:128.
- [36] Zheng B. A new fractional Jacobi elliptic equation method for solving fractional partial differential equations. *Adv Difference Equ* 2014;2014:228.