



## Regular Articles

Versatility of putative aromatic aminotransferases from *Candida albicans*

Kamila Rząd, Sławomir Milewski, Iwona Gabriel\*

Department of Pharmaceutical Technology and Biochemistry, Gdańsk University of Technology, 11/12 Narutowicza Str., 80-233 Gdańsk, Poland



## ARTICLE INFO

## Keywords:

Aminotransferases  
Gene knockout  
*Candida albicans*  
Amino acid metabolism  
Histidine  
Lysine

## ABSTRACT

Amino acids constitute the key sources of nitrogen for growth of *Candida albicans*. In order to survive inside the host in different and rapidly changing environments, this fungus must be able to adapt via its expression of genes for amino acid metabolism. We analysed the *ARO8*, *ARO9*, *YER152C*, and *BNA3* genes with regards to their role in the nutritional flexibility of *C. albicans*. *CaAro8p* is undoubtedly the most versatile enzyme among the aminotransferases investigated. It is involved in the catabolism of histidine, lysine, and aromatic amino acids as well as in L-Lys, Phe and Tyr biosynthesis. *CaAro9p* participates in the catabolism of aromatic amino acids and lysine at high concentrations of these compounds, with no biosynthetic role. Conversely, the *CaYer152Cp* catalytic potential for aromatic amino acid catabolism observed *in vitro* appears to be of little importance *in vivo*. Neither biosynthetic nor catabolic roles of *CaBan3p* were observed for any proteinogenic amino acid. Finally, none of the analysed aminotransferases was solely responsible for the catabolism of a single particular amino acid or its biosynthesis.

## 1. Introduction

In recent years, the number of immunocompromised patients suffering from various infectious diseases, such as those with cancer who have been subjected to extensive chemotherapy and organ transplantation, has increased substantially (Miceli et al., 2011). In particular, invasive fungal infections continue to be a major cause of morbidity and mortality in immunocompromised or severely ill patients (Groll and Lumb, 2012). The main etiological factor of opportunistic fungal infections (63–70% of cases) is *Candida albicans* (López-Martínez, 2010).

During the course of infection, *C. albicans* cells face many different and rapidly changing host environments. The ability to use as many nutrient sources as possible is essential for adaptation to each of these niches. Among different catabolic processes, the degradation of amino acids, necessary for their usage as a nitrogen source, has been demonstrated to be relevant for the survival of many microbial pathogens (Martinez and Ljungdahl, 2005). In the mammalian host infected by a pathogenic fungus, amino acids are present in a free form at low, millimolar concentrations in the plasma or urine (Tan and Gajra, 2006), or can be liberated from proteins and peptides upon the actions of fungal hydrolases. Their utilisation as a nitrogen source requires the presence of a respective aminotransferase in fungal pathogen cells (Fig. 1).

Human pathogenic fungi differ in their ability to use amino acids as a sole nitrogen source. *Histoplasma capsulatum* assimilates ammonium sulphate, aspartic acid and asparagine, although cannot grow with

methionine and gives only poor growth with cysteine, lysine, histidine, tryptophan and phenylalanine. *Sporotrichum schenckii* grows poorly or not at all with aspartic acid, lysine or histidine (Gilardi, 1965). *C. albicans* is able to use each of the 20 proteinogenic amino acids when grown in minimal medium with glucose as a carbon source (Brunke et al., 2014). Conversely, L-cysteine and L-lysine cannot support the growth of *Candida glabrata* or the baker's yeast, *Saccharomyces cerevisiae*, and only *C. glabrata* and *C. albicans* are able to grow on L-histidine. Specifically, the aromatic amino acid aminotransferase *CgAro8p* has been recently identified as an enzyme participating in histidine degradation in *C. glabrata* (Brunke et al., 2014).  $\alpha$ -Aminoadipate, an intermediate of the lysine biosynthetic pathway in fungi, when used as the sole nitrogen source, was found as growth inhibitory and toxic to *S. cerevisiae*, whereas the human pathogenic fungi, *C. albicans*, *C. neoformans* and *A. fumigatus*, are able to use both lysine and  $\alpha$ -aminoadipate as the sole nitrogen source. This unique metabolic property was proposed as a criterium for the identification of these pathogens (Ye et al., 1991).

In fungi, the catabolism of amino acids is complicated by the fact that some species can use them as the sole carbon source and some other as the sole nitrogen source. There are also fungal species that can use amino acids as both carbon and nitrogen source (Kinzel et al., 1983). Even within *Candida* species of the CUG clade, there is a great diversity in that ability. *C. albicans*, *C. tropicalis* and *C. lusitanae* utilize amino acids (in the form of Casamino Acids in YNB) effectively as the carbon source, whereas the growth of *C. dubliniensis* or *C. parapsilosis*

\* Corresponding author.

E-mail address: iwogabri@pg.gda.pl (I. Gabriel).




Fig. 1. The schematic overview of reactions performed by aromatic and aminoadipate transaminases for L-lysine and aromatic amino acids biosynthesis. The catabolic process is a reversal of the biosynthetic reaction.

under these conditions is much slower (Priest and Lorenz, 2015).

Enzymes catalysing particular steps of amino acid biosynthesis in fungi are considered as potential targets for antifungal therapy, especially those that are absent in mammalian hosts (Jastrzębowska and Gabriel, 2015; Kingsbury et al., 2004; Schöbel et al., 2010). Some inhibitors of such enzymes have already been identified and shown to exhibit antifungal activity (Yamaki et al., 1990; Aoki, 1996). Moderate antifungal activity was also demonstrated for compounds that are structural analogues of substrates or products of the enzymes involved in L-lysine biosynthesis pathway in *C. albicans* (Gabriel et al., 2013; Milewska et al., 2012).

In our efforts to identify an aminotransferase participating in L-Lys biosynthesis and degradation, we have focused our attention on four candidate genes: *CaARO8*, *CaARO9*, *CaYER152C*, and *BNA3*. In this work we present the results of our studies on the actual function of these genes and their products.

## 2. Materials and methods

### 2.1. Strains and growth conditions

*C. albicans* strains used in this investigation are listed in Table 1. The strains were routinely grown in YPG liquid medium (1% yeast extract, 1% peptone, 2% glucose) in a shaking incubator. For growth on solid media, 1.5% agar was added to the medium (YPG agar medium).

*Escherichia coli* TOP 10F' strain from Invitrogen was used in all cloning procedures. *E. coli* Rosetta (DE3) pLysS strain from Novagen was used for the overproduction of wild-type *CaAro8p* and *CaYer152Cp*, and the *E. coli* BL21 Star (DE3) strain from Invitrogen was used for the overproduction of *CaAro9p*. *E. coli* strains were cultured at 37 °C on Luria-Bertani (LB) solid medium (1.0% (w/v) NaCl, 1.0% (w/v) tryptone, 0.5% yeast extract, and 1.5% (w/v) agar) and LB liquid medium supplemented with 100  $\mu\text{g mL}^{-1}$  ampicillin and/or 34  $\mu\text{g mL}^{-1}$  chloramphenicol, when required.

### 2.2. Mutant strain construction

*C. albicans* SC5314 genomic DNA was isolated using a method described previously (Reuss et al., 2004). Isolation of plasmid DNA was carried out according to the protocol of the NucleoSpin Plasmid kit (Macherey-Nagel). DNA fragments were isolated from agarose gels following the standard procedure of the NucleoSpin Gel and PCR Clean-up kit (Macherey-Nagel). DNA digestion with the use of restriction enzymes was performed according to the enzyme supplier's instructions.

The pSFS5 plasmid (Sasse et al., 2011) was used for the construction of deletion cassettes. Restriction enzymes and DNA polymerase were purchased from New England Biolabs. The gene knock-out cassettes were prepared by amplification of upstream and downstream regions of each particular gene: *ARO8* (orf19.2098), *ARO9* (orf19.1237), *YER152C* (orf19.1180), and *BNA3*, present in two copies as *BNA31* (orf19.5809) and *BNA32* (orf19.1589.1), from *C. albicans* SC5314 genomic DNA followed by *SacI*-*SacII* and *XhoI*-*Apal* digestion. The primers used in this study are described in Table 2. The amplified regions were cloned on both sides of the *SAT1* flipper cassette in the pSFS5 plasmid, resulting in pARO8M1, pARO9M1, p1180M1, pBNA31M1, and pBNA32M1 plasmids. The knock-out cassettes were isolated by restriction digestion using *SacI* and *Apal* enzymes. The entire genes were deleted using the *SAT1* flipping strategy (Reuss et al., 2004). The reintroduced strains were prepared in accordance with the above mentioned procedure. The complementation cassettes were generated using the appropriate primers as described in Table 2. The complete ORF and flanking sequences were cloned as *SacI*-*SacII* fragment and substituted for the upstream sequence in appropriate plasmids, generating pARO8K1, pARO9K1, p1180K1, pBNA31K1 and pBNA32K1. The *SacI*-*Apal* regions were integrated into one of the alleles of homozygous mutants. Two independent mutants were constructed from their respective wild-type strain. The correct genomic integration of each construct and excision of the *SAT1* flipper cassette were confirmed by Southern hybridization with the upstream and downstream flanking sequences.

**Table 1**  
*C. albicans* strains used in this study.

| Strain(s)                         | Relevant characteristic or genotype                            | References           |
|-----------------------------------|--|----------------------|
| SC5314                            | Wild-type reference strain                                     | Gillum et al. (1984) |
| SCAR08M2A and -B                  | <i>ARO8/aro8Δ::FRT</i>   | This study           |
| SCΔ <i>aro8A</i> and -B           | <i>aro8Δ::FRT/aro8Δ::FTR</i>                                   | This study           |
| SCAR08K2A and -B                  | <i>aro8Δ::FTR/ARO8::FTR</i>                                    | This study           |
| SCAR09M2A and -B                  | <i>ARO9/aro9Δ::FRT</i>   | This study           |
| SCΔ <i>aro9A</i> and -B           | <i>aro9Δ::FRT/aro9Δ::FTR</i>                                   | This study           |
| SCAR09K2A and -B                  | <i>aro9Δ::FTR/ARO9::FTR</i>                                    | This study           |
| SCYER152CM2A and -B               | <i>YER125C/yer152CΔ::FRT</i>                                   | This study           |
| SCΔ <i>yer152CA</i> and -B        | <i>yer152CΔ::FRT/yer152CΔ::FTR</i>                             | This study           |
| SCYER152CK2A and -B               | <i>yer152CΔ::FTR/YER152C::FTR</i>                              | This study           |
| SCBNA31M2A and -B                 | <i>BNA31/bna31Δ::FRT</i>                                       | This study           |
| SCΔ <i>bna31A</i> and -B          | <i>bna31Δ::FRT/bna31Δ::FTR</i>                                 | This study           |
| SCBNA31K2B                        | <i>bna31Δ::FTR/BNA31::FTR</i>                                  | This study           |
| SCBNA32M2A and -B                 | <i>BNA32/bna32Δ::FRT</i>                                       | This study           |
| SC Δ <i>bna32A</i> and -B         | <i>bna32Δ::FRT/bna32Δ::FTR</i>                                 | This study           |
| SCΔ <i>aro8 ARO9M2A</i> and -B    | <i>aro8Δ, ARO9/aro9Δ::FRT</i>                                  | This study           |
| SCΔ <i>aro8 Δaro9A</i> and -B     | <i>aro8Δ, aro9Δ::FRT/aro9Δ::FTR</i>                            | This study           |
| SCΔ <i>aro8 ARO9K2A</i> and -B    | <i>aro8Δ, aro9Δ::FTR/ARO9::FTR</i>                             | This study           |
| SCAR08K2 Δ <i>aro9A</i> and -B    | <i>aro8Δ::FTR/ARO8::FTR, aro9Δ::FRT/aro9Δ::FTR</i>             | This study           |
| SCAR08K2 ARO9K2A and -B           | <i>aro8Δ::FTR/ARO8::FTR, aro8Δ, aro9Δ::FTR/ARO9::FTR</i>       | This study           |
| SCΔ <i>aro8 YER152CM2A</i> and -B | <i>aro8Δ, YER152C/yer152CΔ::FRT</i>                            | This study           |
| SCΔ <i>aro8 Δyer152CA</i> and -B  | <i>aro8Δ, yer152CΔ::FRT/yer152CΔ::FTR</i>                      | This study           |
| SCΔ <i>aro8 YER152CK2A</i> and -B | <i>aro8Δ, yer152CΔ::FTR/YER152C::FTR</i>                       | This study           |
| SCAR08K2 Δ <i>yer152CA</i>        | <i>aro8Δ::FTR/ARO8::FTR, yer152CΔ::FRT/yer152CΔ::FTR</i>       | This study           |
| SCAR08K2 YER152CK2A and -B        | <i>aro8Δ::FTR/ARO8::FTR, aro8Δ, yer152CΔ::FTR/YER152C::FTR</i> | This study           |
| SCΔ <i>aro8 BNA31M2A</i> and -B   | <i>aro8Δ, BNA31/bna31Δ::FRT</i>                                | This study           |
| SCΔ <i>aro8 Δbna31A</i> and -B    | <i>aro8Δ, bna31Δ::FRT/bna31Δ::FTR</i>                          | This study           |
| SCΔ <i>aro8 BNA31K2A</i> and -B   | <i>aro8Δ, bna31Δ::FTR/BNA31::FTR</i>                           | This study           |
| SCΔ <i>aro8 BNA31K2A</i> and -B   | <i>aro8Δ, BNA31::FTR/bna31Δ::FTR</i>                           | This study           |
| SCAR08K2 Δ <i>bna31A</i>          | <i>aro8Δ::FTR/ARO8::FTR, bna31Δ::FRT/bna31Δ::FTR</i>           | This study           |
| SCAR08K2 BNA31K2A                 | <i>aro8Δ::FTR/ARO8::FTR, aro8Δ, bna31Δ::FTR/BNA31::FTR</i>     | This study           |
| SCΔ <i>aro8 Δbna31 BNA32M2A</i>   | <i>aro8Δ, bna31Δ, BNA32/bna32Δ::FRT</i>                        | This study           |
| SCΔ <i>aro8 Δbna31 Δbna32A</i>    | <i>aro8Δ, bna31Δ, bna32Δ::FRT/bna32Δ::FTR</i>                  | This study           |

### 2.3. Cloning of *ARO8*, *ARO9*, and *YER152C* genes

Cloning of *ARO8* and *YER152C* was performed as previously described (Rząd and Gabriel, 2015) with the use of the pET Directional TOPO Expression Kit and the pET101/D-TOPO plasmid (Invitrogen). The same kit

was used for the construction of recombinant expression plasmids encoding *ARO9*. The fragment of the *ARO9* gene was amplified from the *C. albicans* SC5314 genomic DNA by PCR. The primers used in the amplification were: *ARO9.f* 5'-CACCATGTCGTACTCTACTCATTTAATTTCTAAG-3' and *ARO9.r* 5'-CTTAGGCTAGCTTTTAAACTCTAACCCATTAC-3'. Primers were

**Table 2**  
Primers used in the construction of knock-out cassettes.

| Name    | Sequence                          | Destination   |
|---------|-----------------------------------|---|
| ARO8.01 | tttGAGCTCaccacttaaaactgataagaaa   | Amplification of <i>ARO8</i> (orf19.2098) upstream region   |
| ARO8.02 | gtatCCGCGGtcattatgagcagtagaaa     |   |
| ARO8.03 | gtaaCTCGAGcttctgtgtgcaacaagt      | Amplification of <i>ARO8</i> (orf19.2098) downstream region   |
| ARO8.04 | caaaGGGCCCGctcgtaaagatttctctg     |   |
| ARO8.05 | tggacCCGCGGaaagattatctggaatg      | With ARO8.01 amplification of <i>ARO8</i> (orf19.1237) gene with upstream and downstream regions    |
| ARO9.01 | taaaGAGCTCacattatgatgatgatg       | Amplification of <i>ARO9</i> (orf19.1237) upstream region   |
| ARO9.02 | atatCCGCGGgtaacaaagttctctaatg     |   |
| ARO9.03 | tgggCTCGAGttttaaagactgacc         | Amplification of <i>ARO9</i> (orf19.1237) downstream region   |
| ARO9.07 | cttattgaattcagtaGGGCCcttacc       |   |
| ARO9.09 | gtaaCCGCGGgaaattgaatgacattgga     | Amplification of <i>ARO9</i> (orf19.1237) gene with upstream and downstream regions                 |
| ARO9.11 | aatgGAGCTCtttaaatgactctgtaag      |   |
| 1180.01 | aatgGAGCTCtttattatgagttgtttt      | Amplification of <i>YER152C</i> (orf19.1180) upstream region  |
| 1180.02 | tttaCCGCGGgagtaaaagttcatttc       |   |
| 1180.03 | acttCTCGAGtagttagtaaacagttttaa    | Amplification of <i>YER152C</i> (orf19.1180) downstream region                                      |
| 1180.04 | aataGGGCCcCaataatataatgataggg     |   |
| 1180.05 | caataCCGCGGcaaaaatataatgataggg    | With 1180.01 amplification of <i>YER152C</i> (orf19.1180) gene with upstream and downstream regions |
| 5809.01 | ccttGAGCTCtggaaaggtgtaattc        | Amplification of <i>BNA31</i> (orf19.5809) upstream region  |
| 5809.02 | ggaaaCCGCGGcttaacataaaaaggtgaa    |   |
| 5809.03 | taaaaCTCGAGcgtttgtagatgtagctg     | Amplification of <i>BNA31</i> (orf19.5809) downstream region  |
| 5809.04 | ctaGGGCCGgagtttttagtacaataacc     |   |
| 5809.06 | ctaCCGCGGgagtttttagtacaataacc     | With 5809.01 amplification of <i>BNA31</i> (orf19.5809) gene with upstream and downstream regions   |
| 1589.02 | aattCCGCGGgatgagttgagttgaaga      | With 5809.01 amplification of <i>BNA32</i> (orf19.1589.1) upstream region                           |
| 1589.03 | gttctcagtagtagctgCTCGAGcatat      | Amplification of <i>BNA32</i> (orf19.1589.1) downstream region                                      |
| 1589.04 | ttacatGGGCCcattgaaataaaccaactctg  |   |
| 1589.06 | ttacatCCGCGGgattgaaataaaccaactctg | With 1589.01 amplification of <i>BNA32</i> (orf19.1589.1) gene with upstream and downstream regions |

Bolded sequences are specific regions recognized by restriction enzymes: *SacI* (GAGCTC), *SacII* (CCGCGG), *XhoI* (CTCGAG), *ApaI* (GGGCC).

designed according to manufacturer instruction. The PCR products (1572 bp) were purified from an agarose gel and cloned directionally into the pET101/D-TOPO vector, thus yielding the recombinant expression plasmid pET101/D-TOPO + *ARO9* (7334 bp). The identity of the plasmid was confirmed by restriction analysis and DNA sequencing.

#### 2.4. Overexpression of *CaARO9* and purification of the gene product

Construction of the overexpression plasmid containing the *CaARO9* gene, conditions for *E. coli* transformation, overexpression of *CaARO9*, and isolation of Aro9p were the same as described previously for the *CaARO8* and *CaYER152* genes and their products (Rząd and Gabriel, 2015), except that *E. coli* BL21 Star (DE3) cells were used instead of *E. coli* Rosetta (DE3) pLysS as a host for the heterologous expression. Samples of purified protein were then held frozen ( $-20\text{ }^{\circ}\text{C}$ ) in 50 mM phosphate buffer pH 7.5, with pyridoxal 5'-phosphate (PLP, at slight molar excess with respect to the proteins) and 10% glycerol.

#### 2.5. Growth in liquid medium with different nitrogen sources

*C. albicans* cells were grown overnight at  $30\text{ }^{\circ}\text{C}$  in YPG medium. Cells from the overnight culture were washed twice with PBS and suspended to  $2 \times 10^5$  cells  $\text{mL}^{-1}$  in YNB medium w/o amino acids with 2% glucose, 100 mM phosphate buffer pH 5.8. Aliquots of 100  $\mu\text{L}$  were used to inoculate the microtiter wells containing 100  $\mu\text{L}$  of YNB medium w/o amino acids, containing a particular amino acid, aminoadypic acid, or ammonium sulphate to a final concentration of 10 mM, 5 mM, and 0.25%, respectively. If necessary, the aminotransferase inhibitor (aminoxy)acetate (AOA) was added (10 mM). The cell suspensions were cultivated for 48 h at  $30\text{ }^{\circ}\text{C}$  and the cell density was measured at time intervals spectrophotometrically ( $\lambda = 531\text{ nm}$ ) with a microplate reader (Victor<sup>3</sup> Perkin Elmer). Additional experiments were performed for L-Lys, L-His and L-Tyr at different amino acids concentration. Serial 2-fold dilution of that amino acids were made in 100  $\mu\text{L}$  YNB medium w/o nitrogen source with 2% glucose, 100 mM phosphate buffer pH 5.8 in 96-well plate. Aliquots of 100  $\mu\text{L}$  containing  $2 \times 10^5$  *C. albicans* cells  $\text{mL}^{-1}$  were used to inoculate microtiter wells containing 100  $\mu\text{L}$  of serial dilutions of selected amino acid.

#### 2.6. Determination of aminotransferase activity

The aminotransferase activity of isolated enzymatic proteins was determined as previously described (Rząd and Gabriel, 2015). In all assays, 2.5  $\mu\text{g}$  of the purified enzyme was used. Briefly, in assays of aromatic aminotransferase activity, the product formation was measured spectrophotometrically at  $\lambda = 320\text{ nm}$  for phenylpyruvate (PhP),  $\epsilon = 17\,700\text{ dm}^3\text{ cm}^{-1}\text{ mol}^{-1}$ ;  $\lambda = 331\text{ nm}$  for 3-indolepyruvate,  $\epsilon = 19\,900\text{ dm}^3\text{ cm}^{-1}\text{ mol}^{-1}$ ; and  $\lambda = 338\text{ nm}$  for 4-hydroxyphenylpyruvate (4-hydroxyPhP),  $\epsilon = 9\,300\text{ dm}^3\text{ cm}^{-1}\text{ mol}^{-1}$ . The rate of Phe or Tyr formation was estimated by quantification of the decrease in the PhP or 4-hydroxyPhP derived absorbance, respectively.

$\alpha$ -Amino adipate aminotransferase activity was followed with the use of L-glutamate and 2-oxoadipate (2-OA) as substrates. Estimation of the quantity of 2-oxoglutarate (2-OG) formed was performed using an L-glutamic dehydrogenase assay in which formation of  $\text{NAD}^+$  was measured at  $\lambda = 340\text{ nm}$ ;  $\epsilon = 6\,200\text{ dm}^3\text{ cm}^{-1}\text{ mol}^{-1}$ .

Catabolic reactions of  $\alpha$ -amino adipate or L-histidine degradation were analysed using the same method, in reaction mixtures containing  $\alpha$ -amino adipate/2-oxoglutarate or L-histidine/2-oxoglutarate as substrates, respectively. Formation of L-glutamate was measured spectrophotometrically with the use of the L-Glutamate Assay Kit (Sigma), according to the manufacturer's instructions.

The effect of (aminoxy)acetate (AOA) on aminotransferase activity was determined by measuring the enzyme activity in standard assay mixtures containing various concentrations of Phe, 2-OG (10 mM), and different concentrations of AOA. The level of PhP formation was estimated spectrophotometrically as mentioned above.

### 3. Results

Our previous studies showed that the *ARO8* gene of *C. albicans* encodes a transaminase, exhibiting aromatic and amino adipate aminotransferase activity (Rząd and Gabriel, 2015). It has also been known that in *S. cerevisiae* aromatic aminotransferase exists in two isoforms encoded by the *ARO8* and *ARO9* genes (Iraqi et al., 1998). Apart from these two genes, the phylogenetic analysis of aromatic transaminases in *S. cerevisiae* genome performed by Hébert et al. (2011) resulted in identification of the *YER152C* ORF as a gene encoding putative aromatic amino acid aminotransferase. On the other hand, the putative amino adipate aminotransferase activity has also been suggested for the products of *ARO8*, *BNA3* and *YER152C* from *S. cerevisiae* by King et al. (2009).

A BLAST search, using the nucleotide sequence of the respective ORFs from the *Saccharomyces* Genome Database ([www.yeastgenome.org](http://www.yeastgenome.org)) performed on the *Candida* Genome Database ([www.candidagenome.org](http://www.candidagenome.org)) revealed orf19.1237, (named *ARO9*), orf19.1180 (*YER152C*) and orf19.5809 or orf19.1589.1, both associated with *BNA3* (being present in two copies, *BNA31* and *BNA32*), highly homologous to their *S. cerevisiae* counterparts (55%, 64% and 69% of similarity, respectively). The nucleotide sequences of *CaARO9*, *CaYER152C* and *CaBNA3* exhibit 31%, 24%, 23% identity and 48%, 44%, 37% similarity to *CaARO8*, respectively.

#### 3.1. Growth in minimal media with different nitrogen sources

To investigate the functions of the aminotransferases encoded by *ARO8*, *ARO9*, *YER152C*, and *BNA3* in nitrogen utilisation by *C. albicans*, deletion mutants of the SC 5314 strain lacking either one, two, or three of these genes were constructed and tested for their ability in using ammonium sulphate, proteinogenic amino acids, and intermediates of lysine biosynthesis and breakdown as the sole nitrogen source. The results data presented in Fig. 2 indicate that all mutant strains grew equally as well as the wild-type strain in minimal medium with ammonium sulphate as the sole nitrogen source. This result suggested that all mutants were able to synthesise all amino acids necessary for protein biosynthesis; i.e., none of the already identified or putative aminotransferases encoded by *ARO8*, *ARO9*, *YER152C*, or *BNA3* is essential for biosynthesis of any of the proteinogenic amino acids. The majority of the mutants were also able to use each of the 19 proteinogenic amino acids as the sole nitrogen source to support their growth. However, differences in histidine, lysine, and tyrosine utilisation were observed between the mutants lacking the *ARO8* gene and the wild-type SC 5314 cells. The difference can be better seen in Fig. 3, in which the growth kinetics are presented. The growth rates of the all *ARO8*-deficient mutants in minimal media containing L-His or L-Lys as the sole nitrogen source were much lower than that of the wild-type strain. The respective difference was lower for L-Tyr utilisation, but still evident. Moreover, a single gene disruption of *ARO9*, *YER152C*, or *BNA3* resulted in no effect on *C. albicans* growth with L-His, L-Lys, or L-Tyr, thus indicating that none of these genes is essential for the catabolism of these amino acids.

Notably, all the mutants tested, including the *ARO8*-deficient mutants, demonstrated similar ability as that of the wild-type strain to use L- $\alpha$ -amino adipate (AA) as the sole nitrogen source, whereas deletion of *ARO8* apparently affected growth on L-Lys (Fig. 4). AA is an intermediate in the L-lysine catabolic pathway, specifically functioning as a direct substrate for a putative aminotransferase (Fig. 1).

The growth ability of *ARO8*-deficient mutants could be to a large extent restored by supplementation of the minimal medium with L-Lys or L-Tyr in an amino acid concentration-dependent manner ( $62.3 \pm 4.2\%$  and  $74.0 \pm 4.9\%$  restoration, in comparison with the wild-type cells at 4 mM L-Lys or L-Tyr, respectively), as shown in Fig. 5. Conversely, supplementation with L-His apparently did not restore growth of these mutants (only  $19.9 \pm 3.7\%$  of that of SC5314 at 4 mM L-His). Moreover, the growth ability was not related to the concentration of the nitrogen source.




Fig. 2. Growth of *C. albicans* wild-type and deletion mutant strains in YNB media containing ammonium sulphate (AS) or particular amino acids as the sole nitrogen source. Optical density of cell suspensions and liquid media (negative control) was measured at 531 nm after incubation for 48 h at 30 °C. All data represent the means ± SD.




Fig. 3. Growth kinetics of *C. albicans* wild-type and deletion mutants in the YNB media containing either L-Lys, L-His, or L-Tyr as the sole nitrogen source (10 mM) or ammonium sulphate (AS). Cell density was measured at time intervals spectrophotometrically ( $\lambda = 531$  nm). Optical density of liquid medium (YNB) serves as a negative control.




Fig. 4. Growth of *C. albicans* wild-type and deletion mutant strains in YNB media containing L-Lys or α-aminoacidate as the sole nitrogen source. Optical density of cell suspensions and liquid media (negative control) was measured at 531 nm after incubation for 48 h at 30 °C. All data represent the means ± SD.




Fig. 5. Growth of *C. albicans* wild-type and deletion mutants on various concentrations of L-His, L-Lys, or L-Tyr supplementing the YNB media as the sole nitrogen source. Optical density of cell suspensions and liquid media (negative control) was measured at 531 nm after incubation for 48 h at 30 °C.




Fig. 6. Effect of (Aminoxy)acetate (AOA) on *C. albicans* wild-type and mutant growth in YNB with ammonium sulphate (AS) medium and YPD. Dashed lines, no AOA present; solid lines, 10 mM AOA. Optical density at 531 nm of the appropriate liquid medium served as a control.

### 3.2. Effect of aminotransferase inhibition on *C. albicans* growth

(Aminoxy)acetate (AOA) is a broad-spectrum inhibitor of pyridoxal phosphate-dependent enzymatic reactions, such as transaminations, which was also shown to inhibit some fungal transaminases (Brunke et al., 2014; Preuss et al., 2013; Wallach, 1961). AOA has also been reported to affect the growth of *C. glabrata* and *C. albicans* cells (Brunke et al., 2014). In our hands, presence of 10 mM AOA in cultures of *C. albicans* wild-type and mutant cells in minimal medium with AS as the sole nitrogen source resulted in growth retardation; the magnitude of this effect was roughly the same, regardless of strain type (Fig. 6). Growth retardation by AOA was not observed in the YPD complex medium, thus demonstrating that AOA is not a general toxin and that its growth inhibitory effect could be reversed by peptides and amino acids present in YPD. AOA also strongly inhibited the growth of *C. albicans* in minimal medium with selected amino acids as the sole nitrogen source in an amino acid concentration-dependent manner (Fig. 7). AOA strongly inhibited the growth of *C. albicans* SC5314 and mutant strains on 5 mM L-His as the nitrogen source. The same results were obtained by Brunke et al. (2014) for *C. albicans* SC5314 and *C. glabrata* ATCC2001. In contrast, when 10 mM L-His was used, after a long lag phase growth mostly resumed for *C. albicans* SC5314, SCΔ*yer152C*, and SCΔ*aro9*. In this instance, the main role in L-His degradation could be acquired by other remaining transaminases, such as CaAro8p or CaBna3p. Despite the fact that AOA inhibits CaAro8p and CaAro9p at micromolar concentrations *in vitro* with  $IC_{50} 96 \pm 3.5$  M,  $K_i$  L-Phe  $8.7 \pm 1.4$  M and  $IC_{50} 11.8 \pm 3$  M,  $K_i$  L-Phe  $3 \pm 0.3$  M, respectively, these or other unknown aminotransferases might not be completely inhibited *in vivo*.

The growth of strains deficient in the *ARO8* gene, including SCΔ*aro8*, SCΔ*aro8Δaro9*, SCΔ*aro8Δyer152C*, and SCΔ*aro8Δbna31* remained reduced. For the *C. albicans* SCΔ*aro8Δbna31Δbna32* strain, no significant growth with any analysed concentration of L-His was detected even after 48 h in the presence of AOA. Our results, in combination with the data from the growth phenotypes of deletion mutants in minimal media with ammonium sulphate, demonstrate that L-histidine biosynthesis is not dependent on the aminotransferase activity of Aro8p, Aro9p, Yer152Cp, or Bna3p in *C. albicans*, although the presence of CaAro8p is crucial for degradation of this amino acid.

*C. albicans* SC5314 and mutant strain growth inability was detected with 10 mM L-Lys even after 48 h in the presence of 10 mM AOA (Fig. 7). These results demonstrate that lysine utilisation is strongly dependent on the aminotransferase activity. It is probable that the

growth phenotype of the wild-type SC5314 and of SCΔ*yer152C* with L-Lys as the sole nitrogen source and AOA is the same as that for mutant strains (SCΔ*aro8*, SCΔ*aro9*, SCΔ*aro8Δaro9*, SCΔ*aro8Δyer152C*, SCΔ*aro8Δbna31*, and SCΔ*aro8Δbna31Δbna32*), which are deficient in CaAro8p and/or CaAro9p aminotransferase activity, partially owing to the inhibition of both enzymes by AOA *in vivo* and the inhibition of unknown transaminases involved in L-Lys degradation. Overall, these results, in combination with the data obtained from the growth phenotypes of deletion mutants in minimal media with ammonium sulphate, demonstrate that lysine biosynthesis is not dependent on the aminotransferase activity of CaAro8p and other analysed aminotransferases. However, degradation of this amino acid is strongly dependent on CaAro8p aminotransferase activity, with an important role of transaminases not examined in this study also being probable.

L-Phe and L-Trp utilisation is also dependent on aminotransferase activity. Addition of the inhibitor AOA to *C. albicans* cultures strongly inhibited the growth of *C. albicans* SC5314 and mutant strains on 5 mM L-Phe and L-Trp as a nitrogen source. At higher concentrations of analysed amino acids, growth mostly resumed for all strains, regardless of the presence or absence of the tested genes. These results demonstrate that in *C. albicans*, L-Phe and L-Trp utilisation is not as strongly dependent on the aminotransferase activity as is L-Lys and L-His degradation. In our hands, in contrast to results obtained by Brunke et al. (2014), significant growth of *C. albicans* SC5314 in the presence of 5 or 10 mM L-Tyr and 10 mM AOA was detected. Notably, according to our results, L-Tyr utilisation is not performed mainly by aromatic aminotransferases as assumed by other authors.

### 3.3. Catalytic properties of CaAro8p, CaAro9p, and CaYer152Cp

The CaARO9 gene was heterologously expressed in *E. coli* and its product, CaAro9p, was isolated and purified in several steps including ammonium sulphate precipitation, desalting, and ion exchange chromatography, under conditions previously reported for CaAro8p and CaYer152Cp (Rzaq and Gabriel, 2015).

The CaAro8p, CaAro9p, and CaYer152Cp proteins were tested for their catalytic properties. The kinetic parameters of reactions catalysed by CaAro8p and CaAro9p are summarized in Table 3. CaYer152Cp catalysed the transamination of L-Phe, L-Tyr and L-Trp with 2-oxoglutarate as an amino group acceptor (kinetic parameters:  $K_{mPhe} = 0.24 \pm 0.024$  mM,  $V_{maxPhe} = 184 \pm 4$  nmol min<sup>-1</sup> mg<sup>-1</sup>;  $K_{mTyr} = 0.46 \pm 0.122$  mM,  $V_{maxTyr} = 164 \pm 15$  nmol min<sup>-1</sup> mg<sup>-1</sup>; and  $K_{mTrp} = 1.07 \pm 0.182$  mM,  $V_{maxTrp} = 383 \pm 16$  nmol min<sup>-1</sup> mg<sup>-1</sup>, Rzaq and Gabriel, 2015). No




Fig. 7. Effect of (Aminoxy)acetate (AOA) on *C. albicans* wild-type and mutant growth in YNB media containing particular amino acids as the sole nitrogen source. Dashed lines, no AOA present; solid lines, 10 mM AOA. Optical density at 531 nm of the appropriate liquid medium served as a control.

activity of this enzyme was detected with L- $\alpha$ -amino adipate and L-His as substrates for 2-oxoglutarate amination, nor were any transamination products formed with phenylpyruvate or 4-hydroxyphenylpyruvate as

amino acceptors from L-Glu.

Analysis of the kinetic parameters of reactions catalysed by the three homologous enzymes, determined under *in vitro* conditions,



**Table 3**  
Kinetic parameters of catabolic and anabolic reactions catalysed by *CaAro8p* and *CaAro9p*.

| Aro8p   |               |  | Aro9p        |             |  |
|---|---------------|--|--------------|-------------|--|
| Substrate   | $K_m$ [mM]    | $V_{max}$ [nmol min <sup>-1</sup> mg <sup>-1</sup> ] | Substrate    | $K_m$ [mM]  | $V_{max}$ [nmol min <sup>-1</sup> mg <sup>-1</sup> ] |
| <i>Catabolic activity; amino group acceptor: 2-oxoglutarate</i> |               |  |              |             |  |
| L-Phe   | 0.053 ± 0.003 | 1200 ± 27  | L-Phe        | 4.76 ± 1.17 | 7457 ± 457   |
| L-Tyr   | 0.09 ± 0.008  | 1528 ± 34  | L-Tyr        | 2.80 ± 0.25 | 6922 ± 240   |
| L-Trp   | 1.68 ± 0.28   | 6634 ± 357   | L-Trp        | 1.80 ± 0.18 | 4988 ± 168   |
| L-AA <sup>a</sup>   | 0.016 ± 0.008 | 779 ± 87   | L-AA         | 4.28 ± 1    | 6466 ± 511   |
| L-His   | 11.5 ± 2.42   | 6520 ± 567   | L-His        | ND          | ND   |
| <i>Anabolic activity; amino group donor: L-Glu</i>              |               |  |              |             |  |
| PhP   | 0.017 ± 0.004 | 696 ± 68   | PhP          | ND          | ND   |
| 4-hydroxyPhP  | 0.026 ± 0.004 | 870 ± 50   | 4-hydroxyPhP | ND          | ND   |
| 2-oxoadipate  | 0.1 ± 0.033   | 9954 ± 2032  | 2-oxoadipate | 0.13 ± 0.04 | 7743 ± 577   |

<sup>a</sup>  $K_m$  values of 2-oxoglutarate in this reaction were: 0.27 ± 0.07 mM for Aro8p and 0.41 ± 0.036 mM for Aro9p, respectively; ND – not determined, very low activity.

indicated that under *in vivo* conditions, *CaAro8p* is probably mainly involved in L-Phe, L-Tyr, and L-Lys catabolism when concentration of these amino acids in the cytosol is low, such as at the micromolar level, whereas at higher, millimolar concentrations, *CaAro9p* takes over this duty. Both enzymes exhibit similar affinity for L-Trp as an amino donor in the transamination of 2-oxoglutarate and show comparable catalytic efficiency in this reaction. The most intriguing finding is a low affinity (high  $K_m$  value) of *CaAro8p* for L-His as a substrate in 2-oxoglutarate transamination, assuming that *C. albicans* mutants lacking the *ARO8* gene displayed the most noticeable growth defects in minimal medium with L-His as the sole nitrogen source. Nevertheless, the high catalytic efficiency of *CaAro8p* in reaction with L-His, reflected by its high  $V_{max}$ , may compensate for the low affinity for this substrate. It is noteworthy therefore, that a similar  $K_m$  value was found for Aro8p from *C. glabrata*, namely 8.80 ± 1.41 mM, although the  $V_{max}$  was in that case much lower; i.e., 274.9 ± 19.8 nmol min<sup>-1</sup>. However, the *C. glabrata*  $\Delta$ aro8 strain was not able to grow in minimal medium with L-His as the sole nitrogen source, which identified *CgAro8p* as the principal transaminase involved in histidine catabolism (Brunke et al., 2014). Our results obtained for the *C. albicans* version of Aro8p (high  $K_m$  and  $V_{max}$  and growth impairment but not complete inability of the  $\Delta$ aro8 mutant) suggest that activity of this enzyme is important for L-His degradation, especially at high L-His concentration, although another, as-yet unidentified transaminase is probably operative at lower concentrations of this amino acid. The kinetic parameters determined for transaminations with L-Glu as an amino donor clearly suggest that *CaAro8p* but not *CaAro9p* can participate in the biosynthesis of aromatic amino acids (confirmed for L-Phe and L-Tyr and probably also L-Trp) and that both enzymes may be involved in L-Lys biosynthesis. In comparison, the undisturbed growth of  $\Delta$ aro8,  $\Delta$ aro9, and  $\Delta$ aro8 $\Delta$ aro9 mutants in minimal medium with AS as the sole nitrogen source indicates that under *in vivo* conditions neither *CaAro8p* nor *CaAro9p* serves as the only transaminase participating in the biosynthesis of lysine and aromatic amino acids.

Despite the high degree of *CaYer152Cp* sequence similarity to *CaAro8p* (44%), the *YER152C* gene product exhibited a narrower substrate spectrum (aromatic amino acids only) and relatively low catalytic efficiency (low  $V_{max}$  values) in catabolic transaminations, with apparently no anabolic activity. These properties are in contrast to those of the *S. cerevisiae* *YER152C* gene product, which was identified as an L- $\alpha$ -aminoadipate aminotransferase (King et al., 2009). Taking into account that the  $K_m$  values of *CaYer152Cp* for L-Phe and L-Tyr transamination with 2-oxoglutarate as an amino acceptor are much higher than those of *CaAro8p* and *CaAro9p* and the  $V_{max}$  values are much lower, it appears that the role of this enzyme in the catabolism of aromatic amino acids is of secondary importance. This assumption is supported by the fact that a single gene disruption of *YER152C* did not affect any deficiency of *C. albicans* growth in minimal media with L-Phe, L-Tyr or L-Trp as the only

nitrogen source (Fig. 2). However, the possibility that this protein exhibits any other catalytic activity, not detected in our studies, cannot be excluded.

Although cloning of the *BNA3* gene was performed and the construction of recombinant expression plasmids was completed successfully, we were not able to obtain overproduction of the *CaBNA3p* in the *E. coli* cells used as a host for heterologous expression.

#### 4. Discussion

Nutrient assimilation is a central and fundamental prerequisite for the growth and survival of all living organisms. Among the most important nutrients, nitrogen and carbon are required for almost all biosynthetic processes and must be assimilated in large quantities. The major fungal pathogens display different lifestyles, and, consequently, their metabolic flexibility has been shaped by different evolutionary pressures. *C. albicans* in particular faced especially diverse living conditions. This facultative pathogen can exist primarily as a commensal of the oral cavity as well as the gastrointestinal and urogenital tracts but can also persist within other extracellular (blood, tissues) and intracellular microenvironments, such as inside damaged epithelial and endothelial cells (Brock, 2009).

The unique unlimited ability of *C. albicans* to use different amino acids as the sole nitrogen source may, at least in part, be due to the versatility of its transaminases. In the hemiascomycete *C. glabrata*, histidine degradation was shown to be associated with an aromatic amino acid aminotransferase, Aro8p (Brunke et al., 2014). In the present study, we identified three genes in *C. albicans* of moderate homology to *CaARO8*, encoding putative aminotransferases with possible aminoadipate/aromatic aminotransferase activity, namely *ARO9*, *YER152C*, and *BNA3*. Our results indicated that *C. albicans* differs significantly from *S. cerevisiae* and *C. glabrata* in its ability to survive the lack of Aro8p or Aro9p activity. In contrast to *S. cerevisiae* (Iraqi et al., 1998; Urrestarazu et al., 1998), *C. albicans*  $\Delta$ aro8 $\Delta$ aro9 double mutants appeared not to be auxotrophic for aromatic amino acids. Moreover, the mutant strain behaviour was completely unique in respect to its growth ability when phenylalanine, tyrosine, or tryptophan comprised the only nitrogen source. Neither the  $\Delta$ aro8 nor the  $\Delta$ aro9 mutant exhibited growth impairment, whereas only minor changes were observed for growth of the  $\Delta$ aro8 mutant on tyrosine. In this regard, *C. albicans* differs considerably from *C. glabrata*, for which a severe growth defect of the  $\Delta$ aro8 mutant in tryptophan-containing medium and significantly reduced growth in Phe and Tyr-containing media were observed. *C. glabrata*  $\Delta$ aro9 mutant was only slightly reduced in growth. The  $\Delta$ aro8 mutant also showed a severe growth defect with histidine as the sole nitrogen source (Brunke et al., 2010, 2014). Notably, this is the only amino acid that *C. glabrata* can use for growth, but *S. cerevisiae* cannot (Brunke et al., 2010). Only cysteine and lysine cannot support the

growth of both *C. glabrata* and *S. cerevisiae* (Brunke et al., 2014). The capability of using L-Lys as nitrogen source is unique exclusively for *C. albicans* and according to our results, it appears that the CaAro8p aminotransferase contributes significantly to the growth ability of *C. albicans* on lysine as the sole nitrogen source.

The *S. cerevisiae*  $\Delta$ aro8 mutant was not auxotrophic but totally failed to grow when the nitrogen source was tryptophan, tyrosine, phenylalanine, or a mixture of these three, whereas the  $\Delta$ aro8 $\Delta$ aro9 double mutants were auxotrophic for both phenylalanine and tyrosine and totally failed to grow when tryptophan or methionine were the nitrogen source (Iraqi et al., 1998; Urrestarazu et al., 1998). These authors postulated that Aro8p from *S. cerevisiae* plays an essentially biosynthetic role with regards to phenylalanine and tyrosine formation. In comparison, they consider that Aro9p is mainly involved in tryptophan degradation and is an inducible catabolic enzyme that nevertheless ensures the formation of Phe and Tyr in *S. cerevisiae*  $\Delta$ aro8 mutants.

In baker's yeast, tryptophan is metabolised by two distinct routes: via the Ehrlich pathway to indole-3-ethanol (common for methionine, aromatic and branched-chain amino acids, in which the first step is catalysed by Aro8p and Aro9p) (Hazelwood et al., 2008) or via the kynurenine pathway to NAD<sup>+</sup> (Panozzo et al., 2002). It was postulated that in *S. cerevisiae*, the aromatic aminotransferase Aro8p participates in L-Trp biosynthesis and Aro9p is involved principally in its degradation. Presumably, the latter constitutes an inducible catabolic enzyme that is normally absent under anaerobic growth conditions (Urrestarazu et al., 1998; Iraqi et al., 1998).

The results of our studies indicate that in *C. albicans*, tryptophan degradation is not mainly performed by the aromatic amino acid aminotransferases CaAro8p or CaAro9p in the first pathway, as the ARO8 and ARO9 deficient mutants did not show any growth defect in minimal medium containing Trp as the sole nitrogen source. Moreover, although the *in vitro* studies of catalytic properties of CaAro8p and CaAro9p revealed that both enzymes use L-Trp as an amino donor in the transamination of 2-oxoglutarate, the kinetic parameters, especially relatively high K<sub>m</sub> values for L-Trp, suggest that in *C. albicans* there must be another aminotransferase that utilises L-Trp as a substrate, operating at lower concentrations of this amino acid. Therefore, we decided to investigate the importance of the second pathway with regards to tryptophan degradation.

In *S. cerevisiae*, the kynurenine pathway is a recognized route for *de novo* NAD<sup>+</sup> biosynthesis; in this pathway, the BNA1 to BNA6 genes encoding enzymes catalysing the conversion of tryptophan to nicotinic acid mononucleotide have been identified. However, whereas BNA1, BNA2, BNA4, BNA5, and BNA6 have orthologues in the genomes of *Debaryomyces hansenii*, *C. albicans*, and *Yarrowia lipolytica*, these all are absent in *C. glabrata*, *Kluyveromyces lactis*, and *Schizosaccharomyces pombe*. Conversely, BNA3 is present in every genome (Li and Bao, 2007). Although BNA3 was initially presumed to encode N-formylkynurenine formamidase (FKF), catalysing the second step of the kynurenine pathway (Panozzo et al., 2002), subsequent studies demonstrated that Bna3p instead most likely functions as the kynurenine aminotransferase, which converts kynurenine to kynurenic acid (Wogulis et al., 2008). Notably, we have found that in the *C. albicans* genome, the BNA3 gene is present in two copies, BNA31 and BNA32, stressing the presumed importance of a putative CaBna3p aminotransferase. However, knock-out of both BNA31 and BNA32 in *C. albicans* resulted in no growth impairment with L-Trp as a sole nitrogen source. Moreover, a similar effect was obtained when CaARO8, CaBNA31, and CaBNA32 genes were simultaneously disrupted.

Finally, it appears that in *C. albicans*, Aro8p and Aro9p likely participate in tryptophan degradation at millimolar concentrations of this amino acid but another, as-yet unidentified aminotransferase is operative when the concentration is lower. This unidentified aminotransferase may also be involved in the biosynthesis of L-Trp.

It is known that *S. cerevisiae* is unable to use histidine as the sole nitrogen source (Large, 1986), while in *C. glabrata*, histidine

degradation was shown to be exclusively associated with CgAro8p (Brunke et al., 2014). Conversely, Our results indicate that in *C. albicans*, CaAro8p serves as a principal but not the only enzyme involved in histidine catabolism, as the growth of the  $\Delta$ aro8 mutant was severely hampered but not completely abolished. The products of three other genes homologous to ARO8, namely ARO9, YER152C, and BNA3 are apparently not involved in histidine catabolism. Because it is well known that hemiascomycetes, including *C. albicans*, do not possess any gene encoding histidine ammonia lyase (histidinase) (Brunke et al., 2014), which is a principal enzyme catabolising histidine in bacteria and in most Eukaryota, we hypothesise that in *C. albicans* some other aminotransferase, not investigated in the present study, may be able to catalyse the amino transfer from L-His to an  $\alpha$ -ketoacid.

Our results indicate that the role of aminotransferases in lysine metabolism in *C. albicans* is to some extent unique in comparison with other hemiascomycetes. As in the other fungi, L-Lys is synthesized in *C. albicans* via the seven-step AA pathway (Xu et al., 2006). The fourth step of this pathway is catalysed by the PLP-dependent  $\alpha$ -aminoacidipate aminotransferase. Although Aro8p from *S. cerevisiae* was considered to be a putative  $\alpha$ -aminoacidipate aminotransferase on the basis of structural studies (Bulfer et al., 2013) and the kinetic parameters of CaAro8p-catalysed transamination with AA as an amino donor confirmed the biosynthetic potential of this enzyme, our  $\Delta$ aro8 mutant was not auxotrophic for lysine. The same was true for the  $\Delta$ aro9 mutant and for the  $\Delta$ aro8 $\Delta$ aro9 double mutant. In comparison, in our previous studies we showed that disruption of the CaLYS21 and CaLYS22 genes encoding isoforms of homocitrate synthase (Gabriel and Milewski, 2016) or the LYS4 gene coding for homoacnitase; i.e., enzymes catalysing two initial steps of the AA pathway, resulted in auxotrophy of *C. albicans* cells for lysine (Kur et al., 2010; Gabriel et al., 2014). Further analysis indicated that the  $\Delta$ lys21 $\Delta$ lys22 and  $\Delta$ lys4 mutant cells were able to utilize AA, an intermediate of the lysine biosynthesis pathway, to rescue this auxotrophy. In the present study, we found that AA as the sole nitrogen source supported growth of the ARO8 deficient mutants and that no statistically significant differences in growth kinetics were observed. These results differ significantly from that obtained for *C. albicans*  $\Delta$ aro8 mutant cells in the presence of L-Lys as the sole nitrogen source. Therefore, our results indicate that CaAro8p constitutes the principal but not the only transaminase involved in lysine biosynthesis in *C. albicans*.

The  $\Delta$ aro8 mutant demonstrated general growth impairment in minimal media with lysine as the sole nitrogen source, although this effect was weaker than that observed in L-His-containing minimal medium. The kinetic parameters of CaAro8p, especially the high affinity to AA reflected by the low K<sub>m</sub> value, confirm its potential principal role in lysine degradation. In contrast, the much higher K<sub>m</sub> for AA of CaAro9p is associated with the high V<sub>max</sub> for this substrate, although the  $\Delta$ aro9 mutant did not show any growth defect in lysine-supplemented minimal medium. The most plausible interpretation of these data supposes a principal role of CaAro8p in lysine degradation and auxiliary participation of CaAro9p in this process, when *C. albicans* cells are overloaded with lysine. As previously reported, *C. glabrata* and *S. cerevisiae* were not able to grow in minimal medium with L-Lys as the sole nitrogen source (Brunke et al., 2014), despite the presence of ARO8 genes in the genomes of both species. This indicates the unique role of Aro8p in *C. albicans*.

In fungal microorganisms possessing all enzymes catalysing the complete Ehrlich pathway, for example in *S. cerevisiae*, L-methionine comprises one of the substrates for Aro8p/Aro9p catalysing the first committed step of this pathway. Another possibility for methionine utilisation is via the methionine salvage pathway, in which the  $\alpha$ -amino group of methionine does not serve as a nitrogen source for the cells but is incorporated into polyamines, spermine, and spermidine. In this cyclic pathway, methionine is finally reconstructed by the transamination of 4-methylthio-2-oxobutyrate catalysed by Aro8p, Aro9p, and the branched-chain amino acid aminotransferases Bat1p and Bat2p

## CONCENTRATION OF AMINO ACID IN THE CYTOSOL




Fig. 8. A proposed model of *CaAro8p*, *CaAro9p* and *CaYer152Cp* role in *C. albicans* L-Lys, L-His and aromatic amino acids metabolism.

(Pirkov et al., 2008). This pathway clearly does not have any nutritional role. Nevertheless, if methionine is catabolised by Aro8p or Aro9p, the *ARO8* and *ARO9* deficient mutants should exhibit substantial growth impairment in minimal medium containing methionine as the sole nitrogen source. Accordingly, growth of the *C. glabrata* mutant lacking Aro8p activity was significantly reduced under this condition (Brunke et al., 2010). In our hands, the growth phenotype of the *C. albicans*  $\Delta$ aro8 mutant was comparable to that of the wild-type SC5314 cells. On the basis of previously published results, among five enzymes proposed to be important for L-Met catabolism in the group of hemiascomycetes, namely Bat1p, Bat2p, Aro8p, Aro9p, and Yer152Cp, only two are present in all species: Aro8p and Yer152Cp (Hébert et al., 2011). In our studies, the growth profile of the *C. albicans* SC $\Delta$ aro8 $\Delta$ yer152C mutant strain in minimal media with ammonium sulphate or L-Met as a nitrogen source was comparable to that of the wild-type SC5314. Therefore, it may be confidently concluded that none of the four aminotransferases investigated in the present study is involved in methionine catabolism in *C. albicans*.

In summary, our results indicate that *CaAro8p* is undoubtedly the most versatile enzyme among the aminotransferases investigated. It is involved in the catabolism of histidine, lysine, and aromatic amino

acids as well as in the biosynthesis of L-Lys, Phe and Tyr. *CaAro9p* appears to be an auxiliary enzyme, participating in the catabolism of aromatic amino acids and lysine at high concentrations of these compounds, with no biosynthetic role. The catalytic potential of *CaYer152Cp* for the catabolism of aromatic amino acids deduced from the results of *in vitro* studies appears to be of little importance *in vivo*. The principal activity of this aminotransferase thus remains to be discovered. Finally, *CaBan3p* is likely involved in the kynurenine pathway of tryptophan conversion into NAD<sup>+</sup> but does not take part in the biosynthesis of any proteinogenic amino acids. None of the aminotransferases *CaAro8p*, *CaAro9p*, *CaYer152Cp*, or *CaBan3p* is solely responsible for the catabolism of a single particular proteinogenic amino acid or for its biosynthesis. Having a broad set of catabolic pathways for the specific nutrient sources in the host is clearly advantageous for pathogenic fungi (Brock, 2009). The results of our studies confirm the versatility of *C. albicans* aminotransferases, which constitute a basis for the nutritional flexibility of this human pathogenic fungus. Based on our data, we suggest a model of *CaAro8p*, *CaAro9p* and *CaYer152Cp* role in *C. albicans* L-Lys, L-His and aromatic amino acids metabolism (Fig. 8).

## Acknowledgements

We are very grateful to Professor J. Morschhäuser and Dr. B. Ramírez-Zavala from the Institute for Molecular Infection Biology, University of Würzburg, Germany for their hospitality provided to one of us (KR) during her visit to University of Würzburg and for the invaluable help with mutant strains construction.

Financial support of these studies by the National Science Centre, Poland (grant No. 2015/17/B/NZ6/04248 awarded to Iwona Gabriel) and Polish Ministry of Science and Higher Education (Project No: POKL.04.01.01-00-368/09; travel grant awarded to Kamila Rząd) is gratefully acknowledged.

## References

- Aoki, T., 1996. Antifungal azoxybacillin exhibits activity by inhibiting gene expression of sulfite reductase. *Antimicrob. Agents Chemother.* 40, 127–132.
- Brock, M., 2009. Fungal metabolism in host niches. *Curr. Opin. Microbiol.* 12, 371–376.
- Brunke, S., Seider, K., Almeida, R.S., Heyken, A., Fleck, C.B., Brock, M., Barz, D., Rupp, S., Hube, B., 2010. *Candida glabrata* tryptophan-based pigment production via the Ehrlich pathway. *Mol. Microbiol.* 76, 25–47.
- Brunke, S., Seider, K., Richter, M.E., Bremer-Streck, S., Ramachandra, S., Kiehnopf, M., Brock, M., Hube, B., 2014. Histidine degradation via an aminotransferase increases the nutritional flexibility of *Candida glabrata*. *Eukaryot. Cell* 13, 758–765.
- Bulfer, S.L., Brunzelle, J.S., Trievel, R.C., 2013. Crystal structure of *Saccharomyces cerevisiae* Aro8, a putative  $\alpha$ -aminoacidipate aminotransferase. *Protein Sci.* 22, 1417–1424.
- Gabriel, I., Kur, K., Laforce-Nesbitt, S.S., Pulickal, A.S., Bliss, J.M., Milewski, S., 2014. Phenotypic consequences of *LYS4* gene disruption in *Candida albicans*. *Yeast* 31, 299–308.
- Gabriel, I., Milewski, S., 2016. Characterization of recombinant homocitrate synthase from *Candida albicans*. *Protein Expr. Purif.* 125, 7–18.
- Gabriel, I., Vetter, N.D., Palmer, D.R., Milewska, M.J., Wojciechowski, M., Milewski, S., 2013. Homoisocitrate dehydrogenase from *Candida albicans*: properties, inhibition, and targeting by an antifungal pro-drug. *FEMS Yeast Res.* 13, 143–155.
- Gilardi, G.L., 1965. Nutrition of systemic and subcutaneous pathogenic fungi. *Bacteriol. Rev.* 29, 406–424.
- Gillum, A.M., Tsay, E.Y., Kirsch, D.R., 1984. Isolation of the *Candida albicans* gene for orotidine-5'-phosphate decarboxylase by complementation of *S. cerevisiae* *ura3* and *E. coli* *pyrF* mutations. *Mol. Gen. Genet.* 198, 179–182.
- Groll, A.H., Lumb, J., 2012. New developments in invasive fungal disease. *Fut. Microbiol.* 7, 179–184.
- Hazelwood, L.A., Daran, J.M., van Maris, A.J., Pronk, J.T., Dickinson, J.R., 2008. The Ehrlich pathway for fusel alcohol production: a century of research on *Saccharomyces cerevisiae* metabolism. *Appl. Environ. Microbiol.* 74, 2259–2266.
- Hébert, A., Casaregola, S., Beckerich, J.-M., 2011. Biodiversity in sulfur metabolism in hemiascomycetous yeasts. *FEMS Yeast Res.* 11, 366–378.
- Iraqi, I., Vissers, S., Cartiaux, M., Urrestarazu, A., 1998. Characterisation of *Saccharomyces cerevisiae* ARO8 and ARO9 genes encoding aromatic aminotransferases I and II reveals a new aminotransferase subfamily. *Mol. Gen. Genet.* 257, 238–248.
- Jastrzębowska, K., Gabriel, I., 2015. Inhibitors of amino acids biosynthesis as antifungal agents. *Amino Acids* 47, 227–249.
- King, R.D., Rowland, J., Oliver, S.G., Young, M., Aubrey, W., Byrne, E., Liakata, M., Markham, M., Pir, P., Soldatova, L.N., Sparkes, A., Whelan, K.E., Clare, A., 2009. The automation of science. *Science* 323, 85–89.
- Kingsbury, J.M., Yang, Z., Ganous, T.M., Cox, G.M., McCusker, J.H., 2004. Novel chimeric spermidine synthase-saccharopine dehydrogenase gene (SPE3-LYS9) in the human pathogen *Cryptococcus neoformans*. *Eukaryot. Cell* 3, 752–763.
- Kinzel, J.J., Winston, M.K., Bhattacharjee, J.K., 1983. Role of L-lysine- $\alpha$ -ketoglutarate aminotransferase in catabolism of lysine as a nitrogen source for *Rhodotorula glutinis*. *J. Bacteriol.* 155, 417–419.
- Kur, K., Gabriel, I., Morschhäuser, J., Barchiesi, F., Spreghini, E., Milewski, S., 2010. Disruption of homocitrate synthase genes in *Candida albicans* affects growth but not virulence. *Mycopathologia* 170, 397–402.
- Large, P., 1986. Degradation of organic nitrogen compounds by yeasts. *Yeast* 2, 1–34.
- Li, Y.F., Bao, W.G., 2007. Why do some yeast species require niacin for growth? Different modes of NAD synthesis. *FEMS Yeast Res.* 7, 657–664.
- López-Martínez, R., 2010. Candidosis, a new challenge. *Clin. Dermatol.* 28, 178–184.
- Martínez, P., Ljungdahl, P.O., 2005. Divergence of Stp1 and Stp2 transcription factors in *Candida albicans* places virulence factors required for proper nutrient acquisition under amino acid control. *Mol. Cell. Biol.* 25, 9435–9446.
- Miceli, M.H., Díaz, J.A., Lee, S.A., 2011. Emerging opportunistic yeast infections. *Lancet Infect. Dis.* 11, 142–151.
- Milewska, M.J., Prokop, M., Gabriel, I., Wojciechowski, M., Milewski, S., 2012. Antifungal activity of homoaconitate and homoisocitrate analogs. *Molecules* 17, 14022–14036.
- Panozzo, C., Nawara, M., Suski, C., Kucharczyk, R., Skoneczny, M., Becam, A.M., Rytka, J., Herbert, C.J., 2002. Aerobic and anaerobic NAD<sup>+</sup> metabolism in *Saccharomyces cerevisiae*. *FEBS Lett.* 517, 97–102.
- Pirkov, I., Norbeck, J., Gustafsson, L., Albers, E., 2008. A complete inventory of all enzymes in the eukaryotic methionine salvage pathway. *FEBS J.* 275, 4111–4120.
- Preuss, J., Hort, W., Lang, S., Netsch, A., Rahlfs, S., Lochnit, G., Jortzik, E., Becker, K., Maysr, P.A., 2013. Characterization of tryptophan aminotransferase 1 of *Malassezia furfur*, the key enzyme in the production of indolic compounds by *M. furfur*. *Exp. Dermatol.* 22, 736–741.
- Priest, S.J., Lorenz, M.C., 2015. Characterization of virulence-related phenotypes in *Candida* species of the CUG clade. *Eukaryot. Cell* 14, 931–940.
- Reuss, O., Vik, A., Kolter, R., Morschhäuser, J., 2004. The SAT1 flipper, an optimized tool for gene disruption in *Candida albicans*. *Gene* 341, 119–127.
- Rząd, K., Gabriel, I., 2015. Characterization of two aminotransferases from *Candida albicans*. *Acta Biochim. Pol.* 62, 903–912.
- Sasse, C., Schillig, R., Dierolf, F., Weyler, M., Schneider, S., Mogavero, S., Rogers, D.P., Morschhäuser, J., 2011. The transcription factor Ndt80 does not contribute to Mrr1-, Tac1-, and Upc2-mediated fluconazole resistance in *Candida albicans*. *PLoS ONE* 9, e25623.
- Schöbel, F., Jacobsen, I.D., Brock, M., 2010. Evaluation of lysine biosynthesis as an antifungal drug target: biochemical characterization of *Aspergillus fumigatus* homocitrate synthase and virulence studies. *Eukaryot. Cell* 9, 878–893.
- Tan, I.K., Gajra, B., 2006. Plasma and urine amino acid profiles in a healthy adult population of Singapore. *Ann. Acad. Med. Singap.* 35, 468–475.
- Urrestarazu, A., Vissers, S., Iraqui, I., Grenson, M., 1998. Phenylalanine- and tyrosine-auxotrophic mutants of *Saccharomyces cerevisiae* impaired in transamination. *Mol. Gen. Genet.* 257, 230–237.
- Wallach, D.P., 1961. Studies on the GABA pathway II. The lack of effect of pyridoxal phosphate on GABA-KGA transaminase inhibition induced by amino-oxyacetic acid. *Biochem. Pharmacol.* 8, 328–331.
- Wogulis, M., Chew, E.R., Donohoue, P.D., Wilson, D.K., 2008. Identification of formyl kynurenine formamidase and kynurenine aminotransferase from *Saccharomyces cerevisiae* using crystallographic, bioinformatic and biochemical evidence. *Biochemistry* 47, 1608–1621.
- Xu, H., Andi, B., Qian, J., West, A.H., Cook, P.F., 2006. The  $\alpha$ -aminoacidipate pathway for lysine biosynthesis in fungi. *Cell Biochem. Biophys.* 46, 43–64.
- Yamaki, H., Yamaguchi, M., Imamura, H., Suzuki, H., Nishimura, T., Saito, H., Yamaguchi, H., 1990. The mechanism of antifungal action of (S)-2-amino-4-oxo-5-hydroxypentanoic acid, RI-331: the inhibition of homoserine dehydrogenase in *Saccharomyces cerevisiae*. *Biochem. Biophys. Res. Commun.* 168, 837–843.
- Ye, Z.H., Garrad, R.C., Winston, M.K., Bhattacharjee, J.K., 1991. Use of alpha-aminoacidipate and lysine as sole nitrogen source by *Schizosaccharomyces pombe* and selected pathogenic fungi. *J. Basic Microbiol.* 31, 149–156.